

Modern Algebra 2

Grant Talbert

Spring 2025



College of Arts and Sciences
Department of Mathematics and Statistics
MA 542

Abstract

idk tim kohl

Contents

1	Rings
---	-------

2

Chapter 1

Rings

Lecture 1: Syllabus Day

Wed 22 Jan 2025 13:22

Definition 1.1: Ring

Let R be a set together with two binary operations

$$+, \bullet : R \times R \rightarrow R$$

such that $\forall a, b, c \in R$,

- $a + b = b + a$;
- $(a + b) + c = a + (b + c)$;
- $\exists 0 \in R$ such that $a + 0 = a$ for all $a \in R$;
- $\forall a \in R, \exists b \in R$ such that $a + b = 0$;
- $(ab)c = a(bc)$;
- $a(b + c) = ab + ac$ and $(a + b)c = ac + bc$.

We then call $(R, +, \cdot)$ a ring.

Note that $(R, +)$ is an abelian group. There then clearly exists a forgetful functor $\mathbf{Ring} \rightarrow \mathbf{Grp}$ that maps $(R, +, \cdot) \mapsto (R, +)$. Examples of rings include $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ under the standard definitions of addition and multiplication. However, these rings are all commutative, and many are in fact fields. There is more to ring theory than commutative rings. Oh also $\mathbb{Z}_n$ is a ring $\forall n \in \mathbb{N}$.

Note that since left multiplication is a faithful group action on itself, every row/column of the cayley table in some group has each number appear exactly once.

We take the convention that rings need not have 1. Homomorphisms thus need not preserve 1, except for epimorphisms. An example of a ring without 1 is $2\mathbb{Z}$. Also note that rings need not be commutative; for example, $M_2(\mathbb{R})$. We generally divide rings into commutative rings and noncommutative rings. the category of commutative rings is denoted $\mathbf{CRing}$.

Definition 1.2: Unity

Let R be a ring with some $1 \in R$ such that $\forall r \in R, 1r = r1 = r$. We say 1 is the unity element of R .

Definition 1.3: Polynomial Ring

Let R be a ring. Define $R[x]$ as the polynomial ring

$$R[x] := \left\{ \sum_i a_i x^i \mid a_i \in R \right\}$$

of *finite* linear combinations in the indeterminate x .

Lecture 2: hi

Fri 24 Jan 2025 13:27

Let R be a ring without unity. Define $S = R \times \mathbb{Z}$ or $S = r \times \mathbb{Z}_{\text{char}(R)}$. Define multiplication by

$$(r, n)(s, m) = (rs + ns + mr, nm).$$

Then $(0, 1)$ is a unity and there exists an inclusion $R \hookrightarrow S$ which is also a ring homomorphism by the map $r \mapsto (r, 0)$.

The quaternions $\mathbb{H}$ are defined as the set of linear combinations of the set $\{1, i, j, k\}$ wherein multiplication satisfies $ijk = -1$, $i^2 = -1$, $j^2 = -1$, and $k^2 = -1$. They are intended as an extension of the complex numbers $\mathbb{C}$, in a similar way to how the complex numbers extend $\mathbb{R}$. However, this definition of multiplication necessitates $ij = -ji$, meaning $\mathbb{H}$ is a noncommutative ring. Furthermore, it can be shown (with some difficulty) that all elements of $\mathbb{H}$ have multiplicative inverses. So $\mathbb{H}$ is a noncommutative ring with unity with multiplicative inverses. We call this a division ring.

Definition 1.4: Division Ring

Let R be a ring with unity, such that $\forall r \in R, \exists r^{-1} \in R$ such that $rr^{-1} = r^{-1}r = 1_R$. Then R is a division ring.

Going back down in dimension to $\mathbb{C}$ gives more structure. Elements in $\mathbb{C}$ do, in fact, commute under multiplication; this makes $\mathbb{C}$ into a field.

Definition 1.5: Field

Let k be a commutative ring with unity, such that for all $r \in k$, there exists $r^{-1} \in k$ such that $rr^{-1} = 1_k$. Then k is called a field.

Definition 1.6: Direct Product

Let R, S be rings. Define the direct product $R \times S$ as the cartesian product of the sets R, S together with coordinate-wise addition and multiplication. Moreover, for any set of rings $R_1, \dots, R_n$, define the product $R_1 \times \dots \times R_n$, which is defined in the same way.

The direct product satisfies the universal property for products in **Ring**, but there are more multiplicative structures that can be endowed to the underlying abelian group $R_1 \oplus \dots \oplus R_n$, which we may or may not see when we do polynomial rings. We may write $\oplus$ instead of $\times$ in some commutative cases.

Proposition 1.1

Let R be a ring, and let $a, b, c \in R$. Then

- $a0 = 0a = 0$;
- $a(-b) = (-a)b = -ab$;
- $(-a)(-b) = ab$;
- $a(b - c) = ab - ac$ and $(a - b)c = ac - bc$;
- $(-1)a = -a$;
- $(-1)(-1) = 1$.

Proof. (1) Many of these are trivial, so I will only prove a few. Let $a \in R$. Then

$$0a = (0 + 0)a = 0a + 0a \implies 0a - 0a = 0a - 0a - 0a \implies 0 = 0a.$$

(3) Now let $a, b \in R$ and note that $(-a + a)(-b) = 0(-b) = 0$, and thus

$$(-a)(-b) + a(-b) = 0.$$

By property 2,

$$(-a)(-b) + a(-b) = (-a)(-b) - (ab).$$

Therefore, $(-a)(-b) = ab$.

Statement holds for $n = 1, 2$ trivially. Let n even and suppose the statement holds $\forall m \leq n$. We have

$$x^{n+2} = \left(x^{\frac{n}{2}+1}\right)^2 = 0 \iff x^{\frac{n}{2}+1} = 0.$$

Therefore, $x^{n+2} = 0$ iff $x = 0$. Now we have immediately

$$x(x^{n+1}) = x^{n+2}.$$

If $x \neq 0$ but $x^{n+1} = 0$, then we have $x^{n+2} = 0$ for $x \neq 0$, contradiction. ■