

Higher Topos Theory - Self Study

Grant Talbert

Summer 2025-Winter 2025

Contents

1	An Overview of Higher Category Theory	1
1.1	Foundations of Higher Category Theory	1
1.1.1	Goals and Obstacles	1
1.1.2	∞ -Categories	2
1.1.3	Equivalences of Topological Categories	12

Abstract

I am currently reading Higher Topos Theory to understand the theory of quasi-categories and basic ∞ -category theory. My goal is to understand the generalizations of basic 1-categorical concepts to the ∞ -categorical setting, such as (co)limits, adjunctions, and Yoneda's lemma. I'm also interested in more advanced topics in higher category theory, most of which require a basic understanding of the theory of ∞ -categories as a prerequisite.

I began reading this text on August 8th, 2025, at 11:06 PM, as an independent project. I am hoping to be able to find a professor who is willing to support my readings, but I am unsure that will happen. Nevertheless, I intend to gain a working understanding of ∞ -categories over the next few months, and a more advanced understanding come the spring semester. Since this text is quite advanced and is not meant as a textbook, but rather as a reference, I will be taking extremely detailed and extensive notes on the text. I will also be referencing supplementary resources throughout my reading, so I have decided it worthwhile to maintain a bibliography for these notes. Any notes not credited to a bibliography source are either from [Lur09], or directly from me.

Chapter 1

An Overview of Higher Category Theory

The first chapter introduces quasi-categories as models for ∞ -categories, and then develops ∞ -categorical analogues of standard 1-categorical notions.

1.1 Foundations of Higher Category Theory

Terminology 1.1.0.1. Throughout this paper, the term *space* is short for a topological space, and almost always a compactly generated topological space. The *homotopy type* of a topological space is a choice of a weak homotopy equivalent topological space, or equivalently the collection of weak homotopy equivalent spaces.

1.1.1 Goals and Obstacles

The theory of ∞ -categories finds natural motivation in the fundamental n -groupoids of a topological space X . Define $\Pi_n(X)$ to take points $x \in X$ as objects, paths from x to y as 1-morphisms, homotopies of paths as 2-morphisms, and so on until n . Since vertical composition of homotopies is only associative up to higher homotopy, we are forced to take n -morphisms to be *homotopy classes*, rather than simply homotopies. Of course $\Pi_n(X)$ is a groupoid because all paths and homotopies admit inverses.

One might hope that we can fix this problem by considering an ∞ -groupoid, or an $(\infty, 0)$ -category, which would not take homotopies as n -morphisms for all $n \geq 2$. Interestingly, the ∞ -groupoid $\Pi_\infty(X)$ of a space X encodes the homotopy type of X , meaning the study of ∞ -groupoids simplifies to the study of spaces in classical homotopy theory, and we have many ways of describing homotopy types of spaces. It is also a commonly accepted principle of higher category theory that all $(\infty, 0)$ -categories are of the form $\Pi_\infty(X)$ for some space X . This principle is profound, since it gives us a place to begin our work in higher category theory.

In general, an n -category should be a category enriched over $(n - 1)$ -categories. However, for finite n , this process produces the notion of a *strict* n -category, and n -category theory is more naturally viewed via the theory of *weak* n -categories. These issues make the theory of n -categories fairly difficult to work with, but they vanish when one passes to ∞ -categories. An (∞, n) -category can indeed be viewed as a category enriched over $(\infty, n - 1)$ -categories. This discussion suggests a definition (or more precisely, a blueprint) for ∞ -categories. An ∞ -category $\mathcal{C}$ should consist of objects $\text{Obj}(\mathcal{C})$, and for each $x, y \in \mathcal{C}$, a ∞ -groupoid $\text{Map}_{\mathcal{C}}(x, y)$ with an associative composition law. The theory of ∞ -categories does not care whether this law be strictly associative or only associative up to coherent homotopy, since any ∞ -category with coherently associative multiplication is equivalent to one with strictly associative multiplication.

Definition 1.1.1.1. A *topological category* is a category enriched over $\mathcal{CG}$, the category of compactly generated spaces. The category of topological spaces will be denoted by $\mathcal{Cat}_{\text{top}}$. $\heartsuit$

We recall from our knowledge of enriched category theory that this implies a topological category $\mathcal{C}$ has topological spaces $\text{Map}_{\mathcal{C}}(x, y)$ for each pair $x, y \in \mathcal{C}$, with an associative composition law

$$\text{Map}_{\mathcal{C}}(y, z) \times \text{Map}_{\mathcal{C}}(x, y) \rightarrow \text{Map}_{\mathcal{C}}(x, z),$$

where the product is computed in $\mathcal{CG}$.

Remark 1.1.1.2. As is often the case, working in $\mathcal{CG}$ is done largely as a technical point which can be ignored.

1.1.2 ∞ -Categories

Definition 1.1.1.1 is the most obvious definition of an ∞ -category, but it is also one of the hardest to actually work with. Our use of topological categories in this book will be more as a “benchmark” of sorts. Any model for ∞ -categories which is equivalent to that of topological categories must necessarily be a good one. Many better models exist, and in this text we choose to use the theory of *weak Kan complexes*. These were first introduced by Boardman and Vogt in [BV73], and were more recently and more extensively studied by André Joyal in [Joy08a, Joy08b], who instead called them *quasi-categories*, which the modern literature now uses. However, Lurie’s book almost exclusively uses quasi-categories as our model for ∞ -categories, so for the purposes of these notes we will use the terms “quasi-category” and “ ∞ -category” more or less interchangeably.

Quasi-categories are built atop the theory of *simplicial sets*, and any readers of HTT (or these notes) should have at least an elementary understanding of the subject. I recommend my own exposition [Tal25], which builds the abstract theory of simplicial sets without forsaking geometric intuition, but also without dwelling excessively on the topological implications of the theory. We also recommend [Rie11, Fri23]. The former should prepare the reader for much of the theory in this paper, the rest of which can be picked up along the way, although we feel Riehl strays too far from the geometric intuition of simplices. Conversely, Friedman’s introduction is well-liked for the geometric intuition it provides, but we feel it focuses too much on topological aspects of the theory to be the best preparation for the material covered in HTT.

The theory of simplicial sets is of importance due to the homotopic aspects of the theory. Recall from our discussion in section 1.1.1 that the ∞ -groupoid $\Pi_{\infty}(X)$ of a space X captures the homotopy type of X . We also know from [Qui67] that since $|-| \dashv \text{Sing}$ is a Quillen equivalence, where Sing is the singular complex functor. This means in particular that the unit and counit of the adjunction are natural weak homotopy equivalences, meaning spaces are determined by their singular sets up to homotopy type. Thus, if we are only interested in the study of spaces up to weak homotopy equivalence, we might as well work in the category of simplicial sets. We might also think that ∞ -groupoids should be able to be modeled by singular sets, since they both encode the homotopy type of a space. Indeed this is possible, but we prefer to work slightly more generally. Singular sets satisfy the following important property.

Definition 1.1.2.1. Let K be a simplicial set. We say K is a *Kan complex* if, for all $n \geq 0$ and for

all $0 \leq i \leq n$, any diagram of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow \text{dotted} & \\ \Delta^n & & \end{array}$$

admits a (not necessarily unique) dotted arrow as indicated, rendering the diagram commutative. $\heartsuit$

Recall here that $\Lambda_i^n \subseteq \Delta^n$ represents the i th horn of the standard n -simplex; we defer the reader to [Tal25, §3.4] for a rigorous definition.

Remark 1.1.2.2. The singular complex of a topological space is a Kan complex. Showing this first involves showing that $|\Lambda_i^n|$ is a retract of $|\Delta^n|$ in $\mathcal{J}\text{op}$. There also exist combinatorial approaches to extracting homotopy groups from any Kan complex K , which agree with the homotopy groups on $|K|$. Thus, Kan complexes behave like the homotopy type of a space.

Simplicial sets, and in particular Kan complexes, thus should form a model for ∞ -groupoids, which we note are an important special case of ∞ -categories. Indeed, Kan complexes are a common model for ∞ -groupoids, which we will explore later. However, another special case of ∞ -categories that our definition must reflect is the notion of a basic 1-category. If an ∞ -category $\mathcal{C}$ has no nontrivial n -morphisms for $n > 1$, then clearly $\mathcal{C}$ can be regarded as a 1-category. Luckily, simplicial sets are also closely related to category theory. Recall the nerve $N(\mathcal{C})$ of a category is a simplicial set whose n -simplices are functors $[n] \rightarrow \mathcal{C}$, which carry the data of an n -tuple of composable morphisms.

Remark 1.1.2.3. The nerve encodes the full information of a category, and we can in fact recover the initial category $\mathcal{C}$ from the nerve $N(\mathcal{C})$ with the following list of observations.

1. The 0-simplices $N(\mathcal{C})_0$ are the objects, so we define $\text{Obj}(\mathcal{C}) = N(\mathcal{C})_0$.
2. The 1-simplices $N(\mathcal{C})_1$ are simply morphisms, so we define $\text{Mor}(\mathcal{C}) = N(\mathcal{C})_1$.
3. Given an object $x \in \mathcal{C}$, the degeneracy $s_0 : N(\mathcal{C})_0 \rightarrow N(\mathcal{C})_1$ maps to the identity $x \mapsto 1_x$. Thus, the identity function $x \mapsto 1_x$ is simply the degeneracy $s_0 = 1_\bullet$.
4. Given a morphism $f \in N(\mathcal{C})_1$, the 0th face $d_0 : N(\mathcal{C})_1 \rightarrow N(\mathcal{C})_0$ maps to the codomain $f \mapsto \text{cod } f$, and similarly the first face $d_1 : N(\mathcal{C})_1 \rightarrow N(\mathcal{C})_0$ maps to the domain $f \mapsto \text{dom } f$. We may thus define $\text{dom}, \text{cod} : \text{Mor}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{C})$ as the faces $\text{dom} = d_1$ and $\text{cod} = d_0$.
5. A composable pair (g, f) is a 2-simplex $(g, f) \in N(\mathcal{C})_2$. The 1st face $d_1 : N(\mathcal{C})_2 \rightarrow N(\mathcal{C})_1$ maps composable pairs to their composite $(g, f) \mapsto gf$. Thus, composition is simply the 1st face $d_1 = \circ$.
6. Composites of larger tuples of maps arise from higher degeneracies, and associativity follows from the simplicial identities. The unit law for identities under composition also follows from simplicial identities.

A more explicit description of remark 1.1.2.3 (5,6) will be helpful later. A morphism from x to y is defined as a 1-simplex $f \in N(\mathcal{C})_1$ with $d_1(f) = x$ and $d_0(f) = y$. Given two morphisms $x \xrightarrow{f} y \xrightarrow{g} z$, the composite $g \circ f$ can be characterized by noting there exists a 2-simplex $(g, f) = \sigma \in N(\mathcal{C})_2$ such that

1. $d_0(\sigma) = g$;
2. $d_1(\sigma) = gf$;

3. $d_2(\sigma) = f$.

We say σ witnesses gf as a composite of g and f .

There exists a characterization of simplicial sets that arise as nerves, which is more straightforward than the characterization of Kan complexes.

Notation 1.1.2.4. Let J be a linearly ordered set. Denote by Δ^J the representable functor $\text{Hom}(-, J)$ in the category of linearly ordered sets, restricted along the objects of $\mathbf{\Delta}$. Clearly, $\Delta^{[n]} = \Delta^n$, so this is simply a mild generalization of notation we already had.

Proposition 1.1.2.5. Let K be a simplicial set. The following conditions are equivalent.

1. There exists a small category $\mathcal{C}$ and an isomorphism $K \cong N(\mathcal{C})$.
2. For each $0 < i < n$, and each diagram of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & K \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

there is a unique dashed arrow as indicated, rendering the diagram commutative.

Proof. We mostly follow Lurie's proof in [Lur09, Prop. 1.1.2.2], but we take a different route for showing agreement for (1), and Lurie assumes a slightly more advanced understanding of simplicial sets than I currently have, so at some points in this proof we may prove extra facts. For example, we start by proving that for $0 < i < n$, the vertices of the i th horn are $(\Lambda_i^n)_0 = [n]$. Note that for i to exist, we must have $n \geq 2$. For $n = 2$, there exist at two coface maps $d^j : \Delta^1 \rightarrow \Delta^2$ with $j \neq i$. 0-simplices of Δ^1 are arrows $[0] \rightarrow [1]$, which correspond to elements of $[1]$. More precisely, arrows are uniquely determined by their image. Thus, 0-simplices in the face $\partial_j \Delta_0^n = \text{im } d_0^j$ will also be uniquely determined by their image, so it suffices to show each $x \in [2]$ lands in the image of some coface d^j with $j \neq i$. This is immediate: since d^j only skips j , and since another coface d^k , $k \neq i$ exists which doesn't skip j , every element shows up. Inducting on this same logic proves the statement for $n \geq 2$, and vacuously for all n .

Now we show for $0 < k \leq n$ and $n \geq 2$, there is a simplicial subset $\Delta^{\{k-1, k\}} \subseteq \Lambda_i^n$. Recall that m -simplices f in the horn are composites $[m] \xrightarrow{f} [n-1] \xrightarrow{d^{j \neq i}} [n]$. Let $g : [m] \rightarrow \{k-1, k\} \subseteq [n]$ be an m -simplex in $\Delta^{k-1, k}$. If $k < n$, then $k, k-1 \in [n-1]$. Write g' for the map with codomain viewed as a subset $\{k-1, k\} \subseteq [n-1]$. Since $i \neq n$, we have that $\text{im } d^n \subseteq \Lambda_i^n$. Thus, $d^n g' = g \in \text{im } d_m^n$, and thus is an m -simplex in the horn. If $k = n$, then since $n \geq 2$, we have $k-1 > 0$. Since $i \neq 0$, we do the same argument.

If $n < 2$ then the statement is true vacuously, so let $n \geq 2$. Let K be the nerve of a small category $\mathcal{C}$. Let $f_0 : \Lambda_i^n \rightarrow K$ with $0 < i < n$, and let $x_k \in \mathcal{C}$ denote the image of the vertex $k \in (\Lambda_i^n)_0 = [n]$ under f_0 . Choose some $0 < k \leq n$, and consider the restriction $f_0|_{\Delta^{\{k-1, k\}}}$. Note that the 1-simplices of $\Delta^{\{k-1, k\}}$ are

$$\Delta_1^{\{k-1, k\}} = \left\{ (0, 1) \mapsto k, (0, 1) \mapsto k-1, \begin{matrix} 0 \mapsto k-1 \\ 1 \mapsto k \end{matrix} \right\}.$$

The first two are easily seen to be the degenerates $s_0(k), s_0(k-1)$, respectively. Since f_0 is a morphism of simplicial sets, $f_0 s_0(k) = s_0 f_0(k) = s_0(x_k) = 1_{x_k}$. This is uninteresting, so let g_k denote

the unique nondegenerate simplex. Note that the image under f is a 1-simplex in $N(\mathcal{C})$; that is a morphism $g_k : c \rightarrow d$. We want to show that with notation as above, $c = x_{k-1}$ and $d = x_k$. Indeed, note that for 1-simplices in the nerve, the first face is simply *domain* and the 0th face is *codomain*. Since f_0 commutes with faces, we have

$$\text{dom } g_k = d_1 f_0(g_k) = f_0 d_1(g_k) = f_0(g_k \circ d^1) = f_0(0 \mapsto 0 \mapsto k-1) = x_{k-1},$$

since $0 \mapsto 0 \mapsto k-1 = g_k \circ d^1$ is a 0-simplex, which is uniquely determined (and identified with) its image. The same follows for domain.

We thus have for each $0 < k \leq n$, there is an arrow $g_k : x_{k-1} \rightarrow x_k$, allowing us to form a composable chain

$$x_0 \xrightarrow{g_1} x_1 \xrightarrow{g_2} \cdots \xrightarrow{g_n} x_n.$$

This is an n -simplex in Δ^n . We can similarly construct m -simplices for $m \leq n$ by requiring $k \leq m$. We claim that this relation allows us to define a mapping $f : \Delta^n \rightarrow K$.

To see how this follows, we invoke the Yoneda lemma to recall that there is an isomorphism $\text{Set}^{\Delta^{\text{op}}}(\Delta^n, K) \cong K_n$ given by evaluating the n th component of natural transformations at the identity. In particular, Yoneda says a morphism $f : \Delta^n \rightarrow K$ is determined by $f_n(1_{[n]})$. We choose $f_n(1_{[n]})$ to be the n -simplex $(g_1, \dots, g_n)$ of $N(\mathcal{C})$ considered above. The remainder of the proof will focus on showing this definition indeed agrees with Λ_i^n . For now, we note that f is manifestly unique, since we obtained a unique composite constructed in this way, so any other map $\Delta^n \rightarrow K$ which restricts along the horn to f_0 must necessarily correspond to the same chain of morphisms.

We now begin to show $f|_{\Lambda_i^n} = f_0$. First, we show they agree on cofaces. Let $d^j : [n-1] \rightarrow [n]$ be an $(n-1)$ -simplex in Λ_i^n . Then since f is a morphism of simplicial sets and commutes with faces, $d^j = 1_{[n]} d^j = d_j(1_{[n]})$ implies that $f(d^j) = d_j f(1_{[n]})$. For simplicity, we take d^j to be an inner face $0 < j < n$, and simply note the argument for outer faces is the same. From our knowledge of the nerve, we have that

$$\begin{aligned} f(d^j) &= f d_j(1_{[n]}) = d_j f(1_{[n]}) = d_j \left(x_0 \xrightarrow{g_1} \cdots \xrightarrow{g_n} x_n \right) \\ &= x_0 \xrightarrow{g_1} \cdots \xrightarrow{g_{j-1}} x_{j-1} \xrightarrow{g_{j+1} g_j} x_{j+1} \xrightarrow{g_{j+1}} \cdots \xrightarrow{g_n} x_n \end{aligned}$$

It suffices to show $f_0(d^j)$ agrees with this composite. One can do this by showing each object and morphism align with that of $f(d^j)$, which involves an induction argument that is extremely lengthy due to needing to show many cases, but is in general very simple. We trust the reader to be able to do this argument, and we also trust that the reader knows I can do the argument too and just don't want to, so we prefer not to spend 2 pages on it and instead continue with the proof.

Now, we show this agreement on cofaces implies f and f_0 agree in general. We note that every m -simplex $h \in \Delta_m^n$ is a morphism $h : [m] \rightarrow [n]$ in Δ , which admits a factorization into cofaces and codegeneracies $h = d^{j_1} \circ \cdots \circ d^{j_\ell} \circ s^{p_1} \circ \cdots \circ s^{p_q}$. Since faces and degeneracies in Δ^n are given by precomposing by the corresponding cofaces and codegeneracies in Δ , we have

$$h = d^{j_1} \circ \cdots \circ s^{p_q} = 1_{[n]} \circ d^{j_1} \circ \cdots \circ s^{p_q} = (s_{p_q} \circ \cdots \circ d_{j_1})(1_{[n]}).$$

Since f is a morphism of simplicial sets, we have $f s_j = s_j f$, and likewise for faces. Induction gives

$$f(h) = (f \circ s_{p_q} \circ \cdots \circ d_{j_1})(1_{[n]}) = (s_{p_q} \circ \cdots \circ d_{j_1})(f(1_{[n]})) = (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1}(1_{[n]}))).$$

Now we want to show f agrees with f_0 on m -simplices of Λ_i^n . Since $h \in (\Lambda_i^n)_m$, there is some m -simplex h' such that $h = d^j h'$ for $j \neq i$, so by uniqueness we have that $d^{j-1} \neq d^i$ in the epi-mono factorization of h . Since f and f_0 agree on cofaces,

$$\begin{aligned} f(h) &= (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1}(1_{[n]}))) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1})) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f_0(d_{j_1})) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f_0(d_{j_1}(1_{[n]}))) \\ &\stackrel{!}{=} f_0(h), \end{aligned}$$

where the equality marked by $!$ follows since f_0 is a morphism of simplicial sets and commutes with faces and degeneracies. Thus, the statement $(1 \implies 2)$ is proven.

Now for the converse. Let K satisfy (2), and define a category $\mathcal{C}$ as follows.

1. Objects are vertices: $\text{Obj}(\mathcal{C}) = K_0$;
2. Morphisms are edges: $\text{Mor}(\mathcal{C}) = K_1$. Domain and codomain are faces: $\text{dom} = d_1$ and $\text{cod} = d_0$;
3. Identity $1_\bullet : \text{Obj}(\mathcal{C}) \rightarrow \text{Mor}(\mathcal{C})$ is the degeneracy $1_\bullet = s_0$;
4. Let $f : x \rightarrow y$ and $g : y \rightarrow z$ in $\mathcal{C}$. We want to define a morphism $\sigma_0 : \Lambda_1^2 \rightarrow K$. Define the action on 0-simplices by $0 \mapsto x$, $1 \mapsto y$, and $2 \mapsto z$. Denote by g_k the nondegenerate simplex in $\Delta^{k-1,k}$ for $0 < k \leq 2$. Then define the map as $g_1 \mapsto f$ and $g_2 \mapsto g$. We will show later that this data determines a morphism $\sigma_0 : \Lambda_1^2 \rightarrow K$. By the horn filling condition, this extends to a map $\sigma : \Delta^2 \rightarrow K$. Let $d^1 : [1] \rightarrow [2]$ be the unique 1-simplex of Δ^2 with $d^1(0) = 0$ and $d^1(1) = 2$. Then define $g \circ f$ as $\sigma(d^1)$.

Of course, we need to show the data in (4) defines a morphism $\sigma_0 : \Lambda_1^2 \rightarrow K$. Recall that $\Lambda_1^2 = \partial_0 \Delta^2 \cup \partial_2 \Delta^2$. By universality of union, if we can define maps $\partial_i \Delta^2 \rightarrow K$ for $i = 0, 2$, such that these maps agree on $\partial_0 \Delta^2 \cap \partial_2 \Delta^2$, then they assemble into a map Λ_1^2 . We may thus construct a map in this way using only the data of (4). Recall that $\partial_i \Delta^n = \text{im}(d^i : \Delta^{n-1} \rightarrow \Delta^n)$, and in particular $d^i : \Delta^{n-1} \rightarrow \partial_i \Delta^n$ is an isomorphism by uniqueness of the epi-mono factorization in $\mathbf{\Delta}$, with a well-defined inverse $d^i \circ h \mapsto h$. Thus, if $\tau_0 : \Delta^{n-1} \rightarrow K$ is a morphism of simplicial sets, then there is a morphism of simplicial sets $\tau : \partial_i \Delta^n \rightarrow K$ given by $d^i \circ h \mapsto h \mapsto \tau_0(h)$, since this is simply composition of the inverse of the isomorphism with the map τ_0 . In particular, maps $\Delta^1 \rightarrow K$ induce maps $\partial_i \Delta^2 \rightarrow K$.

By the Yoneda lemma, maps $\tau'_0 : \Delta^1 \rightarrow K$ are specified by the image $\tau'_0(1_{[1]})$. Thus, the map $\tau_0 : \partial_i \Delta^2 \rightarrow K$ is fully specified by the image $\tau'_0(1_{[1]})$ in the aforementioned manner. Define

$$\alpha_0 : \Delta^1 \rightarrow K, \quad 1_{[1]} \mapsto f,$$

$$\beta_0 : \Delta^1 \rightarrow K, \quad 1_{[1]} \mapsto g.$$

These of course assemble into maps $\alpha : \partial_2 \Delta^2 \rightarrow K$ and $\beta : \partial_0 \Delta^2 \rightarrow K$. To show these maps agree on the intersection, it suffices to show they agree on the vertex $1 \in \partial_{0,2} \Delta_0^2$. Indeed, $1 \in \partial_0 \Delta_0^2$ is simply $d^0(0)$, and $1 \in \partial_2 \Delta_0^2$ is simply $d^2(1)$. Thus, they are mapped

$$\alpha : 1 = d^2(1) \mapsto 1 \mapsto \alpha_0(1), \quad \beta : 1 = d^0(0) \mapsto 0 \mapsto \beta_0(0).$$

We simply need to show these agree. Since β_0 and α_0 are morphisms of simplicial sets, they necessarily

commute with faces, meaning the diagrams

$$\begin{array}{ccc} \Delta_1^1 & \xrightarrow{\alpha_0} & K_1 \\ d_0 \downarrow & & \downarrow d_0 \\ \Delta_0^1 & \xrightarrow{\alpha_0} & K_0 \end{array} \quad \begin{array}{ccc} \Delta_1^1 & \xrightarrow{\beta_0} & K_1 \\ d_1 \downarrow & & \downarrow d_1 \\ \Delta_0^1 & \xrightarrow{\beta_0} & K_0 \end{array}$$

commute. Recall that $\alpha_0(1_{[1]}) = f$ and $\beta_0(1_{[1]}) = g$. Following the identity around both squares yields the equalities

$$\begin{array}{ccc} 1_{[1]} & \xrightarrow{\alpha_0} & f \\ d_0 \downarrow & & \downarrow d_0 \\ 1_{[1]} \circ d^0 = 1 & \xrightarrow{\alpha_0} & \alpha_0(1) = d_0(f) \end{array} \quad \begin{array}{ccc} 1_{[1]} & \xrightarrow{\beta_0} & g \\ d_1 \downarrow & & \downarrow d_1 \\ 1_{[1]} \circ d^1 = 0 & \xrightarrow{\beta_0} & \alpha_0(0) = d_1(g) \end{array}$$

Recall that in (2), we defined $d_1(g) = \text{dom } g$ and $d_0(f) = \text{cod } f$. In (4), we defined $\text{cod } f = y = \text{dom } g$, and thus the morphisms agree. We thus have an induced map $\sigma_0 = \alpha \cup \beta : \Lambda_1^2 \rightarrow K$, which is immediately guaranteed to be defined as claimed in (4).

Now that our definitions are indeed defined, we must show $\mathcal{C}$ defines a category, as claimed.

1. First, we show the identity is indeed an identity. Let $y \in \mathcal{C}$, and let $f : x \rightarrow y$ and $g : y \rightarrow z$ be morphisms in $\mathcal{C}$. The data $(f, 1_y)$ constitutes a horn $x \xrightarrow{f} y \xrightarrow{1} y$, which fills to a 2-simplex $\tau \in K_2$ and a morphism $\sigma : \Delta^2 \rightarrow K$ given by $1_{[2]} \mapsto \tau$, such that τ witnesses $1_y \circ f$ as the composite of 1_y and f , as described in (4). We show $1_y \circ f = f$. Recall $1_y \circ f = \sigma(d^1)$. We have

$$\begin{aligned} d_0(\tau) &= d_0\sigma(1_{[2]}) = \sigma d_0(1_{[2]}) = \sigma(d^0) = 1_y = s_0(y), \\ d_1(\tau) &= d_1\sigma(1_{[2]}) = \sigma d_1(1_{[2]}) = \sigma(d^1) = 1_y \circ f, \\ d_2(\tau) &= d_2\sigma(1_{[2]}) = \sigma d_2(1_{[2]}) = \sigma(d^2) = f. \end{aligned}$$

Recall that there is a simplicial identity $s_i d_i = s_i d_{i+1} = 1$. We have

$$s_1(f) = s_1 d_2(\tau) = \tau,$$

and thus τ is degenerate. We also have

$$1_y \circ f = d_1(\tau) = d_1 s_1(f) = f$$

by the same simplicial identity. The converse direction is the same argument, except we find that $s_0(g) = \tau$.

2. Now we show associativity. Let

$$w \xrightarrow{f} x \xrightarrow{g} y \xrightarrow{h} z$$

be a composable triple in $\mathcal{C}$. We want to show $h(gf) = (hg)f$. Define the following 2-simplices of K to be the images of $1_{[2]}$ under the relevant fillers:

$$\sigma_{gf} = \begin{cases} d_0(\sigma_{gf}) = g, \\ d_1(\sigma_{gf}) = gf, \\ d_2(\sigma_{gf}) = f \end{cases} \quad \sigma_{hg} = \begin{cases} d_0(\sigma_{hg}) = h, \\ d_1(\sigma_{hg}) = hg, \\ d_2(\sigma_{hg}) = g \end{cases}$$

$$\sigma_{h(gf)} = \begin{cases} d_0(\sigma_{h(gf)}) = h, \\ d_1(\sigma_{h(gf)}) = h(gf), \\ d_2(\sigma_{h(gf)}) = gf \end{cases}, \quad \sigma_{(hg)f} = \begin{cases} d_0(\sigma_{(hg)f}) = hg, \\ d_1(\sigma_{(hg)f}) = (hg)f, \\ d_2(\sigma_{(hg)f}) = f \end{cases}$$

By the same logic as earlier (and indeed, we will show this holds generally later with an inductive argument), there is a horn $\tau_0 : \Lambda_2^3 \rightarrow K$ and a filler τ defining the 3-simplex $\lambda = \tau(1_{[3]}) \in K_3$. This simplex has faces

$$\begin{aligned} d_0(\lambda) &= d_0\tau(1) = \tau(d^0) = \sigma_{hg}, \\ d_1(\lambda) &= d_1\tau(1) = \tau(d^1) = \sigma_{h(gf)}, \\ d_3(\lambda) &= d_3\tau(1) = \tau(d^3) = \sigma_{gf}. \end{aligned}$$

We first want to show $d_2(\lambda) = \sigma_{(hg)f}$. For now, we note that

$$\begin{aligned} d_0d_2(\lambda) &= d_1d_0(\lambda) = d_1(\sigma_{hg}) = hg, \\ d_2d_2(\lambda) &= d_2d_3(\lambda) = d_2(\sigma_{gf}) = f, \end{aligned}$$

by the simplicial identity $d_id_j = d_{j-1}d_i$ for $i < j$. This defines a horn, which admits a filler. Since $\sigma_{(hg)f}$ agrees with the horn but is a 2-simplex, it necessarily is a filler, and by (2) this filler must be unique. Thus, $d_2(\lambda) = \sigma_{(hg)f}$ is the unique filler. Finally, note that by the same simplicial identity,

$$(hg)f = d_1d_2(\lambda) = d_1d_1(\lambda) = h(gf).$$

Therefore, $\mathcal{C}$ is a category, as claimed.

Finally, we show the isomorphism $K \cong N(\mathcal{C})$. By construction, we have a map $\varphi : K \rightarrow N(\mathcal{C})$ given by mapping n -simplices to the n -fold composite they witness. Of course, this is a morphism of simplicial sets basically by construction, which is checked by a simple (but annoying to type up) proof. We simply show φ is an isomorphism. It suffices to show it is an isomorphism levelwise, meaning $\varphi_n : K_n \cong N(\mathcal{C})_n$. By the Yoneda lemma, this is equivalent to showing isomorphism

$$\text{sSet}(\Delta^n, K) \cong \text{sSet}(\Delta^n, N(\mathcal{C}))$$

for all $n \geq 0$.


We induct on this statement. This is obvious by construction for $n = 0$ and $n = 1$, since $N(\mathcal{C})_i = K_i$ when $i = 0, 1$, so assume the inductive hypothesis for $n \geq 2$, and choose $0 < i < n$ an integer. Precomposition by the canonical inclusion $\Lambda_i^n \hookrightarrow \Delta^n$ and postcomposition by φ induce a commutative diagram

$$\begin{array}{ccc} \text{sSet}(\Delta^n, K) & \longrightarrow & \text{sSet}(\Delta^n, N(\mathcal{C})) \\ \downarrow & & \downarrow \\ \text{sSet}(\Lambda_i^n, K) & \longrightarrow & \text{sSet}(\Lambda_i^n, N(\mathcal{C})) \end{array}$$

Recall by our first part of the proof that $N(\mathcal{C})$ satisfies property (2). Thus, since K and $N(\mathcal{C})$ satisfy (2), the vertical maps are bijective. It suffices to thus show the bottom map is bijective since isomorphisms satisfy 2-out-of-3. By the inductive hypothesis, there is an isomorphism $\text{sSet}(\Delta^{n-1}, K) \cong \text{sSet}(\Delta^{n-1}, N(\mathcal{C}))$. Recall from earlier that any face $\partial_j\Delta^n$ is isomorphic to Δ^{n-1} , and that the horn decomposes as maps for faces. Since the faces are isomorphic to Δ^{n-1} for which the statement follows, the statement must follow for each face, and by universality the statement follows for the horn, completing the proof. $\blacksquare$

Remark 1.1.2.6. We may refer to n -simplices of a simplicial set K as morphisms $\sigma : \Delta^n \rightarrow K$. We recall by our use of Yoneda in the above proof that such a morphism amounts to specifying an n -simplex $\sigma \in K_n$ and an image $1_{[2]} \mapsto \sigma$, justifying this terminology.


Remark 1.1.2.7. Of course, proposition 1.1.2.5 (2) is similar to definition 1.1.2.1, but we require it only for inner horns, and we require the filler be unique. It is fairly easy to see that it would be unreasonable to assume outer horns have unique fillers in proposition 1.1.2.5. For example, let $n = 2$. A horn $\Lambda_2^2 \rightarrow N(\mathcal{C})$ is a diagram of solid arrows



and a filler is a dotted arrow as indicated, rendering the diagram commutative. This is of course unreasonable to ask for, but it is reasonable for *groupoids*.

Simplicial sets are thus good models for ∞ -groupoids when K is a Kan complex, and are good models for regular categories when K satisfies proposition 1.1.2.5 (2). We earlier noted that ∞ -groupoids and regular 1-categories are both special cases of ∞ -categories, so we would expect some more general class of simplicial sets to give us a model. To see how this arises, we use our observations from proposition 1.1.2.5 and remark 1.1.2.3 to consider how a simplicial set K should be interpreted to yield an ∞ -category.

First, clearly objects are vertices, morphisms are edges, and identity, domain, and codomain are the same as remark 1.1.2.3. A 2-simplex $\sigma : \Delta^2 \rightarrow K$ is thought of as a diagram



together with a homotopy $\theta \simeq \psi\phi$ which witnesses the “commutativity” of the diagram (in higher category, commutativity of a diagram implies there are homotopies present as required to witness the composites). Simplices of higher dimension can be thought of similarly as verifying the commutivity of higher dimensional diagrams between higher level morphisms, which we will explore later as an aside from the main text.

However, this discussion has a massive issue: there is in general no way to compose two adjacent morphisms. However, inspecting the proof of proposition 1.1.2.5 gives us a property which makes this possible. A composable pair (g, f) determines a horn $\Lambda_1^2 \rightarrow K$, and the existence of their composite determines a filler $\sigma : \Delta^2 \rightarrow K$. We simply require that fillers for *inner* horns always exist.

The next problem is that of uniqueness. It is in fact fairly unnatural to require that fillers be unique. For example, recall the definition of composites of paths in a topological space, where we compress their domain along $I/2$ instead of I . We could just as easily compress one over $I/3$ and one over $2I/3$, and since the two choices are homotopic, it doesn’t matter which we choose. Since they are homotopic, they should both be valid choices for the composite of these homotopies, but each corresponds to a different filler for the relevant horn. With all of these considerations, we propose the following definition.

Definition 1.1.2.8 (Quasi-category). An ∞ -category (in particular, a quasi-category) is a simplicial set K which admits fillers for all inner horns. More precisely, for any $0 < i < n$, for all diagrams of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & K \\ \downarrow & \nearrow \text{dotted} & \\ \Delta^n & & \end{array}$$

there exists at least one (not necessarily unique) dotted arrow as indicated, rendering the diagram commutative. $\heartsuit$

This definition was first proposed by Boardman and Vogt in ??, motivated by proposition 1.1.2.5. They called them *weak Kan complexes*, and later André Joyal called them *quasi-categories*. Lurie calls them ∞ -categories, although Joyal's terminology is more commonly used when referring to the specific underlying model.

Example 1.1.2.9. Any Kan complex as defined in definition 1.1.2.1 is an ∞ -category. In particular, in view of remark 1.1.2.2, the singular complex $\text{Sing } X$ of any topological space X is an ∞ -category.

Example 1.1.2.10. By proposition 1.1.2.5, the nerve of any category is an ∞ -category. We will oftentimes abuse terminology and regard 1-categories $\mathcal{C}$ as ∞ -categories by identifying them with their nerve $N(\mathcal{C})$. By way of this identification, we may view 1-category theory as a special case of the theory of ∞ -categories.

Notation 1.1.2.11. We will use two different notations for simplicial sets. When viewing a simplicial set K as an ∞ -category, we will generally denote it with a calligraphic letter ($\mathcal{C}, \mathcal{D}, \mathcal{E} \dots$), and otherwise we denote them by capital roman letters ($K, X, Y \dots$). We may also employ the latter notation when comparing different models for ∞ -categories, to avoid confusion.

Before we continue with the text, I want to make a couple remarks that will be useful for later.

Proposition 1.1.2.12. Let K be a simplicial set. Specifying a horn $\Lambda_i^n \rightarrow K$ amounts to specifying $(n-1)$ -simplices τ_j for all $0 \leq j \neq i \leq n$ which satisfy the compatibility condition $d_k(\tau_j) = d_{j-1}(\tau_k)$ for all $k < j$, by defining $d^j \mapsto \tau_j$.

Proof. Recall that unions are pushouts in $\mathbf{sSet}$, meaning morphisms $\Lambda_i^n \rightarrow K$ are precisely pushout diagrams

$$\begin{array}{ccc} \bigcap_{j \in [n], j \neq i} \partial_j \Delta^{n-1} & \hookrightarrow & \partial_j \Delta^{n-1} \\ \downarrow & & \downarrow \\ \partial_k \Delta^{n-1} & \hookrightarrow & \bigcup_{j \in [n], j \neq i} \partial_j \Delta^{n-1} = \Lambda_i^n \\ & \searrow & \downarrow \text{dotted} \\ & & K \end{array}$$

(Note: A curved arrow also goes from $\partial_k \Delta^{n-1}$ to K)

where the pushout is over all $j \in [n] \setminus \{i\}$, but it is hard to draw this many arrows so we only draw

two. The case for $n = 1$ is trivial, so we assume $n \geq 2$.

Recall from the proof of proposition 1.1.2.5 that there is an isomorphism of simplicial sets $\partial_j \Delta^n \cong \Delta^n$ by mapping $\delta^j \alpha \mapsto \alpha$, which is an isomorphism by the epi-mono factorization in $\mathbf{\Delta}$ ([ML98, Tal25]). Note that the map $\delta^j \mapsto \tau_j$ decomposes as

$$\begin{array}{ccc} & d^j \mapsto \tau_j & \\ & \curvearrowright & \\ \partial_j \Delta^{n-1} & \xrightarrow[d^j \alpha \mapsto \alpha]{\cong} \Delta^{n-1} & \xrightarrow[1_{[n-1]} \mapsto \tau_j]{\rightarrow} K, \end{array}$$

where we recall that $1_{[n-1]} \mapsto \tau_j$ is a map of simplicial sets by the Yoneda lemma, as noted in the proof of proposition 1.1.2.5. We review the proof of this in the proof of proposition 1.1.2.13. Thus, $d^j \mapsto \tau_j$ decomposes as a morphism of simplicial sets, and thus is a morphism of simplicial sets for each j . Thus by universality of pushout, and by levelwise computation of universals in functor categories ([ML98]), the claimed morphism $\sigma : \Lambda_i^n \rightarrow K$ exists precisely when the above pushout diagram commutes.

First, we prove the easy direction. Let $k < j$, and suppose the diagram commutes, meaning there is indeed a morphism $\sigma : \Lambda_i^n \rightarrow K$ with $d^j \mapsto \tau_j$ for each $j \neq i$ in $[n]$. Since Λ_i^n is a simplicial subset of Δ^n , faces are given by precomposition. Since σ is a morphism of simplicial sets, we have the string of equalities

$$d_k(\tau_j) = d_k \sigma(d^j) = \sigma d_k(d^j) = \sigma(d^j d^k) \stackrel{!}{=} \sigma(d^k d^{j-1}) = \sigma d_{j-1}(d^k) = d_{j-1} \sigma(d^k) = d_{j-1}(\tau_k),$$

where the equality $!$ is given by the cosimplicial identity $d^j d^k = d^k d^{j-1}$ for $k < j$.

Now we prove the converse. Suppose the compatibility condition holds. We do this in three steps:

1. We show for $n \geq 2$, the intersection $\bigcap_{j \neq i} \partial_j \Delta^{n-1}$ has precisely one 0-simplex i , and all higher levels m have precisely one m -simplex $s_m s_{m-1} \dots s_0(i)$.
2. We show the maps $\partial_{j,k} \Delta^{n-1} \rightarrow K$ agree on i .
3. We show this implies the maps $\partial_{j,k} \Delta^{n-1} \rightarrow K$ agree on the full intersection.

(1) First, we show i is the unique 0-simplex. Note that the 0-simplices of $\Delta^{n-1} = [n-1]$, and the 0-simplices of each $\partial_j \Delta^{n-1}$ are thus $d^j[n-1]$. Since $j \neq i$, none of the cofaces d^j skip i , meaning $i \in \text{im } d^j$ for all j , and thus j is a 0-simplex. Uniqueness follows from the fact that if $k \in [n-1]$ and $k \neq i$, then there exists $\partial_k \Delta^{n-1}$ in the union, the 0-simplices of which are $d^k[n-1]$, which excludes k by definition.

Now we show all higher m -simplices are the degenerate simplices $s_m \dots s_0(i)$ by induction. A 1-simplex σ must have its faces $d_0(\sigma), d_1(\sigma)$ be 0-simplices of the union, meaning both its faces must be i . Recall that since we are working in a simplicial subset of Δ^n , faces are precomposition with cofaces, so this implies

$$0 \mapsto 0 \mapsto \sigma(0) = 0 \mapsto 1 \mapsto \sigma(1) = i,$$

and thus σ is necessarily the map $0, 1 \mapsto i$. Thus there is a unique 1-simplex. This can be viewed as the degenerate simplex

$$s_0(i) = 0, 1 \mapsto 0 \mapsto i,$$

giving the required form. Now assume the inductive hypothesis for m . Since $s_m \dots s_0(i)$ is the unique m -simplex, all faces of an $(m+1)$ -simplex σ must be $s_m \dots s_0(i)$. By the same logic as above,

we immediately see that σ is the map $[m+1] \mapsto i$, and thus admits the representation $s_{m+1} \dots s_0(i)$, concluding this section of the proof.

(2) Let $\sigma_j : \partial_j \Delta^{n-1} \rightarrow K$ and $\sigma_k : \partial_k \Delta^{n-1} \rightarrow K$ be the maps in question. This can be proven directly, but we use induction

(3) Keeping notation as before, we clearly have by inducting on the morphism condition for simplicial sets that for each m -simplex α in the intersection,

$$\sigma_j(\alpha) = \sigma_j s_m \dots s_0(i) = s_0 \dots s_m \sigma_j(i) = s_0 \dots s_m \sigma_k(i) = \sigma_k s_m \dots s_0(i) = \sigma_k(\alpha).$$

■

Proposition 1.1.2.13. *Call an n -tuple $(f_1, \dots, f_n)$ of 1-simplices composable if $d_1(f_i) = d_0(f_{i-1})$ for each $0 < i \leq n$. Then the datum of a composable n -tuple defines an inner horn $\Lambda_i^n \rightarrow K$.*

A filler for an inner horn $\sigma_0 : \Lambda_i^n \rightarrow K$ is an n -simplex $\tau \in K_n$ such that the morphism of simplicial sets $\sigma : 1_{[n]} \mapsto \tau$ agrees with the horn Λ_i^n in the sense that for each $j \neq i$ and each face $d^j : [n-1] \rightarrow [n]$,

$$\sigma_0(d^j) = \sigma(d^j).$$

1.1.3 Equivalences of Topological Categories

We return for a short time to our simpler notion of topological categories, in order to get a feel for what an ∞ -categorical notion of an *equivalence of categories* should be.

Bibliography

- [BV73] J. M. Boardman and Rainer Vogt. *Homotopy invariant algebraic structures on topological spaces*. Lecture notes in mathematics, 347. Springer, Berlin, 1973.
- [Fri23] Greg Friedman. An elementary illustrated introduction to simplicial sets, 2023. URL: <https://arxiv.org/abs/0809.4221>.
- [Joy08a] André Joyal. Quasi-categories and kan complexes. Lecture notes, 2008.
- [Joy08b] André Joyal. The theory of quasi-categories and its applications. Unpublished manuscript, 2008.
- [Lur09] Jacob Lurie. *Higher Topos Theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, 2009. Available online at <https://www.math.ias.edu/~lurie/papers/HTT.pdf>.
- [ML98] Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer, second edition, 1998.
- [Qui67] Daniel G. Quillen. *Homotopical Algebra*, volume 43 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, Heidelberg, 1967. Unofficial revised and modernized version at <https://aareyanmanzoor.github.io/assets/books/homotopical-algebra.pdf>.
- [Rie11] Emily Riehl. A leisurely introduction to simplicial sets. Expository notes available online, 2011. URL: <https://math.jhu.edu/~eriehl/ssets.pdf>.
- [Tal25] Grant Talbert. Introduction to simplicial sets. Unpublished manuscript, 2025. URL: <https://github.com/granttalbert/ssets>.