The Null Space of a Matrix

Benjamin Ochoa

January 16, 2015

Below is a summary of the (right) null space and left null space of a matrix, and how to compute them using singular value decomposition (SVD).

(Right) null space

The (right) null space of a matrix $A \in \mathbb{R}^{m \times n}$ is the matrix X = null(A) such that

$$AX = 0$$

where $X \in \mathbb{R}^{n \times (n-r)}$ and $r = \text{rank}(A) \leq \min(m, n)$.

Left null space

The left null space of a matrix $A \in \mathbb{R}^{m \times n}$ is the matrix Y such that

$$YA = 0$$

where $Y \in \mathbb{R}^{(m-r)\times m}$ and $r = \operatorname{rank}(A) \leq \min(m, n)$. The left null space may be calculated using the (right) null space as $Y = (\operatorname{null}(A^{\top}))^{\top}$.

Computation of the right and left null space using SVD

The singular value decomposition (SVD) of a matrix $A \in \mathbb{R}^{m \times n}$ may be written as

$$A = U\Sigma V^{\top}$$

where the orthogonal matrix $U \in \mathbb{R}^{m \times m}$, the diagonal matrix $\Sigma = \operatorname{diag}(\sigma_1, \sigma_2, \dots, \sigma_{\min(m,n)}) \in \mathbb{R}^{m \times n}$, where $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{\min(m,n)} \geq 0$, and the orthogonal matrix $V \in \mathbb{R}^{n \times n}$. σ_i is the *i*-th singular value of A, and the *i*-th column of V and *i*-th column of V are the corresponding left singular vector and right singular vector, respectively, of V. The rank V of V is the number of nonzero singular values. The (right) null space of V is the columns of V corresponding to singular values equal to zero. The left null space of V is the rows of V corresponding to singular values equal to zero (or the columns of V corresponding to zero, transposed).