

A-Levels Math Notes

Grass

Contents

1	Inequalities and Equations	1
1.1	Solving Inequalities	1
1.2	Modulus Inequalities	1
1.3	System of Linear Equations	1
1.4	Summary	2
2	Sequences and Series	3
2.1	Binomial Theorem and Series	3
2.2	APGP	4
2.3	Summation	4
2.4	Method of Differences	5
3	Recurrence Relations	6
4	Induction	8
5	Differentiation	9
6	Integration Techniques	10
6.1	Basic Integration (IBS, IBP, etc)	10
6.2	Areas & Volumes	11
6.3	Numerical Methods	12
6.3.1	Trapezium Rule	12
6.3.2	Simpson's Rule	13
7	Complex Numbers	14
7.1	Complex Number I	14
7.2	Complex Numbers II	15
8	Linear Algebra	19
9	Numerical Methods	24
10	Graphing Techniques	25
10.1	Graphing 'Familiar' Functions and Asymptotic bois	25
10.2	Conics	26
10.3	Parametric Equations	27
10.4	Scaling, Translations, and Reflections	27
10.5	$ f(x) $ and $f(x)$	28
10.6	$y = \frac{1}{f(x)}$	28

11 Polar Curves	29
12 Conic Sections	33
13 Functions	35
14 Permutations and Combinations	36
15 Vectors	38
16 Probability	40
17 Differential Equations	41
17.1 First Order D.E.s	41
17.1.1 Elementary Solving Techniques	41
17.1.2 Numerical Methods	41
17.2 Second Order D.E.	42
17.3 Applications	43
17.3.1 Exponential Growth	43
17.3.2 Logistics Growth	43
17.3.3 Harvesting	45
17.3.4 Physics	45
18 Discrete Random Variables	47
19 Special Discrete Random Variables	48
20 Continuous Random Variables	51
21 Special Continuous Random Variables	52
22 Sampling and Estimation	55
23 Statistics: Hypothesis Testing	59
24 Correlation and Linear Regression	61
25 Bibliography	63

Inequalities and Equations

1.1 Solving Inequalities

General Methods

1. Quadratic formula for factorisation / finding roots (of polynomial).
2. Completing the square.
3. Discriminant/Completing the square to eliminate factors which are *always* positive or negative (e.g. removing $x^2 - 3x + 4$). *Note to include coefficient of x^2 in the argument.*
4. GC (include sketch).
5. *Rational Functions*: Move everything to one side by adding or subtracting, then use a number line.

1.2 Modulus Inequalities

Fact

Given $x \in \mathbb{R}$, we have that

- $|x| \geq 0$,
- $|x^2| = |x|^2 = x^2$,
- $\sqrt{x^2} = |x|$.

And as long as $x \in \mathbb{R}^+$,

- $\sqrt{x^2} = |x|$.

Useful Properties

For every $x, k \in \mathbb{R}$:

- (a) $|x| < k$ iff $-k < x < k$.
- (b) $|x| > k$ iff $x < -k$ or $x > k$.

1.3 System of Linear Equations

General Information

- For more complicated real-world-context qns, try playing around with the values (e.g. use simultaneous equations) first. It may work out nicer than expected.

1.4 Summary

G.C. Skills

1. Plotting curves $y = f(x)$ in G.C.
2. How to use simultaneous equation solver.

Important Notes

- Eliminating Factors — *only* works for $c = 0$ in $f(x) \geq c$ or $f(x) \leq c$.
Counterexample: It is false that $P(x) = x(3x^2 - 9x + 10) \leq 2$ iff $x \leq 2$. Notice that $P(1.8) = 6.336 \not\leq 2$.
- Discriminant — include coefficient of x^2 in argument.
- When using factor elimination to remove some $f(x)$, we only need to say that “ $f(x)$ is negative”.
- Rational functions — exclude the values that causes division by zero to occur.
- With inequalities, be really careful about multiplication! If $x > y$ and $z > 0$, then $xz > yz$.
- Cross multiplication preserves order for $\frac{x}{y} < \frac{x'}{y'}$ iff y and y' are *both* positive or negative.
Note the counterexample $\frac{1}{2} < \frac{1}{-3}$.
- Squaring preserves/reverses order for $x < y$ iff x and y are *both* positive or negative.
- Can't necessarily use differentiation to solve if qns asks for algebraic method.
- Safer to graph out the two functions separately!
- Be careful about whether to include equality! Don't forget to account for it!
- Note that when solving for $|x| = y$, $|x| < y$, etc, y must be greater than or equal to 0. In other words, there may be solutions we will need to reject.
- Carelessness: Look at the question carefully! If they ask for a *set* of values, then rmb to give it as a *set*!
- Exponentiation and Logarithms: Simply use \ln and avoid \log_c for $c < 1$.
Order is *Preserved* under exponentiation/logarithms if the base is *larger than* one. Otherwise, when it is *less than* one, the order is *reversed*. <https://www.desmos.com/calculator/gd8z5fa0bg>
- For more complicated real-world-context qns, try playing around with the values (e.g. use simultaneous equations) first. It may work out nicer than expected.

Sequences and Series

2.1 Binomial Theorem and Series

Theorem 2.1: The Binomial Theorem

$$(a + b)^n = \sum_{r=0}^n \binom{n}{r} a^{n-r} b^r,$$

where $n \in \mathbb{Z}^+$.

Theorem 2.2: The Binomial Series

$$(1 + x)^p = \sum_{r=0}^{\infty} \binom{p}{r} x^r,$$

where $p \in \mathbb{Q}$, $|x| < 1$, and

$$\binom{p}{r} := \frac{p(p-1) \cdots (p-r+1)}{r!}.$$

Corollary 2.3

Clearly,

$$(a + x)^p = a^p \left(1 + \frac{x}{a}\right)^p = a^p \sum_{r=0}^{\infty} \binom{p}{r} \frac{x^r}{a^r},$$

under the same conditions.

Fact

We can expand $(a + x)^p$ in descending powers of x by using $(a + x)^p = x^p \left(1 + \frac{a}{x}\right)^p$.

Note

Sometimes computing a couple terms can be useful in finding a pattern. For example, to get the coefficient of x^k explicitly.

2.2 APGP

Basics

	AP	GP
u_n	$u_n = S_n - S_{n-1}$ $u_n = a + (n-1)d$	$u_n = ar^{n-1}$
S_n	$S_n = \frac{n}{2}[2a + (n-1)d]$ $= \frac{n}{2}(a + \ell)$	$S_n = \frac{a(1-r^n)}{1-r}$ $= \frac{a(r^n-1)}{r-1}$
S_∞	Always diverges	$S_\infty = \frac{a}{1-r}$ when $ r < 1$
Prove AP/GP	I Show $u_n - u_{n-1}$ is a constant / independent of n . II Show $u_n = a + (n-1)d$ explicitly.	I Show $\frac{u_n}{u_{n-1}}$ is constant / independent of n . II Show $u_n = ar^{n-1}$ explicitly
Mean	$u_{n+1} = \frac{u_n + u_{n+2}}{2}$. (Arithmetic Mean)	$\frac{u_{n+1}}{u_n} = \frac{u_{n+2}}{u_{n+1}}$ $u_{n+1}^2 = u_n \cdot u_{n+2}$ (Geometric Mean)

Important Notes

Applications: Write out a few terms in a table and observe the trend. (You can literally say “By observing a trend, ...”)

G.C. Skills

Table function

1. Enter eqn into GC.
2. 2nd graph to show table
3. 2nd tblset for setup options

2.3 Summation

Fact

$$\begin{aligned}
 \sum_{i=m}^n f(i) + g(i) &= \sum_{i=m}^n f(i) + \sum_{i=m}^n g(i) \\
 \sum_{i=m}^n af(i) &= a \sum_{i=m}^n f(i) \\
 \sum_{i=m}^n a &= (n-m+1)a, \text{ for any constant } a \\
 \sum_{i=m}^n f(i) &= \sum_{i=1}^n f(i) - \sum_{i=1}^{m-1} f(i)
 \end{aligned}$$

Note

- Look out for sums being AP and GPs.
- Results to be provided:

$$\sum_{i=1}^n i^2 = \frac{n}{6}(n+1)(2n+1)$$

$$\sum_{i=1}^n i^3 = \frac{1}{4}n^2(n+1)^2$$

2.4 Method of Differences**General Information**

$$\sum_{i=1}^n u_i = \sum_{r=1}^n f(r) - f(r-1) = f(n) - f(0).$$

- Explain convergence of a function $h(x) = f(x) + g(x)$: As $n \rightarrow \infty$, $f(x) \rightarrow 0$ and $g(x) \rightarrow 0$. Hence, $h(x)$ converges to...

G.C. Skills

Know how to generate sequences using the two methods:

1. Table method — Present by showing two consecutive values of n so that the values of the sequence are of opposite signs. E.g.:

n	S_n
182	$561.28 < 0$
183	$-1935.91 < 0$

2. 2nd stat seq (& we can use operations on seq, e.g. sum)

Recurrence Relations

General Information

1. Recurrence relation is *homogenous* if constant (b below) is zero.
2. First order linear recurrence relation: $u_n = au_{n-1} + b$, with $a \neq 0$.
3. Second order *homogenous* linear recurrence relation: $u_n = a_1u_{n-1} + a_2u_{n-2}$, $a_2 \neq 0$.
4. Solving RRs in general:
 - (a) Continually expand u_n in terms of u_{n-1} , then in terms of u_{n-2} , ..., till an explicit formula is obtained.
 - (b) Use a_1 to generate a_2, a_3, \dots, a_n .
5. Solving 1st order RRs, $u_{n+1} = au_n + b$:
 - (a) Iteration — Essentially technique 4(a). Will need to use G.P. formula at the end.
 - (b) Rewriting RR + Using G.P. Formulas ((c) is better)
 - i. Write RR as $u_n - k = a(u_{n-1} - k)$, where $k = \frac{b}{1-a}$. Let $v_n = u_n - k$.
 - ii. $\frac{v_n}{v_{n-1}} = a$, a constant and $\{v_n\}$ is a G.P. with first term $v_1 - k$ and common ratio a .
 - iii. So, $v_n = (u_1 - k)a^{n-1}$, and accordingly, $u_n = v_n + k = (u_1 - k)a^{n-1} + k$.
 - (c) ★ Let $u_n = Aa^n + \frac{b}{1-a}$. Then solve for the constant A with info provided.
6. Solving 2nd order (homogenous) RRs, $u_{n+2} = au_{n+1} + bu_n$:
 Assume $u_n = m^n$, then $m^2 - am - b = 0$ (is the *characteristic/auxillary equation* of the RR).
 Solve for the roots, say m_1 and m_2 . Then, the general solution for u_n is

$$u_n = \begin{cases} Am_1^n + Bm_2^n & \text{if } m_1, m_2 \in \mathbb{R}, \text{ and } m_1 \neq m_2, \\ (C + Dn)m_1^n & \text{if } m_1, m_2 \in \mathbb{R}, \text{ and } m_1 = m_2, \\ r^n[A \cos(n\theta) + B \sin(n\theta)] & \text{if } m_1, m_2 \in \mathbb{C}, m_1 = re^{i\theta}, \text{ and } m_2 = re^{-i\theta}. \end{cases}$$

Note

Let $x_{n+1} = f(x_n)$ and $L := \lim x_n$. To find the possible values of L , we can compare the graph of $y = f(x)$ against the identity function $y = x$. This is done by seeing if $f(x) < x$, $f(x) = x$, or $f(x) > x$.

Example 3.1




Figure 3.1: The RR $x_{n+1} = \sqrt{x_n^2 - 3x_n + 4} + x_n - 3$.

Let $f(x) = \sqrt{x^2 - 3x + 4} + x - 3$.

1. Suppose $x_1 \leq \frac{3+\sqrt{29}}{2}$. For $x_1 < \frac{3-\sqrt{29}}{2}$, we see that $f(x) > x$. So x_n increases till $\frac{3-\sqrt{29}}{2}$. While for $\frac{3-\sqrt{29}}{2} < x_1 < \frac{3+\sqrt{29}}{2}$, we have $f(x) < x$. Thus x_n decreases till $\frac{3-\sqrt{29}}{2}$. Notice the graphs intersects at $x = \frac{3-\sqrt{29}}{2}$. So, when $x_n = \frac{3-\sqrt{29}}{2}$, if ever, then $x_{n+1} = x_n$. That is, $L = \frac{3-\sqrt{29}}{2}$.
2. Similarly, if $x_1 = \frac{3+\sqrt{29}}{2}$, then $x_n = \frac{3+\sqrt{29}}{2}$ is a constant function; $L = \frac{3+\sqrt{29}}{2}$.
3. Presume that $x_n > \frac{3+\sqrt{29}}{2}$. Then, $f(x) > x$ tells us x_n is an increasing sequence that is unbounded. In other words, L does not exist.

Induction

General Information

Let $P(x)$ be the statement that “...”.

When $n = 1, \dots$

$\implies P(1)$ is true.

Assume $P(k)$ is true for some $k \in \mathbb{Z}^+$.

Then, ...

$\implies P(k + 1)$ is true.

Therefore, since $P(1)$ is true and $P(k)$ true $\implies P(k + 1)$ true, $P(n)$ is true for all $n \in \mathbb{Z}^+$.

Differentiation

Definition

1. A function f is called (strictly) increasing on an interval I iff $f'(x) > 0$ for all $x \in I$.
2. A function f is called monotonically increasing on an interval I iff $f'(x) \geq 0$ for any $x \in I$.

General Information

1. How to sketch the graph of the integral or derivative of a function f .
2. Relationship btw. a function f and its derivative, f' :

$y = f(x)$	$y = f'(x)$
Vertical asymptote at $x = a$	Vertical asymptote at $x = a$.
Horizontal asymptote at $y = a$	Horizontal asymptote $y = 0$.

3. Recap:

$f(x)$	$f'(x)$
$\sin^{-1}\left(\frac{x}{a}\right)$	$\frac{1}{\sqrt{a^2 - x^2}}, x < a$
$\cos^{-1}\left(\frac{x}{a}\right)$	$-\frac{1}{\sqrt{a^2 - x^2}}, x < a$
$\tan^{-1}\left(\frac{x}{a}\right)$	$\frac{a}{a^2 + x^2}, x \in \mathbb{R}$
$\log_a(f(x))$	$\frac{1}{x \ln(a)}$
a^x	$a^x \ln(a)$

4. Implicit differentiation: $\frac{dz}{dx} = \frac{dz}{dy} \cdot \frac{dy}{dx}$. ★ Makes life much easier (e.g. finding $f^{(n)}(x)$).

5. Parametric Differentiation: $\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$.

6. Small angle approximation:

(a) $\sin(x) \approx x$,

(b) $\cos(x) \approx 1 - \frac{x^2}{2}$,

(c) $\tan(x) \approx x$.

7. Maclaurin Series:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n.$$

Integration Techniques

6.1 Basic Integration (IBS, IBP, etc)

General Information

1. Factor Formulae ★ (must *rm*b):

(a) $\sin(mx) \cos(nx) = \frac{1}{2}[\sin((m+n)x) + \sin((m-n)x)],$

(b) $\cos(mx) \cos(nx) = \frac{1}{2}[\cos((m+n)x) + \cos((m-n)x)],$

(c) $\sin(mx) \sin(nx) = -\frac{1}{2}[\cos((m+n)x) - \cos((m-n)x)].$

2. Common classes of integrals:

- (a) Apply partial fractions:

$$\int \frac{f(x)}{g(x)} dx.$$

- (b) Split $px + q$, then complete the square:

$$\int \frac{px + 1}{\sqrt{ax^2 + bx + c}} dx \quad \text{or} \quad \int \frac{px + 1}{ax^2 + bx + c} dx$$

3. Integration by Substitution:

$$\int f(x) dx = \int f(x) \frac{dx}{du} du.$$

4. Use Pythagoras' Theorem / draw a right-angled triangle to help with trig conversions, e.g.:

$$\tan(\theta) \quad \text{to} \quad \frac{x+1}{\sqrt{2-(x+1)^2}}.$$

5. Integration by Parts:

Let $u = g(x)$, $\frac{dv}{dx} = h(x)$,
 $\frac{du}{dx} = g'(x)$, $v = \int h(x) dx$.

$$\int u \left(\frac{dv}{dx} \right) dx = uv - \int v \left(\frac{du}{dx} \right) dx.$$

6.2 Areas & Volumes

General Information

1. Volume of revolution when rotated about x -axis:

(a) The disc method

$$\int_{x_1}^{x_2} \pi y^2 dx = \int_{x=x_1}^{x=x_2} \pi y^2 \frac{dx}{dt} dt.$$

(b) The shell method:

$$\int_{x_1}^{x_2} 2\pi y x dy.$$

2. Arc length:

$$\int_{x_1}^{x_2} \sqrt{\left(\frac{dy}{dx}\right)^2 + 1} dx = \int_{y_1}^{y_2} \sqrt{\left(\frac{dx}{dy}\right)^2 + 1} dy = \int_{t_1}^{t_2} \sqrt{\left(\frac{dy}{dt}\right)^2 + \left(\frac{dx}{dt}\right)^2} dt = \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$

3. Surface area of revolution when rotated about x -axis:

$$\int_{x_1}^{x_2} 2\pi y \sqrt{\left(\frac{dy}{dx}\right)^2 + 1} dx = \int_{y_1}^{y_2} 2\pi y \sqrt{\left(\frac{dx}{dy}\right)^2 + 1} dy = \int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dy}{dt}\right)^2 + \left(\frac{dx}{dt}\right)^2} dx = \int_{\alpha}^{\beta} \frac{1}{2} r^2 d\theta.$$

★ Rotating about x -axis $\implies y$ in integrand

Rotating about y -axis $\implies x$ in integrand.

6.3 Numerical Methods

6.3.1 Trapezium Rule

General Information

1. Formula for n intervals, or $(n+1)$ ordinates, of width $h := (b - a)/n$:

$$\int_a^b y \, dx = \frac{h}{2}(y_0 + 2y_1 + 2y_2 + \cdots + 2y_{n-1} + y_n).$$

2. Illustration




Figure 6.1: Trapezium rule

3. Error:

- (a) Concave upwards, i.e. $(f'(x)$ is increasing / $f''(x) > 0$) \implies overestimation.
- (b) Concave downwards, i.e. $(f'(x)$ is decreasing / $f''(x) < 0$) \implies underestimation.

6.3.2 Simpson's Rule

General Information

1. Formula for n intervals, or $(n+1)$ ordinates, of width $h := (b - a)/n$:

$$\int_a^b y \, dx = \frac{h}{3}(y_0 + 4y_1 + 2y_2 + 4y_3 + \cdots + 2y_{n-2} + 4y_{n-1} + y_n).$$

Note that the number of intervals n should be *even*, that of ordinates *odd*.

2. Illustration




Figure 6.2: Simpson's rule

Note

Accuracy of the Trapezium rule vs Simpson's Rule: "Simpson's Rule uses *quadratic curves* to interpolate the points on the curve so it usually *gives a better approximation* to the actual curve than the trapezium rule which uses *straight lines* to interpolate the ordinates."

Complex Numbers

7.1 Complex Number I




Figure 7.1: Argand diagram.

General Information

1. Find the square root of $x + iy$: Let $\sqrt{x + iy} = a + bi$. Then square both sides & solve.
2. Simplifying fractions: multiply by the denominator's conjugate

$$\frac{a + bi}{c + di} = \frac{a + bi}{c + di} \cdot \frac{c - di}{c - di} = \dots$$

3. Polynomials:

- (a) Fundamental Theorem of Algebra: If $p(z) := \sum_{i=0}^n a_i z^i$ is a polynomial of degree $n \geq 1$ with complex coefficients, then there exists complex numbers c_i for each $1 \leq i \leq n$ such that

$$p(z) = a_n \prod_{i=1}^n (z - c_i).$$

- (b) If a polynomial in real coefficients only has root $a + bi$, then $a - bi$ is another root.

Example 7.1

Find the roots of $iz^2 + 2z + 3i = 0$.

$$z^2 - 2iz + 3 = 0$$

$$z = \frac{2i \pm \sqrt{(2i)^2 - 4(1)(3)}}{2(1)} = i \pm \frac{\sqrt{-16}}{2} = i \pm 2i$$

So, $z = 3i$ or $z = -i$.

Example 7.2: N2010/2/1

One root of the equation $x^4 + 4x^3 + ax + b = 0$, where a and b are real, is $x = -2 + i$. Find the values of a and b and the other roots.

Substitute $-2 + i$ into the equation:

$$\begin{aligned} (-2 + i)^4 + 4(-2 + i)^3 + (-2 + i)^2 + a(-2 + i) + b &= 0 \\ -12 + 16i &= 2a - b - ai \\ a = -16, \quad 2a - b &= -12 \end{aligned}$$

Therefore, $a = -16$, $b = -20$.

Since all the coefficients of the polynomial are real (**explain**), $-2 - i$ is another root. Now, $x^4 + 4x^3 + ax + b = (x - (-2 + i))(x - (-2 - i))(cx + d)$ for some $c, d \in \mathbb{R}$.

Accordingly, substitute $x = 0$, then $x = 2$, and solve. Alternatively, notice $x^4 + 4x^3 + ax + b = (x^2 - 2(-2)x + ((-2)^2 + 1^2))(x^2 + cx + d) = (x^2 + 4x + 5)(x^2 + cx + d)$. Either ways, we have $c = 0$ and $d = -4$. As such, the last two roots are $x = -2 \pm i$ and $x = \pm 2$.

- (c) Simultaneous equations: Solve as usual.
- (d) Properties of modulus: $|z_1^x z_2^y| = |z_1|^x |z_2|^y$, for any $x, y \in \mathbb{R}$.
- (e) Properties of arguments (same as log): $\arg(z) \in (-\pi, \pi]$ and $\arg(z_1^x z_2^y) = x \arg(z_1) + y \arg(z_2)$ for any $x, y \in \mathbb{R}$.
- (f) Polar form: $z = re^{i\theta}$.
- (g) Polar/Trigonometric form: $z = r[\cos(\theta) + i \sin(\theta)]$.

Note

Show that the value of w^n is either 2^n or 2^{-n} for integers n .

Then we **must** show that $w^n = \dots = \begin{cases} 2^n(1) = 2^n & \text{if } n \text{ even,} \\ 2^n(-1) = -2^n & \text{if } n \text{ odd.} \end{cases}$

7.2 Complex Numbers II

Theorem 7.1: De Moivre's Theorem

Let z be a complex number, n an integer, and θ an angle. Suppose $z = re^{i\theta}$. Then,

$$z^n = e^{in\theta} = r^n[\cos(n\theta) + i \sin(n\theta)].$$

General Information

1. All n th roots of any complex number are the same distance r from the origin and have the same angular separation, π/n .
2. Note that $1 + e^{i\theta} = e^{i\theta/2}(e^{-i\theta/2} + e^{i\theta/2})$.
3. For $z = re^{i\theta}$, we have $z^n + z^{-n} = 2\cos(n\theta)$ and $z^n - z^{-n} = 2i \sin(n\theta)$.
4. The geometric meaning of multiplying by i is a anti-clockwise rotation by π radians.
5. Loci (Use a *compass*)

- (a) The locus represented by $|z - a| = r$ (or $z = a + re^{i\theta}$) is a *circle* of radius r centered at $A(x, y)$ (where $a := x + iy$).




Figure 7.2: The locus of $|z - a| = r$.

- i. Either label the four points to the direct North, South, East, West of the circle, or denote the radius clearly.
- ii. The line segment, representing the furthest distance from a point to a circle, always cuts through the circle's centre. So, the distance

$$OP_{\max} - OP_{\min} = 2 \cdot \text{radius}.$$

- iii. The line segments, from a point to a circle that produces the largest angle, are tangents to the circle.




Figure 7.3: Maximum distance and angle of a point from a circle

- (b) The locus represented by $|z - a| = |z - b|$ is the *perpendicular bisector* of the line segment joining A and B .




Figure 7.4: The locus of $|z - a| = |z - b|$, a perpendicular bisector

- (c) The locus represented by $\arg(z - a) = \theta$ is the *half-line* from A (excluding A) that makes an angle θ with the *positive* real axis.




Figure 7.5: The locus of $\arg(z - a) = \theta$, a half-line.

6. There is no need to find the points of intersection between two loci, unless the questions states so.
7. Suppose we have a locus z represented by the predicate $P(z)$. Then, for any $a \in \mathbb{C}$, the locus of $z + a$ is represented by $P(z - a)$.
8. Say we are given a locus z represented by $|z - a| = r$, where $a = \alpha + \beta i$.
 - (a) The greatest and least value of $|z|$ are $|a| \pm r$, respectively.
 - (b) The greatest and least value of $\arg(z)$ can be obtained geometrically, or by plotting

$$Y_1 = \tan^{-1} \left(\frac{\beta \pm \sqrt{r^2 - (X - \alpha)^2}}{X} \right)$$

and finding the maximum/minimum point, respectively.




Figure 7.6: Brute Force Technique for Finding Maximum/Minimum angles.

Example 7.3: TQ 10(b)

Show that $\cot^2(2\pi/5)$ is a root of the equation $px^2 + qx + r = 0$, where we are given

$$\cot(4\theta) = \frac{\cot^4(\theta) - 6\cot^2(\theta) + 1}{4\cot^3(\theta) - 4\cot(\theta)}.$$

First notice that $\cot(8\pi/5) = -\cot(2\pi/5)$. So,

$$-\cot(2\pi/5) = \frac{\cot^4(2\pi/5) - 6\cot^2(2\pi/5) + 1}{4\cot^3(2\pi/5) - 4\cot(2\pi/5)}.$$

Simplifying gives

$$5[\cot^2(2\pi/5)]^2 - 10[\cot^2(2\pi/5)] + 1 = 0.$$

Thus, $x = \cot^2(2\pi/5)$ is a root of the equation $5x^2 - 10x + 1 = 0$.

Linear Algebra

Definition 8.1

A vector space (or linear space) V over a field \mathbb{F} consists of a set on which two operations (called addition and multiplication respectively here) are defined so that;

- (A) (V is Closed Under Addition) For all $\mathbf{x}, \mathbf{y} \in V$, there exists a unique element $\mathbf{x} + \mathbf{y} \in V$.
- (M) (V is Closed Under Scalar Multiplication) For all elements $a \in \mathbb{F}$ and elements $\mathbf{x} \in V$, there exists a unique element $a\mathbf{x} \in V$.

Such that the following properties hold:

- (VS 1) (Commutativity of Addition) For all $\mathbf{x}, \mathbf{y} \in V$, we have $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$.
- (VS 2) (Associativity of Addition) For all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$, we have $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$.
- (VS 3) (Existence of The Zero/Null Vector) There exists an element in V denoted by $\mathbf{0}$, such that $\mathbf{x} + \mathbf{0} = \mathbf{x}$ for all $\mathbf{x} \in V$.
- (VS 4) (Existence of Additive Inverses) For all elements $\mathbf{x} \in V$, there exists an element $\mathbf{y} \in V$ such that $\mathbf{x} + \mathbf{y} = \mathbf{0}$.
- (VS 5) (Multiplicative Identity) For all elements $x \in V$, we have $1\mathbf{x} = \mathbf{x}$, where 1 denotes the multiplicative identity in \mathbb{F} .
- (VS 6) (Compatibility of Scalar Multiplication with Field Multiplication) For all elements $a, b \in \mathbb{F}$ and elements $\mathbf{x} \in V$, we have $(ab)\mathbf{x} = a(b\mathbf{x})$.
- (VS 7) (Distributivity of Scalar Multiplication over Vector Addition) For all elements $a \in \mathbb{F}$ and elements $\mathbf{x}, \mathbf{y} \in V$, we have $a(\mathbf{x} + \mathbf{y}) = a\mathbf{x} + a\mathbf{y}$.
- (VS 8) (Distributivity of Scalar Multiplication over Field Addition) For all elements $a, b \in \mathbb{F}$, and elements $\mathbf{x} \in V$, we have $(a + b)\mathbf{x} = a\mathbf{x} + b\mathbf{x}$.

Theorem 8.2

Let V be a vector space and W a subset of V . Then W is a subspace of V iff the following 3 conditions hold for the operations defined in V .

- (a) $\mathbf{0} \in W$
- (b) $\mathbf{x} + \mathbf{y} \in W$ whenever $\mathbf{x} \in W$ and $\mathbf{y} \in W$.
- (c) $c\mathbf{x} \in W$ whenever $c \in \mathbb{F}$ and $\mathbf{x} \in W$.

Definition 8.3

A subset S of a vector space V *generates* (or *spans*) V iff $\text{span}(S) = V$. In this case, we also say that the vectors of S generate (or span) V .

Definition 8.4

Let V be a vector space and S a nonempty subset of V . A vector $v \in V$ is called a *linear combination* of vectors of S iff there exists a finite number of vectors u_1, u_2, \dots, u_n in S and scalars a_1, a_2, \dots, a_n in \mathbb{F} such that

$$v = \sum_{i=1}^n a_i u_i.$$

In this case we also say that v is a linear combination of u_1, u_2, \dots, u_n and call a_1, a_2, \dots, a_n the

coefficients of the linear combination

Definition 8.5

A subset S of a vector space V is called *linearly dependent* iff there exists a finite number of distinct vectors u_1, u_2, \dots, u_n in S and scalars a_1, a_2, \dots, a_n not all zero, such that

$$a_1 u_1 + a_2 u_2 + \dots + a_n u_n = \mathbf{0}.$$

Definition 8.6

A *basis* β for a vector space V is a linearly independent subset of V that generates V . If β is a basis for V , we also say that the vectors of β form a basis for V .

Theorem 8.7: The Rank-Nullity Theorem.

For any vector spaces V and W , and a linear operator $T: V \rightarrow W$, it holds that

$$\text{rank}(T) + \text{nullity}(T) = \dim(V).$$

General Information

- Let \mathbf{A} be an $m \times n$ matrix, and \mathbf{a}_j its j th column. For any $\mathbf{x} = (x_1 \ x_2 \ \dots \ x_n)^\top$,

$$\mathbf{A}\mathbf{x} = \sum_{j=1}^n x_j \mathbf{a}_j.$$

- Let \mathbf{A} and \mathbf{B} be matrices having n rows. For any matrix \mathbf{M} with n columns, we have

$$(\mathbf{A} \mid \mathbf{B}) = (\mathbf{MA} \mid \mathbf{MB}).$$

Definition 8.8

A system $\mathbf{Ax} = \mathbf{b}$ is *homogeneous* iff $\mathbf{b} = \mathbf{0}$; otherwise it is *nonhomogeneous*.

Theorem 8.9

For any matrix, its row space, column space, and rank are identical.

Theorem 8.10

A system $\mathbf{Ax} = \mathbf{b}$ of m linear equations in n unknowns has a solution space of dimension $n - \text{rank}(\mathbf{A})$.

Definition 8.11

A system $\mathbf{Ax} = \mathbf{b}$ of linear equations is *consistent* iff its solution set is nonempty; otherwise it is *inconsistent*.

Theorem 8.12: The Rouché-Capelli Theorem.

A system $\mathbf{Ax} = \mathbf{b}$ is consistent iff $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}|\mathbf{b})$.

Definition 8.13

A matrix is said to be in *reduced row echelon form* iff

- Any row containing a nonzero entry precedes any row in which all the entries are zero (if any).
- The first nonzero entry in each row is the only nonzero entry in its column.
- The first nonzero entry in each row is 1 and it occurs in a column to the right of the first

nonzero entry in the preceding row.

- Gaussian elimination.
 - In the forward pass, the augmented matrix is transformed into an upper triangular matrix in which the first nonzero entry of each row is 1 and it occurs in a column to the right of the first nonzero entry of each preceding row.
 - In the backward pass, the upper triangular matrix is transformed into reduced row echelon form by making the first nonzero entry of each row the only nonzero entry of its column.
- Gaussian elimination always reduces a matrix to its rref form.

- Let \mathbf{A} be an invertible $n \times n$ matrix. Then, for some elementary row matrices \mathbf{E}_1 to \mathbf{E}_p ,

$$\mathbf{E}_p \mathbf{E}_{p-1} \dots \mathbf{E}_1 (\mathbf{A} \mid \mathbf{I}_n) = \mathbf{A}^{-1} (\mathbf{A} \mid \mathbf{I}_n) = (\mathbf{I}_n \mid \mathbf{A}^{-1}).$$

In other words, we can perform Gaussian elimination, so that $(\mathbf{A} \mid \mathbf{I}_n) \rightarrow (\mathbf{I}_n \mid \mathbf{A}^{-1})$.

- Let $\mathbf{A} := (\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n)$ be $m \times n$ matrix, and $\mathbf{A}' := (\mathbf{a}'_1 \ \mathbf{a}'_2 \ \dots \ \mathbf{a}'_n)$ its rref. Then, $\{\mathbf{a}_{k_1}, \mathbf{a}_{k_2}, \dots, \mathbf{a}_{k_m}\}$ is linearly independent iff $\{\mathbf{a}'_{k_1}, \mathbf{a}'_{k_2}, \dots, \mathbf{a}'_{k_m}\}$ is. Moreover, the row space of \mathbf{A} and \mathbf{A}' are clearly identical.
- Finding a basis for an intersection of subspaces. Let V and W be subspaces of \mathbb{F}^n generated by the columns of the $n \times m$ matrix \mathbf{A} and $n \times k$ matrix \mathbf{B} , respectively. Find a basis for the subspace $V \cap W$.

1. First notice that $\mathbf{v} \in V \cap W$ iff

$$\mathbf{v} = \mathbf{A}\mathbf{x}_1 = \mathbf{B}\mathbf{x}_2$$

for some $\mathbf{x}_1 \in \mathbb{F}^m$ and $\mathbf{x}_2 \in \mathbb{F}^k$. That is,

$$(\mathbf{A} \ \mathbf{B}) \begin{pmatrix} \mathbf{x}_1 \\ -\mathbf{x}_2 \end{pmatrix} = \mathbf{0}.$$

So, equivalently, we write

$$(\mathbf{A} \ \mathbf{B}) \mathbf{y} = \mathbf{0}.$$

for some $\mathbf{y} \in \mathbb{F}^{m+k}$. As such, by row reducing $(\mathbf{A} \ \mathbf{B})$, we find a basis

$$\beta := \left\{ \begin{pmatrix} \mathbf{u}_1 \\ \mathbf{u}'_1 \end{pmatrix}, \begin{pmatrix} \mathbf{u}_2 \\ \mathbf{u}'_2 \end{pmatrix}, \dots, \begin{pmatrix} \mathbf{u}_r \\ \mathbf{u}'_r \end{pmatrix} \right\},$$

where $\mathbf{u}_i \in \mathbb{F}^m$ and $\mathbf{u}'_i \in \mathbb{F}^k$. Now, a generating set for $V \cap W$ is

$$\Gamma := \{\mathbf{A}\mathbf{u}_1, \mathbf{A}\mathbf{u}_2, \dots, \mathbf{A}\mathbf{u}_r\}.$$

Alternatively, another generating set for $V \cap W$ is

$$\Delta := \{\mathbf{B}\mathbf{u}'_1, \mathbf{B}\mathbf{u}'_2, \dots, \mathbf{B}\mathbf{u}'_r\}.$$

From here, it is simple to choose bases $\gamma \subseteq \Gamma$ and $\delta \subseteq \Delta$ for $V \cap W$.

(Naturally, it holds that $\mathbf{A}\mathbf{u}_i + \mathbf{B}\mathbf{u}'_i = \mathbf{0}$.)

2. An alternative method. By row reduction, we can calculate

$$\begin{aligned} r := \dim(V \cap W) &= \dim(U) + \dim(V) - \dim(U + V), \\ &= \text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}) - \text{rank}(\mathbf{A} \ \mathbf{B}), \\ &= \text{rank}(\mathbf{A}^\top) + \text{rank}(\mathbf{B}^\top) - \text{rank} \begin{pmatrix} \mathbf{A}^\top \\ \mathbf{B}^\top \end{pmatrix}. \end{aligned}$$

Then, a basis for $V \cap W$ can be formed by choosing r linearly independent columns of $(\mathbf{A} \ \mathbf{B})$, or rows of $\begin{pmatrix} \mathbf{A}^\top \\ \mathbf{B}^\top \end{pmatrix}$.

3. **Another alternative**, probably the best option! Skip the row reduction of \mathbf{A} and \mathbf{B} in the above method. We just reduce

$$(\mathbf{A} \ \mathbf{B}) \rightarrow (\mathbf{A}' \ \mathbf{B}').$$

Let \mathbf{c}_i and \mathbf{c}'_i be the i th column of $(\mathbf{A} \ \mathbf{B})$ and $(\mathbf{A}' \ \mathbf{B}')$, respectively. We compare the columns of \mathbf{A}' and \mathbf{B}' to find (with relative ease) a basis $\beta' := \{\mathbf{c}'_{i_1}, \mathbf{c}'_{i_2}, \dots, \mathbf{c}'_{i_r}\}$ for the intersection of the column spaces of \mathbf{A}' and \mathbf{B}' . Then, $\beta := \{\mathbf{c}_{i_1}, \mathbf{c}_{i_2}, \dots, \mathbf{c}_{i_r}\}$ is a basis for $V \cap W$ (the intersection of the column spaces of \mathbf{A} and \mathbf{B}).

4. **A fourth method** for when I learn about orthogonal complements.

Definition 8.14

Let $\mathbf{A} \in M_{n \times n}(\mathbb{F})$. If $n = 1$, so that $A = (a_1 1)$, we define $\det(\mathbf{A}) := a_1 1$. For $n \geq 2$, we define $\det(\mathbf{A})$ recursively as

$$\det(\mathbf{A}) := \sum_{j=1}^n (-1)^{1+j} \mathbf{A}_{1j} \cdot \det(\tilde{\mathbf{A}}_{1j}).$$

The scalar $\det(\mathbf{A})$ is called the *determinant* of \mathbf{A} and is also denoted by $|\mathbf{A}|$. The scalar

$$(-1)^{i+j} \det(\tilde{\mathbf{A}}_{ij})$$

is called the cofactor of the entry of \mathbf{A} in row i , column j .

- A matrix \mathbf{A} is invertible iff its determinant is nonzero.

Theorem 8.15

The determinant of a square matrix can be evaluated by cofactor expansion along any row. That is, if $\mathbf{A} \in M_{n \times n}(\mathbb{F})$, then for any integer $1 \leq i \leq n$,

$$\det(\mathbf{A}) = \sum_{j=1}^n (-1)^{i+j} \mathbf{A}_{ij} \cdot \det(\tilde{\mathbf{A}}_{ij}).$$

Here, $\tilde{\mathbf{A}}_{ij}$ is the $(n-1) \times (n-1)$ matrix obtained from \mathbf{A} by deleting its i th row and j th column.

Theorem 8.16

Let \mathbf{A} be an $n \times n$ matrix. Then,

$$\det(\mathbf{A}) = \det(\mathbf{A}^\top).$$

So, the determinant of a square matrix can also be evaluated by cofactor expansion along any column.

Theorem 8.17

Let \mathbf{A} be an invertible $n \times n$ matrix. Then,

$$\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \text{adj}(\mathbf{A}),$$

where $\text{adj}(\mathbf{A})$ is the adjugate/classical adjoint of \mathbf{A} . That is, the matrix whose (i, j) th entry is the (j, i) th cofactor $(-1)^{j+i} \det(\tilde{\mathbf{A}}_{ji})$.

Definition 8.18

A linear operator T on a finite-dimensional vector space V is called *diagonalisable* iff there is an ordered basis β for V such that $[T]_\beta$ is a diagonal matrix. A square matrix \mathbf{A} is called diagonalisable iff $L_{\mathbf{A}}$ is diagonalisable.

Definition 8.19

Let T be a linear operator on a vector space V . A nonzero vector $\mathbf{v} \in V$ is called an *eigenvector* of T iff there exists a scalar λ such that $T(\mathbf{v}) = \lambda\mathbf{v}$. The scalar λ is called the *eigenvalue* corresponding to the eigenvector \mathbf{v} .

Let \mathbf{A} be in $M_{n \times n}(\mathbb{F})$. A nonzero vector $v \in \mathbb{F}^n$ is called an *eigenvector* of \mathbf{A} iff v is an eigenvector of $L_{\mathbf{A}}$; that is, iff $\mathbf{A}v = \lambda v$ for some scalar λ . The scalar λ is called the eigenvalue of \mathbf{A} corresponding to the eigenvector v .

Definition 8.20

Let $\mathbf{A} \in M_{n \times n}(\mathbb{F})$. The polynomial $f(t) = \det(\mathbf{A} - t\mathbf{I}_n)$ is called the *characteristic polynomial* of \mathbf{A} .

- A matrix $\mathbf{A} \in M_{n \times n}(\mathbb{F})$ is diagonalizable iff there exists an ordered basis $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ for \mathbb{F}^n consisting of eigenvectors of \mathbf{A} , i.e. a eigenbasis. Furthermore, if \mathbf{Q} is the $n \times n$ matrix whose j th column is \mathbf{v}_j , then $\mathbf{A} = \mathbf{Q}^{-1}\mathbf{D}\mathbf{Q}$ is a diagonal matrix such that d_{jj} is the eigenvalue of A corresponding to \mathbf{v}_j . The matrix \mathbf{Q} is said to *diagonalise* \mathbf{A} .

- Hence, we obtain the following procedure to diagonalise a 3×3 matrix \mathbf{A} with three distinct eigenvalues.

1. Find the eigenvalues λ_1, λ_2 , and λ_3 of \mathbf{A} . They are just the roots of the characteristic polynomial of \mathbf{A} . [This can be done using the GC.](#)
2. Find an eigenvector \mathbf{v}_j corresponding to each eigenvalue λ_j by finding the nullspace of $\mathbf{A} - \lambda_j\mathbf{I}$.
3. Let $\mathbf{Q} = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$. Then,

$$\mathbf{D} := \mathbf{Q}^{-1}\mathbf{A}\mathbf{Q}$$

is a diagonal matrix.

Theorem 8.21: The Cayley-Hamilton Theorem.

Let T be a linear operator on a finite dimensional vector space V , and let $f(t)$ be the characteristic polynomial of T . Then $f(T) = T_0$, the zero transformation. That is, T “satisfies” its characteristic equation.

Corollary 8.22: The Cayley-Hamilton Theorem for Matrices.

Let A be an $n \times n$ matrix, and let $f(t)$ be the characteristic polynomial of A . Then, $f(A) = O$, the $n \times n$ zero matrix.

G.C. Skills

Finding eigenvalues of a matrix \mathbf{A} using the GC.

1. `2nd \Rightarrow x-1 (matrix) \Rightarrow Key in the matrix $\mathbf{A} - t\mathbf{I}$, e.g. into [A].`
2. `Plot Y1 = det ([A]).`
3. `2nd \Rightarrow trace \Rightarrow 2:zero \Rightarrow Find the roots.`

Numerical Methods

General Information

- The parity of the degree of a real polynomial is the same as that of its number of real roots.
- Let the real polynomial p given by $p(x) = a_{2n}x^{2n} + a_{2n-1}x^{2n-1} + \dots + a_0$ have coefficients $a_n > 0$ and $a_0 < 0$. Then, it has at least one positive and one negative root.
- Suppose we have some function $f: \mathbb{R} \rightarrow \mathbb{R}$ with a root α , whose value we want to approximate. There are three ways to obtain this approximation.

1. Linear interpolation on an interval $[a, b]$ containing α . We let $x_0 := b$ and

$$x_{i+1} := \frac{a|f(x_i)| + x_i|f(a)|}{|f(a)| + |f(x_i)|}.$$

– *Additional notes.*

2. Fixed-point Iteration. First select a function $F: \mathbb{R} \rightarrow \mathbb{R}$, such that $F(\alpha) = \alpha$, and choose some initial approximation x_0 to α . Then, we recursively define $x_{n+1} := F(x_n)$. The desired convergence behavior is for x_n to approach α .

– *Additional notes.*

G.C. Skills

Linear interpolation: finding an approximation to a root in $[a, b]$ up to n decimal places.

1. $Y_1 = f(x)$,
2. $a \rightarrow A$ and $b \rightarrow B$,
3. $\frac{B|Y_1(A)| + A|Y_1(B)|}{|Y_1(A)| + |Y_1(B)|}$,
4. $\text{Ans} \rightarrow A$ or B (choose the one that has the opposite sign to Ans),
5. Repeat steps 4 to 5,
6. Terminate this process when the approximations are consistent up to n decimal places.

Graphing Techniques

10.1 Graphing ‘Familiar’ Functions and Asymptotic boi

Definition

- Lines of Symmetry:** A *line of symmetry* of a function is a line, such that the function is a reflection of itself about that line.
- Horizontal Asymptotes:** A (horizontal) line $g(x) = c$ is the *horizontal asymptote* of the curve $f(x)$ iff $\lim_{x \rightarrow \infty} f(x) = c$ (or with $-\infty$ instead of ∞).^a
- Vertical Asymptotes:** A (vertical) line $x = c$ is a *vertical asymptote* of the curve $f(x)$ iff $\lim_{x \rightarrow c} f(x) = \infty$ or $-\infty$.
- Oblique Asymptotes:** A line $g(x) = mx + c$ — where $m \neq 0$ — is an *oblique asymptote* of the curve $f(x)$ iff $\lim_{x \rightarrow \infty} [f(x) - g(x)] = 0$ (or with $-\infty$ instead of ∞).

^aOtherwise notated by $f(x) \rightarrow c$ as $x \rightarrow \infty$.

Curve Sketching of Rational Functions

S Stationary points

I Intersection with axes

A Asymptotes

- Know how to sketch the graphs of $y = \frac{ax + b}{cx + d}$ and $y = \frac{ax^2 + bx + c}{dx + e}$.
- Rectangular Hyperbolas (of the form $y = \frac{ax + b}{cx + d}$):
 - Two asymptotes, namely $x = -\frac{d}{c}$ and $y = \frac{a}{c}$.
 - Two lines of symmetry with gradients ± 1 and pass through the intersection point of the aforementioned two asymptotes.
- If $n = \deg P = \deg Q$, then
 - $y = R(x)$ is the *horizontal* asymptote of $\frac{P(x)}{Q(x)} = R(x) + \frac{S(x)}{Q(x)}$.
 - Equivalently, $y = \frac{\text{coeff}_P(x^n)}{\text{coeff}_Q(x^n)}$ is a *horizontal* asymptote.^a
- If $\deg P = \deg Q + 1$, then $R(x)$ is an *oblique* asymptote of $\frac{P(x)}{Q(x)} = R(x) + \frac{S(x)}{Q(x)}$.
- Write down asymptotes and lines of symmetry.^b If none are present indicate with “No lines of symmetry.”

^aE.g.: $y = \frac{1}{15}$ is a horizontal asymptote of $y = \frac{1x^2 + 2x - 3}{(5x + 1)(3x + 2)}$.

^bE.g.:

Asymptotes: $x = 4, y = 20$.



Lines of Symmetry: $y = x + 16, y = -x + 24$.

Important Notes

- The discriminant can be very useful.
- Know how to use the G.C. Transfrm app. It allows you to vary the value of some parameter A for a function $f(Ax)$. Use this to graphically find the values of integer k satisfying some conditions.

10.2 Conics

“Tikz is pain, PGFPlots is suffering” — Wise Man.

	Ellipses	Hyperbolas
Standard Forms	$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$	$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$ $\frac{(y-k)^2}{b^2} - \frac{(x-h)^2}{a^2} = 1$
General Equation	$ax^2 + by^2 + cx^2 + dx + e = 0$, where $\text{sgn}(a) = \text{sgn } b$.	$ax^2 + by^2 + cx^2 + dex + e = 0$, where $\text{sgn}(a) \neq \text{sgn } b$.
Center	(h, k)	
Vertical ‘Radius’ (variables here from <i>standard form</i> !)	b	
Horizontal ‘Radius’ (variables here from <i>standard form</i> !)	a	
Vertical Vertices (variables here from <i>standard form</i> !)	$(h, k \pm b)$	
Horizontal Vertices (variables here from <i>standard form</i> !)	$(h \pm a, k)$	
Shape		
Asymptotes (No need to rmb!)	-	$y = k \pm \frac{b(x-h)}{a}$
Lines of Symmetry	$x = h, y = k$	

General Information

- To find asymptote of hyperbolas, just solve

$$\frac{(x-h)^2}{a^2} = \frac{(y-k)^2}{b^2}.$$

- Label vertices or radii, together with the center and asymptotes.

10.3 Parametric Equations


Important Notes

- ★ Check the qns for any *restrictions* on the parameter! And modify that of the G.C.'s accordingly (Tmin & Tmax).
- ★ Vary the t -step or resolution (when using cartesian coordinates) when the graph is oddly jagged.

10.4 Scaling, Translations, and Reflections

Playing With x		
Function	x is replaced with	(Horizontal) Transformation
$f(x + a)$	$x + a$	Translate a units in the positive ($a \leq 0$) O/R negative x -direction ($a \geq 0$).
$f(-x)$	$-x$	Reflect about the y -axis
$f(ax)$	ax	Scale parallel to the x -axis by a scale factor of $\frac{1}{a}$ if $a \geq 0$.
Playing With $f(x)$		
Function / Change to $f(x)$	(Vertical) Transformation	
$f(x) + a$	Translate a units in the positive ($a \geq 0$) O/R negative y -direction ($a \leq 0$).	
$-f(x)$	Reflect about the x -axis.	
$af(x)$	Scale parallel to the y -axis by scale factor a .	

Important Notes



10.5 $|f(x)|$ and $f(|x|)$


General Information

- For $|f(x)|$, simply flip the part of the graph of $f(x)$ that is below the x -axis, to above the x -axis.
- For $f(|x|)$, its graph is symmetric about the y -axis

10.6 $y = \frac{1}{f(x)}$

Behavior of $f(x)$	Behavior of $1/f(x)$
$f(x) > 0$	$\frac{1}{f(x)} > 0$
$f(x) < 0$	$\frac{1}{f(x)} < 0$
Vertical Asymptote at $x = c$	$\frac{1}{f(x)}$ tends to 0 * $\frac{1}{f(x)}$ is undefined at $x = c$
$\frac{df}{dx} = -\frac{d}{dx} \left(\frac{1}{f(x)} \right)$ <p>i.e. when $f(x)$ increases, $\frac{1}{f(x)}$ decreases.</p>	
(a, b) is a <i>minimum</i> pt	$\left(a, \frac{1}{b}\right)$ is a <i>maximum</i> pt
(a, b) is a <i>maximum</i> pt	$\left(a, \frac{1}{b}\right)$ is a <i>minimum</i> pt

Polar Curves



Definition

1. The *pole* is the origin, i.e. the point 0.
2. The *initial line* / *polar axis* is the *half line* $\theta = 0$.






General Information

- Coordinate Conversion

$r = \sqrt{x^2 + y^2}$	$x = r \cos(\theta)$
$\theta = \tan^{-1}\left(\frac{y}{x}\right)$	$y = r \sin(\theta)$

- Standard Functions

Polar Equation	Cartesian Equation
$\begin{array}{c} \theta = \frac{\pi}{2} \\ \left \begin{array}{c} \text{---} \\ \text{---} \end{array} \right. \\ \text{---} \\ \left \begin{array}{c} \text{---} \\ \text{---} \end{array} \right. \\ 0 \end{array}$ <p style="text-align: center;">$r \cos(\theta) = a$</p>	$x = a$

	$y = a$
	$y = x \tan(\alpha)$
	$x^2 + y^2 = a^2$
	$\left(x - \frac{a}{2}\right)^2 + y^2 = \frac{a^2}{4}$
	$x^2 + \left(y - \frac{a}{2}\right)^2 = \frac{a^2}{4}$

- Tangent lines at the pole are obtained by solving $r = 0$.
- Know how to find range of r and θ (given a func/eqn).
- $r = f(\theta)$ is symmetrical about the polar (horizontal) axis iff $f(\theta) = f(-\theta)$.
 - Suppose r is a function of $\cos(n\theta)$ ^a only. Then, the lines of symmetry are $n\theta = 0, \pi, 2\pi, \dots$
- $r = f(\theta)$ is symmetrical about the vertical line $\theta = \pi/2$ iff the equation $f(\theta) = f(\pi - \theta)$.
 - Suppose r is a function of $\sin(n\theta)$ only. Then, the lines of symmetry are

$$n\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \dots, \frac{(2m+1)\pi}{2}, \dots$$

- $r = f(\theta)$ is symmetrical about the pole iff (r, θ) is a point on the curve whenever $(-r, \theta)$ is.
- R -formula may be necessary
- Area of a sector,

$$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta,$$

where $\alpha < \beta$.

- Arc length,

$$\ell = \int_{\theta_1}^{\theta_2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$

^aE.g.: $r = a\sqrt{(4 + \sin^2(\theta))^2 \cos(\theta)}$

Important Notes

1. r is normally ≥ 0 . But, in some questions, it can be negative.
2. No need to fully expand; a final answer such as $(x^2 + y^2)^2 = 3y(x^2 + y^2) - 4y^2$ suffices.
3. Polar curve sketching essentials:
 - (a) Shape of curve
 - (b) Intersection(s) with ('axial') half lines
 - (c) Nothing else *unless* the qns asks for it
 - ☐ Be careful of the sharpness / smoothness of points! Points supposed to be sharp should be sufficiently so, and those supposed to be smooth should be sufficiently rounded / not look sharp.
 - ☐ Check whether the polar axes are tangents at pole. Use this information to draw accurate graphs.
 - ☐ Best to add a small dotted line to show tangentiality at intercepts.
 - ☐ Careful about constants like a in $r = a \sin(\theta)$ for axial intercepts.
 - ☐ No need to state points at the pole unless they are 'axial', i.e. $\theta = 0$, or $\pi/2$, etc.
4. When finding maximum / minimum y values ($dy/dx = 0$), we need to check its nature (1st/2nd deriv test). However, this is unnecessary for max / min r values.
5. For stuff like dy/dx , try to keep it in polar form if possible instead of converting to cartesian form.

6. As usual, be *careful*! E.g. Which values need to be rejected.
7. When reflecting/rotating, a diagram may be useful in finding the necessary angle/expression to replace θ with. E.g.:
 - (a) To reflect about $r = \theta$ or $y = x$, we map $(r, \theta) \rightarrow (r, \pi/2 - \theta)$.
 - (b) Reflect about the half-line $\theta = \pi/2$ is obtained by mapping $(r, \theta) \rightarrow (r, \pi - \theta)$.

G.C. Skills

1. To display a nicely scaled polar curve, we use **Zoom fit**, followed by **Zoom square**
2. Simply press alpha trace 1 to get r_1 . In fact, this works for the other modes available in the GC as well.
3. We can type

$$\left. \frac{d}{d\theta} r_1 \right|_{\theta=\theta}$$

info formulas (like the one for arc length) without having to manually differentiate it!

Conic Sections

Definition 12.1

Eccentricity, e , is defined as

$$\frac{\text{distance of } P \text{ from focus}}{\text{distance of } P \text{ from directrix}}.$$

General Information

◦ Shapes associated with the value of e

- $e = 0$: Circle
- $0 < e < 1$: Ellipse
- $e = 1$: Parabola
- $e > 1$: Hyperbola

Conic	Parabolas		Ellipses		Hyperbolas	
Equation	$x^2 = 4py$	$y^2 = 4px$	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, b > a$	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	$\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$
Foci	$(0, p)$	$(p, 0)$	$(\pm c, 0)$	$(0, \pm c)$	$(\pm c, 0)$	$(0, \pm c)$
a, b, c	N.A.		$c^2 = a^2 - b^2$	$c^2 = b^2 - a^2$	$c^2 = a^2 + b^2$	
Directrices	$y = -p$	$x = -p$	$x = \pm \frac{a}{e} = \pm \frac{a^2}{c}$	$y = \pm \frac{b}{e} = \pm \frac{b^2}{c}$	$x = \pm \frac{a}{e} = \pm \frac{a^2}{c}$	$y = \pm \frac{b}{e} = \pm \frac{b^2}{c}$
e	$e = 1$		$0 < e < 1$		$e > 1$	
	N.A.		$e = \frac{c}{a}$ $= \frac{\sqrt{a^2 - b^2}}{a}$ $= \sqrt{1 - \frac{b^2}{a^2}}$	$e = \frac{c}{b}$ $= \frac{\sqrt{b^2 - a^2}}{b}$ $= \sqrt{1 - \frac{a^2}{b^2}}$	$e = \frac{c}{a}$ $= \frac{\sqrt{a^2 + b^2}}{a}$ $= \sqrt{1 + \frac{b^2}{a^2}}$	$e = \frac{c}{b}$ $= \frac{\sqrt{a^2 + b^2}}{b}$ $= \sqrt{1 + \frac{a^2}{b^2}}$
Reflective Property	When light parallel to its axis of symmetry ($x = 0$ or $y = 0$) hits its concave side, the light is reflected to the focus.		For any point P on the ellipse with $a > b$, $PF_1 + PF_2 = 2a$		For any point P on the hyperbola with $\text{coeff}(x^2) > 0$, $ PF_1 - PF_2 = 2a$	

- Polar Form: $x = p$, $x = -p$, $y = p$, or $y = -p$ being the directrix

Top		
$r = \frac{ep}{1 + e \sin(\theta)}$		
Left		Right
$r = \frac{ep}{1 - e \cos(\theta)}$		$r = \frac{ep}{1 + e \cos(\theta)}$
Bottom		
$r = \frac{ep}{1 - e \sin(\theta)}$		

Definition

- Major / minor axes \implies lengths of longest and shortest diameters respectively.
- Semi-major / semi-minor \implies half of major / minor axes respectively.
- Focal radius \implies distance from point on conic section to focus.

Note

Some possible things to try:

- Using the fact that $PF_1 + PF_2 = 2a$ to do simultaneous equations.
- Converting to polar form (when $e < 1$ so $r \geq 0$) for distances.
- Congruent/Similar triangles.
- Classic use of discriminants.
- Sum and product of roots: Given any polynomial $ax^2 + bx + c$ with the roots α and β ,

$$\alpha + \beta = -\frac{b}{a} \quad \text{and} \quad \alpha\beta = \frac{c}{a}.$$

Functions

General Information

1. Horizontal Line Test:

- (a) Fail: Since^a $y = k$ intersects the graph of $y = f(x)$ more than once, therefore f is not injective.
- (b) Success: Since *any* horizontal line $y = k$ will intersect the graph of $y = g(x)$ *at most once*, so $f(x)$ is one-one.

2. The inverse function, f^{-1} , of a function f exists iff f is one-one.

3. $y = f^{-1}$ is a reflection of $y = f(x)$ about the line $y = x$.

4. The composite function gf exists iff $R_f \subseteq D_g$.

5. $D_{gf} = D_f$ & $R_{gf} = R_g$.

6. Finding the range:

(a) Graphing method:

(b) Mapping method, e.g.: $D_f = (0, \infty) \xrightarrow{f} (1, \infty) \xrightarrow{g} (0, \infty) = R_{gf}$

^asome specific k , e.g. $y = 1/2$

Permutations and Combinations

Definition 14.1

The terms n pick r and n choose r respectively denote

$${}^nP_r := \frac{n!}{(n-r)!} \quad \text{and} \quad \binom{n}{r} = {}^nC_r := \frac{n!}{(n-r)!r!}.$$

General Information

- Addition and multiplication principles
- Know how to ‘bundle’ objects together so as to calculate the total no. of permutations.
- There are

$$\frac{n!}{n_1!n_2!\cdots n_r!}$$

number of ways to arrange n objects, of which n_i are similar, for each i .

Fact

Intuition: If there are n_1 objects are non-distinct out of n objects, then there are $n_1!$ ways to arrange these objects that results in ‘the same’ permutation.

- Case-wise considerations/calculations (then summing together the total number of permutations)
- Unordered circular permutations:
There are $n!/n = (n-1)!$ number of ways of arranging n distinct objects in a circle.

Fact

For unordered circular permutations, we do not care if you rotate the seating arrangement, as long as neighbours are preserved for each object. i.e. $(A, B, C, D) \sim (B, C, D, A)$. As a result, each such collection of n permutations reduces down to one. Thus, explaining the division by n .

- Complementary Method, i.e. taking number of arrangements without restriction - number of arrangements with the opposite of that restriction.

Example 14.1

Number of ways two girls *cannot* sit next to each other = number of arrangements *without restriction* – number of arrangements with girls sitting *together*.

- Insertion Method, place down some of your objects and then insert the rest in the gaps.

Example 14.2

Boys sit at table first: $2!$ ways.

From the 3 gaps, choose 2 for the 2 girls to sit at: 3 ways.

The girls can arrange themselves in $2!$ ways.

So, total no. of ways is $2! \cdot 3 \cdot 2! = 12$.

- Ordered circular permutations: First calculate the number of unordered permutations, then add the ordering at the end.

Note

Circular arrangements are not the same as row arrangements.

We know that A and B are not considered to be seating together in the row arrangement of (A, C, D, E, B) . But, they are seating together in a corresponding row arrangement. The number of row arrangements can be less than, equal to, or more than the number of circular arrangements.

Vectors

Lines	Planes
Equivalent Forms	
<p>1. Vector Equation:</p> $\ell: \mathbf{r} = \mathbf{a} + \lambda \mathbf{m}, \lambda \in \mathbb{R},$ <p>2. Cartesian Equation:</p> $\frac{x - a_1}{m} = \frac{y - a_2}{m_2} = \frac{z - a_3}{m_3}.$	<p>1. Vector Equation:</p> $\Pi: \mathbf{r} = \mathbf{a} + \lambda \mathbf{m}_1 + \mu \mathbf{m}_2 \text{ where } \lambda, \mu \in \mathbb{R},$ <p>2. Scalar Product Form:</p> $\Pi: \mathbf{r} \cdot \mathbf{n} = p$ <p>where the scalar $p := \mathbf{a} \cdot \mathbf{n}$,</p> <p>3. Cartesian Equation:</p> $n_1x + n_2y + n_3z = p$ <p>where the normal vector $\mathbf{n} := (n_1 \ n_2 \ n_3)^\top$.</p>
Foot of Perpendicular	
<p>M1: (a) $\overrightarrow{ON} = \mathbf{a} + \lambda \mathbf{m}$, (b) $\overrightarrow{QN} \cdot \mathbf{m} = 0$, solve for λ, (c) Substitute λ into (a).</p> <p>M2: (a) $\overrightarrow{AN} = (\overrightarrow{AQ} \cdot \hat{\mathbf{m}}) \hat{\mathbf{m}}$, (b) $\overrightarrow{ON} = \overrightarrow{OA} + \overrightarrow{AN}$.</p>	<p>(a) $\ell_{NQ}: \mathbf{r} = \overrightarrow{OQ} + \lambda \mathbf{n}$, where $\lambda \in \mathbb{R}$, and</p> $\Pi: \mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n},$ <p>(b) $(\overrightarrow{OQ} + \lambda \mathbf{n}) \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$, solve for λ,</p> <p>(c) $\overrightarrow{ON} = \overrightarrow{OQ} + \lambda \mathbf{n}$.</p>
Shortest Distance of Point To Line, QN	
<p>M1: $\ \overrightarrow{AQ} \times \hat{\mathbf{m}}\$.</p> <p>M2: (a) $AN = \ \overrightarrow{AQ} \cdot \hat{\mathbf{m}}\$, (b) Pythagoras' Theorem.</p> <p>M3: Using the foot of perpendicular, find distance QN.</p>	<p>M1: $\ \overrightarrow{AQ} \cdot \hat{\mathbf{n}}\$.</p> <p>M2: Distance of plane to <i>origin</i>: If $\Pi: \mathbf{r} \cdot \mathbf{n} = p$, then $\frac{p}{\ \mathbf{n}\ }$ is the shortest distance from the origin to the plane Π. <i>Note:</i></p> <ul style="list-style-type: none"> • If $\frac{p}{\ \mathbf{n}\ } > 0$, then Π 'above' 0. • If $\frac{p}{\ \mathbf{n}\ } < 0$, then Π 'below' 0. <p>M3: Using the foot of perpendicular, then find distance QN.</p>

Relationship Btw 2 Lines		Relationship Btw Line & Plane			
<div>1. Parallel, Non-Intersecting</div> <div>(a) $\mathbf{m}_1 \parallel \mathbf{m}_2$,</div> <div>(b) Solving $\ell_1 = \ell_2$ gives no real solution.</div> <div>2. Parallel, Coinciding</div> <div>(a) $\mathbf{m}_1 \parallel \mathbf{m}_2$,</div> <div>(b) \mathbf{a} lies in ℓ_1 and ℓ_2.</div> <div>3. Non-Parallel, Intersecting</div> <div>(a) \mathbf{m}_1 not $\parallel \mathbf{m}_2$,</div> <div>(b) Solve $\ell_1 = \ell_2$ to find intersection.</div> <div>4. Skew Lines</div> <div>(Non-Parallel, Non-Intersecting)</div> <div>(a) \mathbf{m}_1 not $\parallel \mathbf{m}_2$,</div> <div>(b) Solving $\ell_1 = \ell_2$ gives no real solution.</div>		<div>1. ℓ lies in Π</div> <div>M1: i. $\mathbf{m} \cdot \mathbf{n} = 0$ says $\ell \parallel \Pi$,</div> <div>ii. Combined with $\mathbf{a} \cdot \mathbf{n} = p$, we conclude ℓ lies in Π.</div> <div>M2: Substitute ℓ into Π and show the system (of lin eqns) is consistent for all λ.</div> <div>2. $\ell \parallel \Pi$ but Nonintersecting</div> <div>M1: i. Show $\mathbf{m} \cdot \mathbf{n} = 0$, so $\ell \parallel \Pi$.</div> <div>ii. Then $\mathbf{a} \cdot \mathbf{n} \neq p$, tells us ℓ and Π are nonintersecting.</div> <div>M2: Substitute ℓ into Π, and show the system (of lin eqns) is inconsistent.</div> <div>3. Intersect at 1 point</div> <div>M1: $\mathbf{m} \cdot \mathbf{n} \neq 0$.</div> <div>To find point of intersection: For the plane $\Pi: \mathbf{r} \cdot \mathbf{n} = p$ and ℓ defined by $\ell: \mathbf{r} = \mathbf{a} + \lambda \mathbf{m}$ Solve for λ using simultaneous equations or G.C.</div>			
		-		Relationship Btw 2 Planes	
				<div>1. Parallel Planes: Show there exists an \mathbf{a} for which</div> <div>(a) $\mathbf{a} \cdot \mathbf{n}_1 = p_1$,</div> <div>(b) $\mathbf{a} \cdot \mathbf{n}_2 \neq p_2$.</div> <div>2. Same Plane: Show there exists an \mathbf{a} for which</div> <div>(a) $\mathbf{a} \cdot \mathbf{n}_1 = p_1$,</div> <div>(b) $\mathbf{a} \cdot \mathbf{n}_2 = p_2$.</div> <div>3. Intersect in a line ℓ; To find this line:</div> <div>M1: $\mathbf{n}_1 \times \mathbf{n}_2$ gives the direction vector. So find a common point with simultaneous equations.</div> <div>M2: Solving system of linear equations, from the <i>cartesian</i> form of the planes, using G.C.</div>	
				Point of Reflection	
				<div>1. Find foot of perpendicular \overrightarrow{ON},</div> <div>2. Notice $\overrightarrow{OA'} = \overrightarrow{OA} + 2\overrightarrow{AN} = 2\overrightarrow{ON} - \overrightarrow{OA}$.</div>	
Angle Between					
2 Lines	Line and Plane			2 Planes	
$\theta = \cos^{-1} \widehat{\mathbf{m}}_1 \cdot \widehat{\mathbf{m}}_2 .$	$\theta = \sin^{-1} \widehat{\mathbf{m}} \cdot \widehat{\mathbf{n}} .$	$\theta = \cos^{-1} \widehat{\mathbf{n}}_1 \cdot \widehat{\mathbf{n}}_2 .$			

Probability

General Information

1. Principle of Inclusion and Exclusion for

(a) Two events:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B),$$

(b) Three events:

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C).$$

2. Mutually Exclusive Events:

$$P(A \cap B) = 0,$$

$$P(A \cup B) = P(A) + P(B).$$

3. Independent Events:

$$P(A | B) = P(A),$$

$$P(A \cap B) = P(A)P(B).$$

4. Conditional Probability:

$$P(A | B) = \frac{P(A \cap B)}{P(B)}.$$

5. Use PnC to help compute stuff faster.

6. When we want to find the greatest and least possible probability (e.g. of $P(A^c \cap B^c \cap C^c)$), it is advisable to draw a Venn diagram and fill in all relevant probabilities.

Example 16.1

There are 6 white balls and 5 black balls. Two are randomly drawn. What is the probability that one is white and the other black?

$$\binom{5}{11} \binom{6}{10} + \binom{6}{11} \binom{5}{10} = \frac{6}{11} \quad \text{vs} \quad \frac{\binom{6}{1} \binom{5}{1}}{\binom{11}{2}} = \frac{6}{11}.$$

Differential Equations

17.1 First Order D.E.s

17.1.1 Elementary Solving Techniques

General Information

1. Separable Variables:

$$\frac{dy}{dx} = f(y)g(x), \int \frac{1}{f(y)} dy = \int g(x) dx.$$

2. Integrating Factor:

$$\begin{aligned} \frac{dy}{dx} + P(x)y &= Q(x), \quad \text{let I.F.} = e^{\int P(x) dx} \\ e^{\int P(x) dx} \frac{dy}{dx} + ye^{\int P(x) dx} P(x) &= Q(x)e^{\int P(x) dx}, \\ ye^{\int P(x) dx} &= \int Q(x)e^{\int P(x) dx} dx. \end{aligned}$$

17.1.2 Numerical Methods

General Information

1. Euler's Method:

$$y_{i+1} = y_i + hf(x_i, y_i).$$

Example 17.1

Let (step size) $h = 0.25$ and $f(x, y) = \frac{dy}{dx}$:

$$\begin{aligned} \text{By MF26, } y_2 &= \frac{2}{3} + hf\left(0, \frac{2}{3}\right) \\ &= \frac{13}{18} \\ y_3 &= \frac{13}{18} + hf\left(0.25, \frac{13}{18}\right) \\ &= 0.6701865657. \end{aligned}$$

Therefore, $y(0.5) \approx 0.670$.

1. Improved Euler's Method:

$$i_{i+1} = y_i + hf(x_i, y_i) \quad \& \quad y_{i+1} = y_i + \frac{h}{2}[f(x_i, y_i) + f(x_{i+1}, u_{i+1})].$$

2. Error:

- (a) If $\frac{dy}{dx}$ can be shown to be *increasing* from the calculations of $f(x, y)$, then the curve is *concave upwards*, leading to a *underestimate*.
- (b) If $\frac{dy}{dx}$ can be shown to be *decreasing* from the calculations of $f(x, y)$, then the curve is *concave downwards*, leading to a *overestimate*.

Example 17.2

From the computation, the values of $\frac{dy}{dx}$ increases, i.e. $\frac{d^2y}{dx^2} > 0$, and thus implying the solution curve to be *concave upwards*. Therefore, we have an *underestimation*.

Example 17.3

It is suggested that the estimation in part (ii)^a can be further improved by reducing the step size. Sketch the solution curve and hence comment on this suggestion.

The solution curve has a *stationary point* at $x = 1.47$, which is between 1 and 2 and also the gradient of the curve is close to zero for x value beyond this stationary point. Thus, when the step size is reduced, *tangent* at point close to this stationary point becomes *almost parallel* to the curve, making *little improvement* to the estimation due to *little difference in y* .

^aGiven the point (1,1), we estimated the value of $y(2)$ using the Improved Euler's Method

Example 17.4

It is found that the approximation obtained in (i) for the y -coordinate where $x = 0.75$ is an underestimation and has a percentage error of 120.633%. Explain why there is such a substantial error.

From gradient values calculated above, we suspect sharp changes in gradient values within the interval (from negative to positive). Yet *Euler's Method*^a simply uses a *straight line segment* with gradient^b -4.6409 to estimate the curve for the first iteration, which could have lead to a significant underestimation of the y -value.

^aWe are explaining what it does

^bEmphasising negative gradient (Show its value)

Example 17.5

Compare the relative merits of Euler's Method and the Improved Euler's Method.

Euler's Method: It is computationally simpler.

Improved Euler's Method More accurate as it takes the mean of the initial and next gradient.

17.2 Second Order D.E.

Homogenous	
Roots	Solution y_c
$m_1 \neq m_2$	$y = Ae^{m_1x} + Be^{m_2x}$
$m := m_1 = m_2$	$y = (Ax + B)e^{mx}$
$m = p \pm qi$	$y = e^{px}(A \cos(qx) + B \sin(qx))$
Non-Homogenous , $c_2 \frac{d^2y}{dx^2} + c_1 \frac{dy}{dx} + c_0y = f(x)$	
$y = y_c + y_p$ (C.F. + P.I.)	
$f(x)$	Trial Function for P.I.
Degree n polynomial	$y_p = \sum_{i=0}^n a_i x^i$
ke^{ax}	$y_p = ae^{ax}$
$\alpha \cos(kx) + \beta \sin(kx)$	$y_p = a \cos(kx) + b \sin(kx)$

Note

If y_c and $f(x)$ share some common term, then y_p should be multiplied by x (some least $i \in \mathbb{N}$ times till $x^i y_p$ has no common term with y_c).

Example 17.6

1. If $y_c = A^{-3x}$ and $f(x) = 10e^x$, then $y_p = kxe^x$
2. If $y_c = Ae^x + Be^{-3x}$ and $f(x) = 10e^x$, then $y_p = kxe^x$.
3. If $y_c = Ae^x + Bxe^x + Ce^{-3x}$ and $f(x) = 10e^x$, then $y_p = kx^2e^x$.

17.3 Applications

17.3.1 Exponential Growth

General Information

Let k be the *per-capita growth rate*^a and $P(t)$ be the population at time t . Then we have the model:

$$\frac{dP}{dt} = kP,$$

with the solution

$$P(t) = P_0 e^{kt}.$$

^ai.e. after accounting for births and deaths.

17.3.2 Logistics Growth

General Information

Let k be the *per-capita growth rate*^a, $P(t)$ be the population at time t , and N be the *carrying capacity* of the system. Then we have the model:

$$\frac{dP}{dt} = kP \left(1 - \frac{P}{N} \right).$$

1. Without solving the logistics equation, we can sketch the solution curve by noting the sign of dP/dt :
 - (a) Equilibrium population values occur at $P = 0$ and $P = N$.
 - (b) If, for instance $k > 0$,
 - $0 < P < N$: $1 - \frac{P}{N} > 0$ so $dP/dt > 0$,
 - $P > N$: $1 - \frac{P}{N} < 0$ so $dP/dt < 0$.

“As t increases, the population of _____ increases to the stable population of _____.”

^ai.e. after accounting for births and deaths.

Example 17.7: Neat trick of letting $A = \pm \text{constant}$

$$\begin{aligned}\frac{dP}{dt} &= 3P \left(1 - \frac{P}{200} \right), \\ \int \frac{1}{3P} + \frac{1}{600 - 3P} dP &= \int 1 dt, \\ \ln \left| \frac{3P}{600 - 3P} \right| &= 3t + 3c, \\ \frac{3P}{600 - 3P} &= Ae^{3t}, \text{ where } A = \pm e^{3c}, \\ P &= \frac{200A}{A + e^{-3t}}\end{aligned}$$




Figure 17.1: Logistics curve

17.3.3 Harvesting

General Information

Let k be the *per-capita growth rate*, $P(t)$ be the population at time t , N be the *carrying capacity* of the system, and H the constant *harvesting rate*. Then we have the model:

$$\frac{dP}{dt} = kP \left(1 - \frac{P}{N} \right) - H.$$

1. Bifurcation Point

- (a) When $0 \leq H < \frac{kN}{4}$, there are two equilibrium points, $P = \frac{N}{2} \pm \sqrt{\frac{N^2}{4} - \frac{HN}{k}}$.
- (b) When $H = \frac{kN}{4}$, there is one equilibrium point at $P = \frac{N}{2}$ (the bifurcation point).
- (c) When $H > \frac{kN}{4}$, there is no equilibrium point

2. For Non-Extinction:

$$\frac{N^2}{4} - \frac{HN}{k} \geq 0 \quad \text{and} \quad P_0 \geq 450 - \sqrt{\frac{N^2}{4} - \frac{HN}{k}}.$$

17.3.4 Physics

General Information

MUST rmb the forms.

1. Spring System (where $k > 0$ is the spring constant)

$$\ddot{x} + \frac{k}{m}x = 0.$$

Solution: use **R-formula** to convert to $A \cos(\omega t + \phi)$ where angular frequency $\omega = \sqrt{\frac{k}{m}}$. Period $T = \frac{2\pi}{\omega} = 2\pi\sqrt{\frac{m}{k}}$.

2. Simple Pendulum (where ℓ is its length)

$$\ddot{x} + \frac{g}{\ell}x = 0.$$


Angular frequency $\omega = \sqrt{\frac{g}{\ell}}$ and period $T = 2\pi\sqrt{\frac{\ell}{g}}$.

3. Spring-Mass-Dashpot System (where $c > 0$ is the damping constant)

$$\ddot{x} + \frac{c}{m}\dot{x} + \frac{k}{m}x = 0.$$

Solution

- (a) Real and Distinct Roots: *Overdamped*
- (b) Identical Real Roots: *Critically Damped*
- (c) Complex Conjugate Roots: *Underdamped*
"It will oscillate about the equilibrium position with decreasing amplitude."

**Figure 17.2:** Oscillatory behaviors

Discrete Random Variables

General Information

1. Expectation / Mean,

$$E(X) := \sum_{\text{all } x} x P(X = x).$$

2. Variance

$$\text{Var}(X) := E(X^2) - [E(X)]^2.$$

3. Standard Deviation

$$\sigma := \sqrt{\text{Var}(X)}.$$

4. Properties for two *independent* random variables X and Y ; two *independent observations* X_1 and X_2 of X :

- (a) $E(aX + bY + c) = aE(X) + bE(Y) + c$,
- (b) $E(X_1 + X_2) = E(X_1) + E(X_2) = 2E(X)$.
- (c) $\text{Var}(aX + bY + c) = a\text{Var}(X) + b\text{Var}(Y)$,
- (d) $\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) = 2\text{Var}(X)$.

5. Probability Distribution Table:

x	1	\dots	n
$P(X = x)$	$P(X = 1)$	\dots	$P(X = n)$

Special Discrete Random Variables

Definition 19.1

A discrete random variable X which takes all values in \mathbb{Z}_0^+ is a *binomial distribution* with probability of success p , denoted by $X \sim B(n, p)$, iff

$$P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}.$$

Definition 19.2

A discrete random variable X which takes all values in \mathbb{Z}^+ has a *geometric distribution* with probability of success p , denoted by $X \sim \text{Geo}(p)$, iff

$$P(X = x) = (1-p)^{x-1} p.$$

Note

We can assume $X \sim B(n, p)$ (or $W \sim \text{Geo}(n, p)$) iff the following three conditions hold

1. The event of a [trial in context] is independent of that of another [trial in context].
2. The probability of each [trial in context] is constant.
3. Each trial has only 2 mutually exclusive outcomes.

Note

Defining random variables:

1. Binomial distribution: Let X be the number of [trial in context], out of [number of trials n in context].
2. Geometric distribution: Let W be the number of [trial in context], up to and including the first [successful trial in context].

Note

Let $W \sim \text{Geo}(p)$, and $q := 1 - p$. Then,

$$P(W > m) = q^m \quad \text{and} \quad P(X > m + n \mid X > n) = P(X > m) = q^m.$$

Definition 19.3

A discrete random variable X which takes all values in \mathbb{Z}_0^+ has a *Poisson Distribution* with parameter $\lambda > 0$, denoted by $X \sim \text{Po}(\lambda)$, iff

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}.$$

Note

We can assume $Y \sim \text{Po}(\lambda)$ iff the following three conditions hold

1. The event of a [trial in context] is *independent* of that of another [trial in context].
2. The *mean number of occurrences* of [trial in context] is *constant* over an fixed interval of time/space.
3. The *mean number of occurrences* of [trial in context] is *proportional* to the length of the space/time interval.

Note

Additive property of the Poisson distribution: If $U \sim \text{Po}(\mu)$ and $V \sim \text{Po}(\lambda)$ are *independent* variables, then

$$U + V \sim \text{Po}(\mu + \lambda).$$

Note

Defining random variables: Let Y be the number of [event in context], in [space/time interval in context].

General Information

1. Expectation and Mean:

Distribution	Expectation	Variance
$X \sim \text{B}(n, p)$	np	$np(1 - p)$
$Y \sim \text{Po}(\lambda)$	λ	
$W \sim \text{Geo}(p)$	p^{-1}	$(1 - p)p^{-2}$

2. Use graphing or a table to deal with questions involving inequalities
3. It is helpful to remember the following formulas for when you're asked to derive a formula for mean/mode:

$$\sum_{r=1}^{\infty} rx^{r-1} = (1 - x)^{-2} \quad \text{and} \quad \sum_{r=1}^{\infty} r^2 x^{r-1} = \frac{1 + x}{(1 - x)^3}.$$

4. Why is the probability for (b) is smaller than that for (a): The case of (b) is a proper subset of (a).
5. A discrete random variable M can have other probability distributions. In such cases, defining a random variable W having a Binomial/Poisson/Geometric distribution, and then writing M as a function of W may help.

For example, it may be that $M = W - 1$, or $M = W_1 + W_2$.

Note

When the question asks for the *most likely* number of [event], it is asking for the *mode*.

G.C. Skills

Finding *mode* (e.g. for binomial distributions):

1. Set $Y_1 = \text{binompdf}(n, p, X)$.
2. Go to table.
3. Find the value of X for which the highest value of Y_1 occurs.

G.C. Skills

1. 2nd + Vars + 'A' $\Rightarrow \text{binompdf}(n, p, x) = P(X = x)$
2. 2nd + Vars + 'B' $\Rightarrow \text{binomcdf}(n, p, x) = P(X \leq x)$

Note

Let X be the random variable such that $X \sim \text{B}(n, p)$. If $P(X = n)$ is the *highest probability* that occurs, $X = n$ is the modal value. So, we solve the two inequalities $P(X = 5) > P(X = 4)$ and $P(X = 5) > P(X = 6)$. This gives the *strictest* range of values that p can take (Fig 17.1).




Figure 19.1: In this case, $X = 10$ is the mode.

Example 19.1: 2018 TPJC JC2 H2 MYE P2 8

On average, 3.5% of a certain brand of chocolate turn out misshapen. The chocolates are sold in packets of 25.

- (i) State, in context, two assumptions needed for the number of misshapen chocolates in a packet to be well modelled by a binomial distribution.
- (ii) Explain why one of the assumptions stated in part (i) may not hold in this context.

Answer:

- (i)
 1. Each chocolate is *equally likely* (3.) to be misshapen.
 2. The event that a chocolate is misshapen is *independent* (2.) of the event that another chocolate is misshapen.
- (ii) While on average, the probability that a chocolate is misshapen is 3.5%, it is possible that there are more misshapen chocolates at certain times, possible due to equipment malfunction, which would mean the probability is not constant.

OR

Misshapen chocolates could be the result of equipment used and as the equipment used would not be the same for the same portion of the chocolate produced, whether a chocolate is misshapen may not be independent of another chocolate being misshapen.

Continuous Random Variables

General Information

- A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is a *probability mass function* (pdf) of a continuous random variable X iff f is nonnegative and $\int_{-\infty}^{\infty} f(x) dx = 1$.
- For any probability mass function f , we have $P(a \leq X \leq b) = \int_a^b f(x) dx$. Whether the inequality is strict or nonstrict does not affect the above identity.
- A *mode* of X is any value m such that $f(m)$ is maximum.
- A *cumulative distribution function* (cdf) $F: \mathbb{R} \rightarrow [0, 1]$ of a random variable X is defined by

$$F(x) := P(X \leq x) = \int_{-\infty}^x f(x) dx.$$

- When writing out the cdf as a piecewise function, we explicitly write out the range of values for each case. We reserve the use of “otherwise” for pdf’s.
- Any cdf is continuous and nondecreasing.
- Let X be a continuous random variable with cdf F . To find the pdf g of any $y(X)$, we first find its cdf, then differentiate. We achieve this by reverse engineering $y(X) \leq y$ to find an inequality that relates X with y . E.g. $e^X \leq y$ iff $X \leq \ln(y)$.
- A *median* of X is any value m such that $P(X \leq m) = F(m) = 1/2$.
- Mean/Expectation:

$$\mu = E(X) := \int_{-\infty}^{\infty} x f(x) dx \quad \text{and} \quad E(g(X)) = \int_{-\infty}^{\infty} g(x) f(x) dx.$$

- Important property:

$$E(ag(X) \pm bh(x)) = a E(g(X)) \pm E(h(X)).$$

- Variance:

$$\text{Var}(X) := E(X^2) - [E(X)]^2.$$

- Important property:

$$\text{Var}(aX \pm b) = a^2 \text{Var}(X).$$

Special Continuous Random Variables

Definition 21.1

A continuous random variable X has a *normal distribution* with mean μ and standard deviation σ , denoted by $X \sim N(\mu, \sigma^2)$, iff its pdf f is such that

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

General Information

- A normal distribution is symmetrical about the line $x = \mu$. That is

$$P(X \leq \mu - \delta) = P(X \geq \mu + \delta)$$

for each $\delta > 0$. Note that the mean, median, and mode coincide with μ .

- Properties of the normal distribution. Let X and Y be independent, such that $X \sim N(\mu, \sigma^2)$ and $Y \sim N(m, s^2)$. Then, for any $n \in \mathbb{N}$ and $x, y \in \mathbb{R}$,
 - $nX \sim N(n\mu, n^2\sigma^2)$,
 - $X_1 + X_2 + \cdots + X_n \sim N(n\mu, n\sigma^2)$,
 - $aX \pm bY \sim N(a\mu \pm bm, a^2\sigma^2 + b^2s^2)$.
- At times, the question may be phrased in a misleading manner. Try using some inference to figure out the intended interpretation.

Example 21.1

“The mass of the padding is 30% of the mass of a randomly selected light bulb of mass L . Find the probability that a light bulb with padding has mass c .”

Then for any light bulb of mass L_1 , the mass of the padding is $0.3L_2$ (and *not* $0.3L_1$). i.e. we are to find $P(L_1 + 0.3L_2)$.

- A variable $Z \sim N(0, 1)$ is said to follow the *standard* normal distribution.

Note: Z is reserved for this purpose.

- Let $X \in N(\mu, \sigma^2)$. Then, $\frac{X-\mu}{\sigma}$ follows the standard normal distribution.
- What **Tail** do we select for **invNorm**?


$P(X < x) = p$	LEFT
$P(-x < X < x) = p$	CENTER
$P(X > x) = p$	RIGHT

- When using **invNorm** on an inequality, what should the sign be? For simplicity, we write $\mathcal{L}(p) = \text{invNorm}(p, 0, 1, \text{RIGHT})$, and $\mathcal{R}(p) = \text{invNorm}(p, 0, 1, \text{LEFT})$. Then,

$P(Z > z) \geq p$	$z \leq \mathcal{L}(p)$
$P(Z > z) \leq p$	$z \geq \mathcal{L}(p)$
$P(Z < z) \geq p$	$z \geq \mathcal{R}(p)$
$P(Z < z) \leq p$	$z \leq \mathcal{R}(p)$

Example 21.2

Suppose we want to find the least integer value of m for which $P(Z > 1 - m) \geq 1/2$. Then, using `invNorm (RIGHT)`, we infer that $z \leq 0$, *not* $z \geq 0$. An illustration:

**Definition 21.2**

A continuous random variable X has a *uniform distribution* over the interval (a, b) , which is denoted by $X \sim U(a, b)$, iff its pdf f is such that

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } a < x < b, \\ 0 & \text{otherwise.} \end{cases}$$

Definition 21.3

A continuous random variable Y has an (negative) exponential distribution, which we denote with $Y \sim \text{Exp}(\lambda)$, iff its pdf g is such that

$$g(Y) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

(An exponential distribution models time between occurrences.)

Note

Let $Y \sim \text{Exp}(\lambda)$, then

$$P(Y > z + y \mid Y > y) = P(Y > z).$$

In general, it also holds that

$$P(X > z + y \mid Y > y) = 1 - P(X < z + y \mid Y > y)$$

for any random variable X .

- Expectation and variance:

Distribution	Expectation	Variance
$X \sim U(a, b)$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
$Y \sim \text{Exp}(\lambda)$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$

Note: We need to remember the expectation and variance for the uniform distribution, as it is not provided in the MF26 formula sheet (unlike all other distributions).

- *Warning:* The G.C. tends to incorrectly process an integral if its upper and lower bounds contain $\pm E99$.
- Let T be the time taken between two consecutive arrivals and $\# \sim \text{Po}(\lambda t)$ the number of arrivals in time t . Then,

$$P(T > t) = P(\# = 0) = e^{-4t}.$$

As such, the probability that there is at least one arrival in an interval of time t is

$$P(T \leq t) = 1 - e^{-4t}.$$

Sampling and Estimation

Definition 22.1

A sample is a finite subset of the population.

Definition 22.2

A random sample is a sample selected such that each member of the population has an equal probability of being selected into the sample.

Note

State, in context, what it means for the sample to be random.

It means that every [a member of the population] has an equal probability of being selected into the sample.

Note

Explain why the sample would actually not be random.

[Contextual reason], so not all the [members of the population] have an equal probability of being selected into the sample.

Definition 22.3

Any statistic T derived from a random sample and used to estimate an unknown population parameter θ is known as an *estimator*. It is an *unbiased* estimator iff $E(T) = \theta$. If T is unbiased we commonly write $\hat{\theta}$ for T .

General Information

- Either write $\hat{\mu} \equiv \bar{x} = \dots$ or write out “Unbiased estimate of the population mean μ , $\bar{x} = \dots$ ” Same holds for other population parameters θ .

- Estimators you should know:

Parameter	Estimator	Unbiased?	Formula
Population Mean μ	Sample Mean \bar{X}	✓	$\frac{X_1 + X_2 + \dots + X_n}{n}$
Population Variance σ^2	Sample Variance σ_n^2	×	$\frac{\sum (X_i - \bar{X})^2}{n}$ $\frac{\sum X_i^2}{n} - \bar{X}^2$
	S^2	✓	$\frac{\frac{n}{n-1} \sigma_n^2}{n-1}$ $\frac{\sum (X_i - \bar{X})^2}{n-1}$ $\frac{1}{n-1} \left[\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right]$
Population Proportion p	Sample Proportion P_s	✓	$\frac{X}{n}$

- Let X be a random variable following *any distribution*, and suppose we have a random sample X_1, X_2, \dots, X_n of size $n \geq 50$. Then by CLT (Central Limit Theorem), since $n \geq 50$ is large,

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) \quad \text{and} \quad X_1 + X_2 + \dots + X_n \sim N(n\mu, n\sigma^2)$$

approximately.

- Assumptions when using CLT:
 - The sample is random.
 - Each X_i is independent and identically distributed.
- Suppose $X \sim N(\mu, \sigma^2)$ is known and we pick a *particular* sample. Then,

Distribution	Is An Approximation?
$\bar{X} \sim N(\mu, \sigma^2)$	No
$\bar{X} \sim N(\bar{x}, \sigma^2)$	Yes
$\bar{X} \sim N(\mu, s^2)$	Yes
$\bar{X} \sim N(\bar{x}, s^2)$	Yes

So, if we obtain any of the latter three in solving a question, we must write “ $X \sim N(_, _) \text{approximately}$ ” (even though we knew X *exactly* follows a normal distribution!)

- Pooled estimators. First assume we have two populations, from which we select a random sample of size n_1 and n_2 . We let \bar{X}_1 and S_1^2 denote the sample mean and unbiased estimator for variance, respectively, for the first sample. Similarly define \bar{X}_2 and S_2^2 , for the second sample.

Parameter	Unbiased Pooled Estimator
Mean	$\hat{\mu} = \frac{n_1 \bar{X}_1 + n_2 \bar{X}_2}{n_1 + n_2}$
Variance	$S_p^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}$

The following definition is found in [Hogg-McKean-Craig](#). Similar definitions are also found in [Wackerly-Mendenhall-Schaefer](#) and [Nitis Mukhopadhyay](#).

Definition 22.4

Let X_1, X_2, \dots, X_n be a sample on a random variable X , where X has pdf $f(x; \theta)$, $\theta \in \Omega$. Let $0 < \alpha < 1$ be specified. Let $L = L(X_1, X_2, \dots, X_n)$ and $U = U((X_1, X_2, \dots, X_n))$ be two statistics. We say that the interval (L, U) is a $(1 - \alpha)100\%$ *confidence interval* for θ iff

$$1 - \alpha = P_\theta[\theta \in (L, U)].$$

That is, the probability that the interval contains θ is $1 - \alpha$, which is called the *confidence coefficient* or *confidence level* of the interval.

- We cannot write “a $1 - \alpha$ (e.g. 0.95) confidence interval”. The $1 - \alpha$ must always be expressed as a *percentage*.
- Let $\hat{\theta}$ be a statistic that is normally distributed with mean θ and standard error $\sigma_{\hat{\theta}}$. We see that

$$\frac{\hat{\theta} - \theta}{\sigma_{\hat{\theta}}} = Z \sim N(0, 1).$$

Rewriting $P(-z_{1-\alpha/2} < Z < z_{1-\alpha/2}) = 1 - \alpha$ gives

$$P(\hat{\theta} - z_{1-\alpha/2}\sigma_{\hat{\theta}} < \theta < \hat{\theta} + z_{1-\alpha/2}\sigma_{\hat{\theta}}) = 1 - \alpha.$$

Hence, a $(1 - \alpha)100\%$ confidence interval for θ is

$$(\hat{\theta} - z_{1-\alpha/2}\sigma_{\hat{\theta}}, \hat{\theta} + z_{1-\alpha/2}\sigma_{\hat{\theta}}).$$

(Wackerly-Mendenhall-Schaefer)

- Let $0 < \alpha < 1$ and X_1, X_2, \dots, X_n be a sample on a random variable X with mean μ , where n is large. Then, an approximate $(1 - \alpha)100\%$ confidence interval for μ is

$$\left(\bar{x} - z_{1-\alpha/2} \frac{s}{\sqrt{n}}, \bar{x} + z_{1-\alpha/2} \frac{s}{\sqrt{n}} \right).$$

When the variance σ^2 is known, we can replace s with σ . If the distribution of X is known to be normal, in addition to σ^2 being known exactly, then the confidence interval is exact; it is not just an approximation.

(Hogg-McKean-Craig)

- Let X be a Bernoulli random variable with probability of success p , where X is 1 or 0 if the outcome is success or failure, respectively. Suppose X_1, X_2, \dots, X_n is a random sample from the distribution of X , where n is large. Let $\hat{p} = \bar{X}$ be the sample proportion of successes. Then, an approximate $(1 - \alpha)100\%$ confidence interval for p is given by

$$\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right).$$

(Letting $Y = X_1 + X_2 + \dots + X_n \sim B(n, p)$ gives $\hat{p} = Y/n$, which is the presentation used in the school's notes.)

(Hogg-McKean-Craig)

Note

Standard phrasing for the interpretation of a $(1 - \alpha)100\%$ confidence interval (a, b) .

The probability that the interval (a, b) contains the true value of the [population mean/proportion in context] is $1 - \alpha$.

Note

Standard phrasing for what is a $(1 - \alpha)100\%$ confidence interval for θ ?

It is an interval which has probability $1 - \alpha$ of containing the true value of θ .

Note

Standard phrasing for whether mean/proportion in context has likely increased/decreased, when given suitable confidence intervals.

1. There is no conclusive result.

Since the old and new $(1 - \alpha)\%$ confidence intervals overlap, we are unable to conclude whether the [mean/proportion in context] has decreased or not. Hence, it is inconclusive from these figures as to whether the [context (e.g. an awareness campaign)] has been effective.

2. It has likely increased/decreased.

The old $(1 - \alpha)\%$ confidence interval is to the left/right of the new $(1 - \alpha)\%$ confidence interval, such that they do not overlap. So, can conclude that the [mean/proportion in context] likely increased/decreased. Hence, these figures suggests that the [context (e.g. an awareness campaign)] has been effective.

Note

Advantage and disadvantage of a $(1 - \beta)\%$ confidence interval compared to a $(1 - \alpha)\%$ confidence interval, where $\beta < \alpha$.

Advantage: A $(1 - \beta)\%$ CI is more likely to contain the true mean.

Disadvantage: A $(1 - \beta)\%$ CI is less precise (or wider).

Note. Clearly state which is the advantage and disadvantage, as illustrated above.

G.C. Skills

Calculating statistics (i.e. \bar{x} , s , etc) by G.C. given data for a sample.

1. Keying in the data: **stat** \Rightarrow **1:Edit** \Rightarrow Key in the data into one of the lists L_i .
2. Calculating the statistic: **stat** \Rightarrow **CALC** \Rightarrow **1-Var Stats (List: L_i)** \Rightarrow **Calculate**.
3. Getting the statistic for further calculations: **vars** \Rightarrow **5:Statistics** \Rightarrow Select the desired statistic.

G.C. Skills

Calculating the symmetric confidence interval by G.C.

Mean: **stat** \Rightarrow **TESTS** \Rightarrow **7:ZInterval...**

Proportion: **stat** \Rightarrow **TESTS** \Rightarrow **A:1-PropZInt...**

Statistics: Hypothesis Testing

Definition 23.1

The *null hypothesis* H_0 and *alternative hypothesis* H_1 are the hypotheses that we hope to reject and accept, respectively.

General Information

- Without going into details, a *critical region* C is just a set that defines the decision rule / test

$$\text{Reject } H_0 \text{ (Accept } H_1) \quad \text{if } (X_1, X_2, \dots, X_n) \in C,$$

for any random sample X_1, X_2, \dots, X_n from the distribution of a random variable X .

Definition 23.2

The *significance level* $100\alpha\%$ of a test is the probability of rejecting H_0 when it is in fact true. i.e. $\alpha = P(H_0 \text{ is rejected} \mid H_0 \text{ is true})$.

Note

Explain, in context, the meaning of ‘at the $\alpha\%$ level of significance’.

The probability that [H_1 in context], when actually [H_0 in context], is $\alpha\%$.

Definition 23.3

The *p-value* is the lowest level of significance for which the null hypothesis will be rejected. In other words, for the null hypotheses

$$(a) \mu < \mu_0, \quad (b) \mu \neq \mu_0, \quad (c) \mu > \mu_0,$$

we have

$$(a) p\text{-value} = P(Z \leq z_{\text{calc}}), \quad (b) p\text{-value} = P(|Z| \geq |z_{\text{calc}}|), \quad (c) p\text{-value} = P(Z \geq z_{\text{calc}}).$$

Note

Explain what the *p-value* means in context.

The *p-value* is the least level of significance to conclude that [H_1 in context].

- A large sample hypothesis test for the mean.

- Let [X in context] and μ be the population mean.

- Test against H_1 : (a) $\mu < \mu_0$, (b) $\mu \neq \mu_0$, or (c) $\mu > \mu_0$, at the $100\alpha\%$ significance level.

- Under H_0 , we have $\bar{X} \sim N(\mu_0, \hat{\sigma}^2/n)$ approximately. Or, if σ^2 is known exactly, then by CLT $\bar{X} \sim N(\mu_0, \sigma^2/n)$ approximately.

- Test statistic:

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1).$$

4. Find $z_{1-\alpha}$ or $z_{1-\alpha/2}$, which satisfies

(a) $P(Z < z_{1-\alpha}) = \alpha$,

(b) $P(-z_{1-\alpha/2} < Z < z_{1-\alpha/2}) = \alpha$,
or

(c) $P(Z > z_{1-\alpha})$.

5. Find the test statistic value

$$z_{\text{calc}} = \frac{\hat{\mu} - \mu_0}{\sigma/\sqrt{n}}.$$

6. Reject H_0 iff

(a) $z_{\text{calc}} < z_{1-\alpha}$,

(b) $|z_{\text{calc}}| > z_{1-\alpha/2}$, or

(c) $z_{\text{calc}} > z_{1-\alpha}$.

7. Since (a) $z_{\text{calc}} < z_{1-\alpha}$, (b) $|z_{\text{calc}}| > z_{1-\alpha/2}$, (c) $z_{\text{calc}} > z_{1-\alpha}$, or $p\text{-value} < \alpha$, we reject H_0 . There is sufficient evidence at the significance level $100\alpha\%$ that [H_1 in context].

Note. For *not* rejecting H_0 , simply change to the appropriate inequality (such that z_{calc} is outside the critical region) and write “insufficient” instead of “sufficient”.

- If we have a null hypothesis, such as

$$H_0: \mu \leq \mu_0 \quad \text{or} \quad H_0: \mu \geq \mu_0,$$

we can just use $H_0: \mu = \mu_0$ instead.

4. Find the p -value using GC.

5. Reject H_0 iff p -value is less than α .

G.C. Skills

Calculating the p -value of a sample.

`stat` \Rightarrow TESTS \Rightarrow 1:Z-Test.

Note

Explain why there is no need to assume that the distribution of X is normal/know anything about the population distribution of X .

As the sample size n is large, by the Central Limit Theorem, the sample mean of [random variable X in context] will approximately follow a normal distribution.

Note. Spell “Central Limit Theorem” and “the sample mean” out *in full*. Do not use CLT or \bar{X} for this question.

Correlation and Linear Regression

Note

A good scatter diagram should follow the guidelines below.

- The relative position of each point on the scatter diagram should be clearly shown.
- The range of values for the set of data should be clearly shown by marking out the extreme x and y values on the corresponding axis.
- The axes should be labeled clearly with the variables.


General Information

- The Product Moment Correlation Coefficient is a measure of the linear correlation between two variables. It is defined by

$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sqrt{\sum (x - \bar{x})^2 \sum (y - \bar{y})^2}} = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sqrt{\left[\sum x^2 - \frac{(\sum x)^2}{n} \right] \left[\sum y^2 - \frac{(\sum y)^2}{n} \right]}}$$

which takes on a value from 0 to 1.

- When $r = 0$, there is no linear relationship. But, a nonlinear relationship may be present. Additionally, the regression lines are perpendicular.
- The closer the value of r is to 1 (or -1), the stronger the positive (or negative) linear correlation. Furthermore, the regression lines coincide.



- The regression line of y on x minimises the sum of squares deviation (error) in the y -direction. (i.e. we are assuming x is the independent variable whose values are known exactly.) It is given by

$$y = \bar{y} + b(x - \bar{x}), \quad \text{where} \quad b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2} = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sum x^2 - \frac{(\sum x)^2}{n}}.$$

- The point (\bar{x}, \bar{y}) always lies on both the regression lines of y on x , and x on y .
- Say we are given the value of one variable, and asked to approximate the value of the other variable. Then, we should always use the line of the *dependent* variable on the *independent*.
- Estimations should not be taken for data outside the range of the sample provided, even if the value of r is close to 1.

Bibliography

1. Fig 6.1 Trapezium rule <https://tex.stackexchange.com/a/110618>
2. Fig 6.2 Simpson's rule <https://tex.stackexchange.com/a/439119>
3. Fig 7.1 Argand Diagram <https://tex.stackexchange.com/a/466846>
4. Oscillatory behavior of DEs modelling physical phenomena Fig 15.2 https://tikz.net/dynamics_oscillator/
5. Fig 17.1 Mode of a binomial distribution <https://math.oxford.emory.edu/site/math117/normalApproxToBinomial/>
6. Product moment correlation https://www.ncl.ac.uk/webtemplate/ask-assets/external/maths-resources/images/R_value.png
7. Used some inspiration from the beautiful preamble by tearfox and det.uwu from Discord, to make my environments look better.