# A-Level Mathe

## Shao Hong

## Contents

1	<b>A1:</b>	Inequalities and Equations
	1.1	Solving Inequalities
		1.1.1 Rational Inequalities
	1.2	Modulus Inequalities
<b>2</b>	B1(.	A): Graphing Techniques (Part I)

### A1: Inequalities and Equations

#### 1.1 Solving Inequalities

#### 1.1.1 Rational Inequalities

### General Methods

- 1. Quadratic formula for factorisation / finding roots (of polynomial).
- 2. Completing the square.
- 3. Discriminant/Completing the Sq to eliminate\* factors which are always positive or negative (e.g. removing  $x^2 - 3x + 4$ ). Note to include coefficient of  $x^2$  in argument.
- 4. GC (include sketch).
- 5. Rational Functions<sup>a</sup>: Move everything to one side (+,-), then use number line.
- 6. Number line (more complicated functions).

### Important Notes

- $\Box$  Eliminating Factors only a works for c = 0 in  $f(x) \ge c$  or  $f(x) \le c$ .
- $\Box$  Discriminant include coefficient of  $x^2$  in argument.
- □ Rational functions exclude the values that causes division by zero to occur.
- $\Box$  Cross multiplication preserves order for  $\frac{x}{y} < \frac{x'}{y'}$  iff y and y' are both positive or negative.
- $\Box$  Squaring preserves order for x < y iff x and y are both positive or negative.

#### Modulus Inequalities 1.2

Fact. Given  $x \in \mathbb{R}$ , we have that

- $|x| \geq 0$ ,
- $|x^2| = |x|^2 = x^2$ ,
- $\bullet \ \sqrt{x^2} = |x|.$

And as long as  $x \in \mathbb{R}^+$ ,

$$\bullet \ \sqrt{x}^2 = |x|.$$

<sup>&</sup>lt;sup>a</sup>Fractions of Polynomials

<sup>&</sup>lt;sup>a</sup>Counterexample:  $P(x) = x(3x^2 - 9x + 10) \le 2$  iff  $x \le 2$  is false. E.g.:  $P(1.8) = 6.336 \le 2$ . <sup>b</sup>Otherwise, note the counterexample  $\frac{1}{2} < \frac{1}{-3}$ .

### Useful Properties

For every  $x, k \in \mathbb{R}$ :

- (a) |x| < k iff -k < x < k.
- (b) |x| > k iff x < -k or x > k.

(of course, similarly applies for the non-strict ordering  $\leq)$ 

<sup>a</sup>Notice that k > 0 here since  $0 \le |x| < k$ .

### Important Notes

 $\circ$  Note that when solving for |x| = y, |x| < y, etc, y must be greater than or equal to 0. In other words, there may be solutions we will need to reject.

(For <, equality is of course not allowed.)

## B1(A): Graphing Techniques (Part I)

### General Definitions

- 1. **Lines of Symmetry**: A *line of symmetry* of a function is a line, such that the function is a reflection of itself about that line.
- 2. Horizontal Asymptotes: A (horizontal) line g(x) = c is the horizontal asymptote of the curve f(x) iff  $\lim_{x\to\infty} f(x) = c$  (or with  $-\infty$  instead of  $\infty$ ).
- 3. Vertical Asymptotes: A (vertical) line x = c is a vertical asymptote of the curve f(x) iff  $\lim_{x\to c} f(x) = \infty$  or  $-\infty$ .
- 4. **Oblique Asymptotes**: A line g(x) = mx + c where  $m \neq 0$  is an *oblique asymptote* of the curve f(x) iff  $\lim_{x\to\infty} [f(x)-g(x)]=0$  (or with  $-\infty$  instead of  $\infty$ ).

<sup>a</sup>Otherwise notated by  $f(x) \to c$  as  $x \to \infty$ .