

# Day 4

## Outline

1. Statistical Terminology
2. Sampling Distributions

## Statistical Terminology

### Tidy Data

Each column represents a variable

Header row will contain the name of the variable

Each row represents a case

Each value goes in its own cell.

**Good form:** left most column contains label variable whose values are unique IDs for the cases

One row could represent:

- a patient
- a particular test for that patient
- all patients seen by a doctor

Merging datasets: you need to pair like data

### Data Dictionary

For each variable:

- name of the variable
- type of the variable
- units of measurement
- description

### Type of Variable

Numerical (Quantitative): int, float, double

Categorical (Qualitative): string, classes, char, software-specific variable type

Typically, we do not select only one case from the population. We instead select a subset of the population: sample. A sample will always exist in the real world.

## Statistical Terminology (Continued)

Variables vary between cases.

Statistics vary between samples

Parameters vary between populations.

## Frequentist Statistics

Parameters are constants, but we don't know their value.

Statistics are random variables that describe “randomly select a sample of some fixed size, record values of a variable for each case in the sample, and summarize the value”.

## In this class

Numerical variables: we use  $\mu$  to represent population mean and  $\bar{X}$  to represent sample mean

Categorical variables: we use “p” to represent population proportion of outcome in a particular category.

We use  $\hat{p}$  to represent sample proportion of outcomes in a particular category.

## Example

A clinical trial compares two bladder cancer drugs:

- Drug A (Company's "new" drug)
- Drug B (Current best drug)

They recruit 200 subjects with bladder cancer and assign 100 to take Drug A and 100 to take Drug B

## Questions

- What is the case
- We can consider this study to have 2 "hypothetical" populations. What are they?
- What are the two samples from those hypothetical populations? (subset of a larger group/population)
- Name an outcome the drug company might be interested in. Is that outcome a numerical or categorical variable?
- What statistic might we use to summarize that outcome in a sample
- What is the corresponding parameter in the hypothetical population.

## Answers

- A case is one patient with bladder cancer
- Everyone on Drug A (all bladder cancer patients, if they took Drug A) and everyone on Drug B
- 100 people who took Drug A and 100 people who took Drug B
- How much more effective is Drug A compared to Drug B. This would yield a numerical value. (Reduction in tumor size, cancer remission/not in remission)
- Sample mean ( $\bar{X}$ ) tumor reduction. Sample proportion/sample percent of patients who are in remission.
- Population mean tumor reduction and population proportion in remission

## Sampling Distribution

These are things that do not have real world equivalences.

The probability distribution of a statistic is its sampling distribution

Distribution of a statistic over all possible samples of a given size from the sample population. **Must** specify size of sample.

To find a sampling distribution:

1. Simulation: approximate the sampling distribution by simulating samples.
2. Asymptotic behavior: as the number of samples  $\rightarrow \infty$ , what does the distribution look like?

## Properties of Sampling Distributions

Let  $X$  be the statistic we use to estimate a parameter  $\theta$  for a population.  $X$  is an unbiased estimator of  $\theta$  if  $\mu_x = \theta$ . Otherwise  $X$  is biased and the amount of **bias** is  $\mu_x - \theta$ .

The variability of  $X$  describes the amount by which individual realizations of  $X$  are “spread” about  $\mu_x$ .

We can summarize variability by variance, standard deviation, standard error, margin of error