

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the same field \mathbb{F} .

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.
2. Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$, let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be defined by $T(x) = Ax$, $x \in \mathbb{R}^n$. Then T is a linear transformation.

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.
2. Let $A \in \mathbb{M}_{m \times n}(R)$, let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be defined by $T(x) = Ax$, $x \in \mathbb{R}^n$. Then T is a linear transformation.
3. Let $T : C[a, b] \rightarrow \mathbb{R}$ be defined by $T(f) = \int_a^b f(x)dx$. Then T is a linear transformation.

-
1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.

-
1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
 2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.
3. Let \mathbb{V} be a vector space over the field \mathbb{F} and let $\lambda \in \mathbb{F}$. Let

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.
3. Let \mathbb{V} be a vector space over the field \mathbb{F} and let $\lambda \in \mathbb{F}$. Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = \lambda x, x \in \mathbb{V}$. This transformation is called **scalar transformation**.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

$T : \mathbb{R} \rightarrow \mathbb{R}$ be a map defined by $T(x) = x + 1$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

$T : \mathbb{R} \rightarrow \mathbb{R}$ be a map defined by $T(x) = x + 1$. Using above theorem you can say that T is not linear.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^n \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

$$T(x) = \sum_{i=1}^n \alpha_i x_i$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

$$T(x) = \sum_{i=1}^n (\alpha_1^i, \dots, \alpha_m^i) x_i$$

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{i=1}^n x_i \alpha_i^i$$

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

Could it be possible to get the linear map explicitly?

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

Could it be possible to get the linear map explicitly?

Answer: Yes. Let $(x_1, x_2) \in \mathbb{R}^2$. Then $(x_1, x_2) = x_1 e_1 + x_2 e_2$.

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

Could it be possible to get the linear map explicitly?

$$\begin{aligned}\text{Then } T(x_1, x_2) &= x_1 T(e_1) + x_2 T(e_2) \\ &= x_1(1, 1) + x_2(-1, 1) \\ &= (x_1 - x_2, x_1 + x_2)\end{aligned}$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

Could it be possible to get the linear map explicitly?

Answer: No it is not possible.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} .

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} .

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$.

Define $T(x) = \sum_{i=1}^n c_i w_i$. It is clear that T is well defined because $x = \sum_{i=1}^n c_i u_i$, this expression unique.

We first show that T is a linear transformation. Take $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$ and $y = \sum_{i=1}^n d_i u_i$.

We first show that T is a linear transformation. Take $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$ and $y = \sum_{i=1}^n d_i u_i$.

Let $\alpha, \beta \in \mathbb{F}$. $T(\alpha x + \beta y) = T(\sum_{i=1}^n (\alpha c_i + \beta d_i) u_i)$.

$$\begin{aligned} T(\alpha x + \beta y) &= \sum_{i=1}^n (\alpha c_i + \beta d_i) w_i. \\ &= \alpha \sum_{i=1}^n c_i w_i + \beta \sum_{i=1}^n d_i w_i. \\ &= \alpha T(x) + \beta T(y). \end{aligned}$$

Hence T is linear.

Uniqueness: Suppose that there is another linear transformation U such that $U(u_i) = w_i$.

Uniqueness: Suppose that there is another linear transformation U such that $U(u_i) = w_i$.

To show that $U = T$. Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n a_i u_i$. Using definition of T

$$\text{we have } T(x) = T\left(\sum_{i=1}^n a_i u_i\right) = \sum_{i=1}^n a_i w_i.$$

$$U(x) = U\left(\sum_{i=1}^n a_i u_i\right)$$

$$= \sum_{i=1}^n a_i U(u_i) \text{ (applying the definition of linear transformation)}$$

$$= \sum_{i=1}^n a_i w_i.$$

Then $U(x) = T(x)$ for all $x \in \mathbb{V}$. Hence $U = T$.

- [Example]

Take the basis $\{e_1, e_2, e_3\}$ in \mathbb{R}^3 . Take $1, 2, 3 \in \mathbb{R}$. Then using previous theorem we have a unique linear transformation T from \mathbb{R}^3 to \mathbb{R} such that $T(e_1) = 1$, $T(e_2) = 2$, $T(e_3) = 3$ and $T(x_1, x_2, x_3) = x_1 + 2x_2 + 3x_3$.

Take the basis $\{e_1, e_2, e_3\}$ in \mathbb{R}^3 . Take $1, 2, 3 \in \mathbb{R}$. Then using previous theorem we have a unique linear transformation T from \mathbb{R}^3 to \mathbb{R} such that $T(e_1) = 2$, $T(e_2) = 1$, $T(e_3) = 3$ and $T(x_1, x_2, x_3) = 2x_1 + x_2 + 3x_3$. This transformation is different between the previous transformation.

The previous theorem gives a technique to construct a linear transformation from a finite dimensional vector space to another dimensional vector space over the same field \mathbb{F} .

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspaces $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$.

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$. you can easily check that $R(T)$ is a subspace of \mathbb{W} .

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$. you can easily check that $R(T)$ is a subspace of \mathbb{W} .

The subspace $R(T)$ is called the **range space** of T .

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

$$\implies N(T) := \text{ls}(\{(1, 1, 1)\})$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

$$\implies N(T) := \text{ls}(\{(1, 1, 1)\})$$

$$\dim(N(T)) = 1.$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \text{ls}(\{(1, 0), (0, 1)\})$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \text{ls}(\{(1, 0), (0, 1)\})$$

$$\dim(R(T)) = 2$$

-
- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the field \mathbb{F} .

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the field \mathbb{F} .

The $\dim(R(T))$ is called the **rank** of T and $\dim(N(T))$ is called the **nullity** of T .

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$. This implies $T(x_1 - x_2) = 0$. Hence $x_1 - x_2 \in \text{Ker}(T)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$. This implies $T(x_1 - x_2) = 0$. Hence $x_1 - x_2 \in \text{Ker}(T)$. Therefore $x_1 - x_2 = 0$. This implies $x_1 = x_2$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } T(u_1), \dots, T(u_n) \text{ are LI.}$$

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } T(u_1), \dots, T(u_n) \text{ are LI.}$$

Converse is not true in general. That is if u_1, \dots, u_n are LI, then $T(u_1), \dots, T(u_n)$ may or may not be LI.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$$c_i u_1 + \dots + c_n u_n = 0_{\mathbb{V}} \text{ as } T \text{ is one-one.}$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$$c_i u_1 + \dots + c_n u_n = 0_{\mathbb{V}} \text{ as } T \text{ is one-one.}$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } u_1, \dots, u_n \text{ are LI}$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$T(x) = T(c_1 u_1 + \dots + c_n u_n)$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$\begin{aligned} T(x) &= T(c_1 u_1 + \dots + c_n u_n) \\ &= c_1 T(u_1) + \dots + c_n T(u_n) \end{aligned}$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$T(x) = T(c_1 u_1 + \dots + c_n u_n)$$

$$= c_1 T(u_1) + \dots + c_n T(u_n) = c_{k+1} T(u_{k+1}) + \dots + c_n T(u_n)$$

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Since $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$ is basis of \mathbb{V} . Then $a_1 = \dots = a_{n-k} = 0$.

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Since $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$ is basis of \mathbb{V} . Then $a_1 = \dots = a_{n-k} = 0$.

Therefore $T(u_{k+1}), \dots, T(u_n)$ are LI. Hence $\{T(u_{k+1}), \dots, T(u_n)\}$ is a basis of $R(T)$. Then $\dim(R(T)) = n - k$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Let $\mathbb{S} = \text{ls}(\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\})$. So \mathbb{S} is a subspace of \mathbb{V} and $\dim(\mathbb{S}) = m + k + 1$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Let $\mathbb{S} = \text{ls}(\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\})$. So \mathbb{S} is a subspace of \mathbb{V} and $\dim(\mathbb{S}) = m + k + 1$.

$T_{\mathbb{S}}$ is a LT from \mathbb{S} to \mathbb{W} . Then $Ker(T_{\mathbb{S}}) \subseteq Ker(T)$ and $R(T_{\mathbb{S}}) \subseteq R(T)$.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

1. T is one-one.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

1. T is one-one.
2. T is onto.

-
- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) < \dim(\mathbb{W})$. Then there is no onto linear transformation from \mathbb{V} to \mathbb{W} .

- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) < \dim(\mathbb{W})$. Then there is no onto linear transformation from \mathbb{V} to \mathbb{W} .
- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) > \dim(\mathbb{W})$. Then there is no one-one linear transformation from \mathbb{V} to \mathbb{W} .