

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

If $x = (0, 0, \dots, 0)$, then $x \cdot x = 0$.

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

If $x = (0, 0, \dots, 0)$, then $x \cdot x = 0$.

$$\text{ii) } x \cdot y = \sum_{i=1}^n x_i y_i = \sum_{i=1}^n y_i x_i = y \cdot x.$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.
- How do we understand the geometry of an arbitrary vector space?

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.
- How do we understand the geometry of an arbitrary vector space?
- Inner product can be used to understand the geometry of abstract vector spaces.

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

An inner product is a mapping $\langle ., . \rangle : \mathbb{V} \times \mathbb{V} \rightarrow \mathbb{K}$ which satisfies the following conditions.

-
1. $\langle x, x \rangle \geq 0$ for all $x \in V$ (**positivity**)

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

4. $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$ for all $x, y, z \in \mathbb{V}$ (**additivity**).

Examples:

- Dot product on \mathbb{R}^n is an inner product.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_i^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Ans: This is an inner product.

- $V = C[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in C[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

- $\mathbb{V} = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

$$\text{Then } \langle \alpha f, g \rangle = \int_a^b (\alpha f(x))g(x)dx = \alpha \int_a^b f(x)g(x)dx = \alpha \langle f, g \rangle.$$

4. Let $f, g, h \in \mathbb{C}[a, b]$.

4. Let $f, g, h \in \mathbb{C}[a, b]$.

$$\begin{aligned}\text{Then } \langle f + g, h \rangle &= \int_a^b (f(x) + g(x)) h(x) dx \\ &= \int_a^b (f(x)h(x) + g(x)h(x)) dx \\ &= \langle f, h \rangle + \langle g, h \rangle.\end{aligned}$$

- Let $f : [a, b]$ be integrable and $f(x) \geq 0$. Let f be continuous at $x = c$ and $f(c) > 0$ (resp. $f(c) < 0$). Then $\int_a^b f(x)dx > 0$ (resp. $\int_a^b f(x)dx < 0$).

Sol: Case I. Let $a < c < b$.

f is continuous at $x = c$. Take $\epsilon = \frac{f(c)}{2}$ then there exists $\delta > 0$ such that

for each $x \in (c - \delta, c + \delta) \implies |f(x) - f(c)| < \frac{f(c)}{2}$.

Then $-\frac{f(c)}{2} < f(x) - f(c) < \frac{f(c)}{2}$ for all $x \in (c - \delta, c + \delta)$.

Therefore $f(x) > \frac{f(c)}{2} > 0$ for all $x \in (c - \delta, c + \delta)$.

$$\int_a^b f^2(x)dx = \int_a^{c-\delta} f(x)dx + \int_{c-\delta}^{c+\delta} f(x)dx + \int_{c+\delta}^b f(x)dx > 0$$

as $\int_{c-\delta}^{c+\delta} f^2(x)dx > 0$.

Case II. $c = a$

When $c = a$, we have a neighborhood $(a, a + \delta)$ where f is positive.

$$\int_a^b f^2(x) dx = \int_a^{a+\delta} f(x) dx + \int_b^{a+\delta} f(x) dx > 0$$

Case-III. $c = b$. Same as Case II.

To show $\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0$.

Here f^2 is nonnegative function and continuous. If f is not identically zero, then there exists $c \in [a, b]$ such that $f(c) \neq 0$. Therefore $f^2(c) > 0$. Using above result we have $\int_a^b f(x)f(x)dx > 0$, a contradiction. Hence $f \equiv 0$.

-
- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

- $V = C(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in C(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in C(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

Hence $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ is not an inner product on $C(\mathbb{R})$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, k\}$ be a basis of \mathbb{V} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

We now show that \langle, \rangle is an IP on \mathbb{V} .

-
1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

Therefore $x = \sum_{i=1}^k a_i u_i = 0$.

2. Let $x, y \in \mathbb{V}$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, y \rangle = \sum_{i=1}^k a_i \overline{b_i} = \overline{\sum_{i=1}^k \overline{a_i} b_i} = \overline{\langle y, x \rangle}.$$

3. This part is same as part 2.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Then $x + y = \sum_{i=1}^k (a_i + b_i) u_i$. Therefore,

$$\begin{aligned}\langle x + y, z \rangle &= \sum_{i=1}^k (a_i + b_i) \overline{c_i} \\ &= \sum_{i=1}^k (a_i \overline{c_i} + b_i \overline{c_i}) \\ &= \langle x, z \rangle + \langle y, z \rangle.\end{aligned}$$

Case II. \mathbb{V} is infinite dimensional.

Exercise.