

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.
3. Basis deletion theorem is valid.

Infinite Dimensional

1. It is not easy to check whether a set is linearly independent or not.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.
3. Basis deletion theorem is valid.
4. Extension theorem is valid.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.
3. Basis deletion theorem is valid.
4. Extension theorem is valid.

Infinite Dimensional

1. It is not easy to check whether a set is linearly independent or not.
2. It is not easy to find a basis of a given infinite dimensional vector space. For example, $C[a, b]$ over the field \mathbb{R} is an infinite dimensional vector space. But we do not have any explicit basis for $C[a, b]$.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.
3. Basis deletion theorem is valid.
4. Extension theorem is valid.

Infinite Dimensional

1. It is not easy to check whether a set is linearly independent or not.
2. It is not easy to find a basis of a given infinite dimensional vector space. For example, $C[a, b]$ over the field \mathbb{R} is an infinite dimensional vector space. But we do not have any explicit basis for $C[a, b]$.
3. Basis deletion theorem is not valid.

Finite Dimensional

1. It is easy to check whether a set is linearly independent or not.
2. It is easy to find a basis of a given finite dimensional vector space.
3. Basis deletion theorem is valid.
4. Extension theorem is valid.

Infinite Dimensional

1. It is not easy to check whether a set is linearly independent or not.
2. It is not easy to find a basis of a given infinite dimensional vector space. For example, $C[a, b]$ over the field \mathbb{R} is an infinite dimensional vector space. But we do not have any explicit basis for $C[a, b]$.
3. Basis deletion theorem is not valid.
4. Extension theorem is not valid.

Finite Dimensional

1. $\dim(\mathbb{V}) = n$ and you have a LI set B of n vectors then B is a basis.

Finite Dimensional

1. $\dim(\mathbb{V}) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(\mathbb{V}) = n$ and you have a set B of n vectors such that $\text{ls}(B) = \mathbb{V}$ then B is a basis.

Finite Dimensional

1. $\dim(\mathbb{V}) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(\mathbb{V}) = n$ and you have a set B of n vectors such that $\text{ls}(B) = \mathbb{V}$ then B is a basis.
3. Gram Schimdt Process is valid only for finite set.

Finite Dimensional

1. $\dim(\mathbb{V}) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(\mathbb{V}) = n$ and you have a set B of n vectors such that $\text{ls}(B) = \mathbb{V}$ then B is a basis.
3. Gram Schimdt Process is valid only for finite set.
4. It is easy to prove any theorem which is valid here.

Infinite Dimensional

1. We do not have this type result for infinite dimensional vector space.

Finite Dimensional

1. $\dim(V) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(V) = n$ and you have a set B of n vectors such that $\text{ls}(B) = V$ then B is a basis.
3. Gram Schimdt Process is valid only for finite set.
4. It is easy to prove any theorem which is valid here.

Infinite Dimensional

1. We do not have this type result for infinite dimensional vector space.
2. We do not have this type result for infinite dimensional vector space.

Finite Dimensional

1. $\dim(\mathbb{V}) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(\mathbb{V}) = n$ and you have a set B of n vectors such that $\text{ls}(B) = \mathbb{V}$ then B is a basis.
3. Gram Schimdt Process is valid only for finite set.
4. It is easy to prove any theorem which is valid here.

Infinite Dimensional

1. We do not have this type result for infinite dimensional vector space.
2. We do not have this type result for infinite dimensional vector space.
3. Gram Schimdt Process is not valid for infinite set.

Finite Dimensional

1. $\dim(V) = n$ and you have a LI set B of n vectors then B is a basis.
2. $\dim(V) = n$ and you have a set B of n vectors such that $\text{ls}(B) = V$ then B is a basis.
3. Gram Schmidt Process is valid only for finite set.
4. It is easy to prove any theorem which is valid here.

Infinite Dimensional

1. We do not have this type result for infinite dimensional vector space.
2. We do not have this type result for infinite dimensional vector space.
3. Gram Schmidt Process is not valid for infinite set.
4. sometime it is difficult to prove any theorem which is valid here.

Questions asked in last class.

Question 1. Let V be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on V such that S is orthogonal.

Questions asked in last class.

Question 1. Let V be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on V such that S is orthogonal.

Answer: There are two cases.

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

Similarly $y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n$.

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

$$\text{Similarly } y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n.$$

$$\text{Then } \langle x, y \rangle = \sum_{i=1}^n c_i \overline{d_i}.$$

Questions asked in last class.

Question 1. Let \mathbb{V} be a vector space over the field \mathbb{K} . Let S be a LI set. Can we have an inner product on \mathbb{V} such that S is orthogonal.

Answer: There are two cases.

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then S is finite set. Let $S = \{u_1, \dots, u_k\}$.

Using extension theorem we can extend S to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

$$\text{Similarly } y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n.$$

$$\text{Then } \langle x, y \rangle = \sum_{i=1}^n c_i \overline{d_i}.$$

You know this one is an inner product.

To check S is orthogonal with respect to above inner product.

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

$$u_j = 0u_1 + 0u_2 + \cdots + u_j + 0u_{j+1} + 0u_n$$

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

$$u_j = 0u_1 + 0u_2 + \cdots + u_j + 0u_{j+1} + 0u_n$$

$\langle u_i, u_j \rangle = 0$, for $i \neq j$. Hence S is orthogonal.

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

$$u_j = 0u_1 + 0u_2 + \cdots + u_j + 0u_{j+1} + 0u_n$$

$\langle u_i, u_j \rangle = 0$, for $i \neq j$. Hence S is orthogonal.

Case-II. \mathbb{V} is infinite dimensional.

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

$$u_j = 0u_1 + 0u_2 + \cdots + u_j + 0u_{j+1} + 0u_n$$

$\langle u_i, u_j \rangle = 0$, for $i \neq j$. Hence S is orthogonal.

Case-II. \mathbb{V} is infinite dimensional.

We cannot apply basis extension theorem. Then how do we proceed?

To check S is orthogonal with respect to above inner product.

Let $u_i, u_j \in S$. Then $u_i = 0u_1 + 0u_2 + \cdots + u_i + 0u_{i+1} + 0u_n$ and

$$u_j = 0u_1 + 0u_2 + \cdots + u_j + 0u_{j+1} + 0u_n$$

$\langle u_i, u_j \rangle = 0$, for $i \neq j$. Hence S is orthogonal.

Case-II. \mathbb{V} is infinite dimensional.

We cannot apply basis extension theorem. Then how do we proceed?

We do not know whether this result is valid for infinite dimensional vector space or not.

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_k \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_n \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

Similarly $y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n$.

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_n \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

$$\text{Similarly } y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n.$$

Then $\langle x, y \rangle$

$$= \langle c_1 u_1 + c_2 u_2 + \dots + c_n u_n, d_1 u_1 + d_2 u_2 + \dots + d_n u_n \rangle$$

$$= \sum_{i=1}^n c_i \overline{d_i}.$$

Question-2 Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Does there exist a basis in \mathbb{V} such that $\langle \cdot, \cdot \rangle$ is equal to the inner product induced by B .

Case-I. \mathbb{V} is finite dimensional and $\dim(\mathbb{V}) = n$. Then there exists a basis $B = \{u_1, \dots, u_n\}$ such that $\langle u_i, u_j \rangle = 0$ for all $i \neq j$ and $\langle u_i, u_i \rangle = 1$ for $i = 1, \dots, n$. (This can be proved using Gram Schmidt process, I will discuss in the next class.)

Let $x, y \in \mathbb{V}$. Then there exist unique $c_1, \dots, c_n \in \mathbb{K}$ such that

$$x = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

$$\text{Similarly } y = d_1 u_1 + d_2 u_2 + \dots + d_n u_n.$$

Then $\langle x, y \rangle$

$$= \langle c_1 u_1 + c_2 u_2 + \dots + c_n u_n, d_1 u_1 + d_2 u_2 + \dots + d_n u_n \rangle$$

$$= \sum_{i=1}^n c_i \overline{d_i}.$$

For finite dimensional inner product space it is true.

Case-I. \mathbb{V} is infinite dimensional.

Case-I. \mathbb{V} is infinite dimensional.

We cannot apply Gram Schimdt process. Then how do we proceed?

Case-I. \mathbb{V} is infinite dimensional.

We cannot apply Gram Schmidt process. Then how do we proceed?

We do not know whether this result is valid for infinite dimensional vector space or not.

- Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be a 2×2 matrix with real entries. Let $f_A : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a map defined by $f_A(x, y) = y^t A x$, where $x, y \in \mathbb{R}^2$. Show that f_A is an inner product on \mathbb{R}^2 if and only if $A = A^t$, $a_{11} > 0$, $a_{22} > 0$ and $\det(A) > 0$.

- Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be a 2×2 matrix with real entries. Let $f_A : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a map defined by $f_A(x, y) = y^t A x$, where $x, y \in \mathbb{R}^2$. Show that f_A is an inner product on \mathbb{R}^2 if and only if $A = A^t$, $a_{11} > 0$, $a_{22} > 0$ and $\det(A) > 0$.

We first assume that f_A is an inner product.

- Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be a 2×2 matrix with real entries. Let $f_A : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a map defined by $f_A(x, y) = y^t A x$, where $x, y \in \mathbb{R}^2$. Show that f_A is an inner product on \mathbb{R}^2 if and only if $A = A^t$, $a_{11} > 0$, $a_{22} > 0$ and $\det(A) > 0$.

We first assume that f_A is an inner product.

Using definition of inner product we have $f_A(x, x) > 0$ for all non-zero $x \in \mathbb{R}^2$.

- Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be a 2×2 matrix with real entries. Let $f_A : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a map defined by $f_A(x, y) = y^t A x$, where $x, y \in \mathbb{R}^2$. Show that f_A is an inner product on \mathbb{R}^2 if and only if $A = A^t$, $a_{11} > 0$, $a_{22} > 0$ and $\det(A) > 0$.

We first assume that f_A is an inner product.

Using definition of inner product we have $f_A(x, x) > 0$ for all non-zero $x \in \mathbb{R}^2$.

$$\text{Then } f_A(e_1, e_1) = e_1^t A e_1 = (1, 0) \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} > 0$$

$$a_{11} > 0.$$

- Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be a 2×2 matrix with real entries. Let $f_A : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a map defined by $f_A(x, y) = y^t A x$, where $x, y \in \mathbb{R}^2$. Show that f_A is an inner product on \mathbb{R}^2 if and only if $A = A^t$, $a_{11} > 0$, $a_{22} > 0$ and $\det(A) > 0$.

We first assume that f_A is an inner product.

Using definition of inner product we have $f_A(x, x) > 0$ for all non-zero $x \in \mathbb{R}^2$.

$$\text{Then } f_A(e_1, e_1) = e_1^t A e_1 = (1, 0) \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} > 0$$

$$a_{11} > 0.$$

Using e_2 we can show that $a_{22} > 0$.

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

$x^t (A^t - A) y = 0$. This is true for all $x, y \in \mathbb{R}^2$.

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

$x^t (A^t - A) y = 0$. This is true for all $x, y \in \mathbb{R}^2$.

Take $x = (1, 0)^t$ and $y = (0, 1)^t$.

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

$x^t (A^t - A) y = 0$. This is true for all $x, y \in \mathbb{R}^2$.

Take $x = (1, 0)^t$ and $y = (0, 1)^t$.

$$(1, 0) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

$x^t (A^t - A) y = 0$. This is true for all $x, y \in \mathbb{R}^2$.

Take $x = (1, 0)^t$ and $y = (0, 1)^t$.

$$(1, 0) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_{21} - a_{12} = 0 \implies a_{12} = a_{21}.$$

Using definition of inner product we have $f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$.

$$f_A(x, y) = \overline{f_A(y, x)} = f_A(y, x)$$

$$y^t A x = x^t A y$$

$$(y^t A x)^t = x^t A y$$

$$x^t A^t y = x^t A y$$

$x^t (A^t - A) y = 0$. This is true for all $x, y \in \mathbb{R}^2$.

Take $x = (1, 0)^t$ and $y = (0, 1)^t$.

$$(1, 0) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_{21} - a_{12} = 0 \implies a_{12} = a_{21}.$$

Hence $A = A^t$.

To prove $\det(A) > 0$, we take $x = (a_{22}, -a_{12})$.

To prove $\det(A) > 0$, we take $x = (a_{22}, -a_{12})$.

Since $f_A(x, x) > 0$, we have $(a_{11}, -a_{12}) \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{pmatrix} \begin{pmatrix} a_{11} \\ -a_{12} \end{pmatrix} > 0$

To prove $\det(A) > 0$, we take $x = (a_{22}, -a_{12})$.

Since $f_A(x, x) > 0$, we have $(a_{11}, -a_{12}) \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{pmatrix} \begin{pmatrix} a_{11} \\ -a_{12} \end{pmatrix} > 0$

Then $a_{11}a_{22} - a_{12}^2 = \det(A) > 0$.

We now prove the converse.

We now prove the converse.

$$1(a). \quad f_A(x, x) = (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

We now prove the converse.

$$\begin{aligned} 1(a). \quad f_A(x, x) &= (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21}) \end{aligned}$$

We now prove the converse.

$$\begin{aligned} 1(a). \quad f_A(x, x) &= (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21}) \\ &= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0) \end{aligned}$$

We now prove the converse.

$$\begin{aligned} 1(a). \quad f_A(x, x) &= (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21}) \\ &= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0) \\ &> 0 \end{aligned}$$

We now prove the converse.

$$1(a). \quad f_A(x, x) = (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21})$$

$$= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0)$$

$$> 0$$

$$1(b). \quad f_A(x, x) = 0. \quad \text{Using } a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2, \text{ this we have } x_1 = 0 \text{ and } x_2 = 0.$$

We now prove the converse.

$$1(a). \quad f_A(x, x) = (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21})$$

$$= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0)$$

$$> 0$$

$$1(b). \quad f_A(x, x) = 0. \quad \text{Using } a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2, \text{ this we have } x_1 = 0 \text{ and } x_2 = 0.$$

2. It is trivial.

We now prove the converse.

$$1(a). \quad f_A(x, x) = (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21})$$

$$= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0)$$

$$> 0$$

1(b). $f_A(x, x) = 0$. Using $a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2$, this we have $x_1 = 0$ and $x_2 = 0$.

2. It is trivial.

3. It is trivial.

We now prove the converse.

$$1(a). \quad f_A(x, x) = (x_1, x_2) \begin{pmatrix} 0 & a_{21} - a_{12} \\ a_{12} - a_{21} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$= a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2 \quad (\text{as } a_{12} = a_{21})$$

$$= a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2 \quad (\text{as } a_{11} > 0)$$

$$> 0$$

1(b). $f_A(x, x) = 0$. Using $a_{11}\left(x_1 + \frac{a_{12}}{a_{11}x_2}\right)^2 + \frac{a_{11}a_{22} - a_{12}^2}{a_{11}}x_2^2$, this we have $x_1 = 0$ and $x_2 = 0$.

2. It is trivial.

3. It is trivial.

4. It is trivial.

-
- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .

- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .
- $f_A(x, y) = x^t A y$ is an inner product on \mathbb{R}^3 if and only if A is symmetric, $a_{ii} > 0$ for $i = 1, \dots, 3$ and all the leading principal minors are non-zero.

- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .
- $f_A(x, y) = x^t A y$ is an inner product on \mathbb{R}^3 if and only if A is symmetric, $a_{ii} > 0$ for $i = 1, \dots, 3$ and all the leading principal minors are non-zero.

Leading Principal minor: There are three leading principal of a matrix of order 3.

- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .
- $f_A(x, y) = x^t A y$ is an inner product on \mathbb{R}^3 if and only if A is symmetric, $a_{ii} > 0$ for $i = 1, \dots, 3$ and all the leading principal minors are non-zero.

Leading Principal minor: There are three leading principal of a matrix of order 3.

Let $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .
- $f_A(x, y) = x^t A y$ is an inner product on \mathbb{R}^3 if and only if A is symmetric, $a_{ii} > 0$ for $i = 1, \dots, 3$ and all the leading principal minors are non-zero.

Leading Principal minor: There are three leading principal of a matrix of order 3.

$$\text{Let } A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}.$$

1. a_{11} is the leading principal minor of size 1.

- If A is a 3×3 matrix. Necessary and sufficient condition for $f_A(x, y) = x^t A y$ to be an inner product on \mathbb{R}^3 .
- $f_A(x, y) = x^t A y$ is an inner product on \mathbb{R}^3 if and only if A is symmetric, $a_{ii} > 0$ for $i = 1, \dots, 3$ and all the leading principal minors are non-zero.

Leading Principal minor: There are three leading principal of a matrix of order 3.

Let $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

1. a_{11} is the leading principal minor of size 1.

$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$ is the leading principal minor of size 2.

$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$ is the leading principal minor of size 3.