

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

If $x = (0, 0, \dots, 0)$, then $x \cdot x = 0$.

- Consider dot product on \mathbb{R}^n . That is

$$x \cdot y = \sum_{i=1}^n x_i y_i.$$

Observe that

$$\text{i) } x \cdot x = \sum_{i=1}^n x_i x_i = \sum_{i=1}^n x_i^2 \geq 0$$

$$x \cdot x = \sum_{i=1}^n x_i^2 = 0 \implies x_i = 0 \text{ for } i = 1, \dots, n, \text{ then } x = (0, 0, \dots, 0).$$

If $x = (0, 0, \dots, 0)$, then $x \cdot x = 0$.

$$\text{ii) } x \cdot y = \sum_{i=1}^n x_i y_i = \sum_{i=1}^n y_i x_i = y \cdot x.$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.
- How do we understand the geometry of an arbitrary vector space?

$$\text{iii) } \alpha x \cdot y = \sum_{i=1}^n (\alpha x_i) y_i = \alpha \sum_{i=1}^n x_i y_i = \alpha (x \cdot y).$$

$$\text{iv) } (x + y) \cdot z = x \cdot z + y \cdot z.$$

- Dot product on \mathbb{R}^n helps us understand the geometry of \mathbb{R}^n with tools to detect angles and distances.
- How do we understand the geometry of an arbitrary vector space?
- Inner product can be used to understand the geometry of abstract vector spaces.

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

An inner product is a mapping $\langle ., . \rangle : \mathbb{V} \times \mathbb{V} \rightarrow \mathbb{K}$ which satisfies the following conditions.

1. $\langle x, x \rangle \geq 0$ for all $x \in V$ (**positivity**)

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

4. $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$ for all $x, y, z \in \mathbb{V}$ (**additivity**).

Examples:

- Dot product on \mathbb{R}^n is an inner product.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_i^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Ans: This is an inner product.

- $V = C[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in C[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

- $\mathbb{V} = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

- $\mathbb{V} = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

$$\text{Then } \langle \alpha f, g \rangle = \int_a^b (\alpha f(x))g(x)dx = \alpha \int_a^b f(x)g(x)dx = \alpha \langle f, g \rangle.$$

4. Let $f, g, h \in \mathbb{C}[a, b]$.

4. Let $f, g, h \in \mathbb{C}[a, b]$.

$$\begin{aligned}\text{Then } \langle f + g, h \rangle &= \int_a^b (f(x) + g(x)) h(x) dx \\ &= \int_a^b (f(x)h(x) + g(x)h(x)) dx \\ &= \langle f, h \rangle + \langle g, h \rangle.\end{aligned}$$

- Let $f : [a, b]$ be integrable and $f(x) \geq 0$. Let f be continuous at $x = c$ and $f(c) > 0$ (resp. $f(c) < 0$). Then $\int_a^b f(x)dx > 0$ (resp. $\int_a^b f(x)dx < 0$).

Sol: Case I. Let $a < c < b$.

f is continuous at $x = c$. Take $\epsilon = \frac{f(c)}{2}$ then there exists $\delta > 0$ such that

for each $x \in (c - \delta, c + \delta) \implies |f(x) - f(c)| < \frac{f(c)}{2}$.

Then $-\frac{f(c)}{2} < f(x) - f(c) < \frac{f(c)}{2}$ for all $x \in (c - \delta, c + \delta)$.

Therefore $f(x) > \frac{f(c)}{2} > 0$ for all $x \in (c - \delta, c + \delta)$.

$$\int_a^b f^2(x)dx = \int_a^{c-\delta} f(x)dx + \int_{c-\delta}^{c+\delta} f(x)dx + \int_{c+\delta}^b f(x)dx > 0$$

as $\int_{c-\delta}^{c+\delta} f^2(x)dx > 0$.

Case II. $c = a$

When $c = a$, we have a neighborhood $(a, a + \delta)$ where f is positive.

$$\int_a^b f^2(x) dx = \int_a^{a+\delta} f(x) dx + \int_b^{a+\delta} f(x) dx > 0$$

Case-III. $c = b$. Same as Case II.

To show $\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0$.

Here f^2 is nonnegative function and continuous. If f is not identically zero, then there exists $c \in [a, b]$ such that $f(c) \neq 0$. Therefore $f^2(c) > 0$. Using above result we have $\int_a^b f(x)f(x)dx > 0$, a contradiction. Hence $f \equiv 0$.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

Hence $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ is not an inner product on $C(\mathbb{R})$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, k\}$ be a basis of \mathbb{V} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

We now show that \langle, \rangle is an IP on \mathbb{V} .

-
1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

Therefore $x = \sum_{i=1}^k a_i u_i = 0$.

2. Let $x, y \in \mathbb{V}$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, y \rangle = \sum_{i=1}^k a_i \overline{b_i} = \overline{\sum_{i=1}^k \overline{a_i} b_i} = \overline{\langle y, x \rangle}.$$

3. This part is same as part 2.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Then $x + y = \sum_{i=1}^k (a_i + b_i) u_i$. Therefore,

$$\begin{aligned}\langle x + y, z \rangle &= \sum_{i=1}^k (a_i + b_i) \overline{c_i} \\ &= \sum_{i=1}^k (a_i \overline{c_i} + b_i \overline{c_i}) \\ &= \langle x, z \rangle + \langle y, z \rangle.\end{aligned}$$

Case II. \mathbb{V} is infinite dimensional.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Define $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_i} f(u_{\alpha_i}, u_{\alpha_i}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_i} f(u_{\alpha_i}, u_{\alpha_i}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k} \text{ where } a_i \in \mathbb{K} \text{ for } i = 1, \dots, k.$$

$$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m} \text{ where } b_j \in \mathbb{K} \text{ for } j = 1, \dots, m.$$

$$z = c_1 u_{\gamma_1} + \dots + c_n u_{\gamma_n} \text{ where } c_l \in \mathbb{K} \text{ for } l = 1, \dots, n.$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle.$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle.$$

- A vectors space together with an inner product is called an inner product space.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
 3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
 3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.
 4. $\langle u, \lambda v \rangle = \overline{\lambda} \langle u, v \rangle$ for all $u, v \in \mathbb{V}$ and $\lambda \in \mathbb{K}$.

1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.
4. $\langle u, \lambda v \rangle = \overline{\lambda} \langle u, v \rangle$ for all $u, v \in \mathbb{V}$ and $\lambda \in \mathbb{K}$.

Proof: 1. $\langle 0, u \rangle = \langle u - u, u \rangle$

$$= \langle u, u \rangle + \langle -u, u \rangle$$

$$= \langle u, u \rangle - \langle u, u \rangle$$

$$= 0.$$

$$2. \langle 0, u \rangle = \overline{\langle 0, u \rangle}$$

$$= \overline{\langle u - u, u \rangle}$$

$$= \overline{\langle u, u \rangle + \langle -u, u \rangle}$$

$$= \overline{\langle u, u \rangle - \langle u, u \rangle}$$

$$= \overline{0}$$

3. Suppose $u, v, w \in \mathbb{V}$. Then

$$\begin{aligned}\langle u, v + w \rangle &= \overline{\langle v + w, u \rangle} \\ &= \overline{\langle v, u \rangle + \langle w, u \rangle} \\ &= \overline{\langle v, u \rangle} + \overline{\langle w, u \rangle} \\ &= \langle u, v \rangle + \langle u, w \rangle.\end{aligned}$$

4. Suppose $\lambda \in \mathbb{K}$ and $u, v \in \mathbb{V}$. Then

$$\langle u, \lambda v \rangle = \overline{\langle \lambda v, u \rangle}$$

$$= \overline{\lambda \langle v, u \rangle}$$

$$= \overline{\lambda} \overline{\langle v, u \rangle}$$

$$= \overline{\lambda} \langle u, v \rangle.$$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

$\langle e_1, e_2 \rangle_1 = -1$. These two vectors are not orthogonal with respect to $\langle \cdot, \cdot \rangle_1$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

$\langle e_1, e_2 \rangle_1 = -1$. These two vectors are not orthogonal with respect to $\langle \cdot, \cdot \rangle_1$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. A subset S of \mathbb{V} is said to be **orthogonal** if $\langle u, v \rangle = 0$ for all $u, v \in S$ and $u \neq v$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$. Take inner product with α_i both side.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$. Take inner product with α_i both side.

$$\langle c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k, \alpha_i \rangle = \langle 0, \alpha_i \rangle$$

$$\sum_{j=1}^k c_j \langle \alpha_j, \alpha_i \rangle = 0.$$

$$c_i \langle \alpha_i, \alpha_i \rangle = 0$$

$c_i = 0$. This is true for $i = 1, \dots, k$.

Hence $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

- The converse of the above theorem is not true. The following is an example.

- The converse of the above theorem is not true. The following is an example.

Example: Let $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$ be an inner product space where $\langle x, y \rangle = \sum_{i=1}^2 x_i y_i$.

- The converse of the above theorem is not true. The following is an example.

Example: Let $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$ be an inner product space where $\langle x, y \rangle = \sum_{i=1}^2 x_i y_i$.

Take $u = (1, 0)$ and $v = (1, 1)$. These two vectors are linearly independent but not orthogonal.

- The converse of the above theorem is not true. The following is an example.

Example: Let $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$ be an inner product space where $\langle x, y \rangle = \sum_{i=1}^2 x_i y_i$.

Take $u = (1, 0)$ and $v = (1, 1)$. These two vectors are linearly independent but not orthogonal.

- [Definition:] Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space.
- [Definition:] Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq \mathbb{V}$. Then $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is called **orthonormal** if $\langle \alpha_i, \alpha_j \rangle = 0$ for $i \neq j$ and $\langle \alpha_i, \alpha_i \rangle = 1$.