

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$.

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$. Then Ax is also a vector in \mathbb{F}^n .

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$. Then Ax is also a vector in \mathbb{F}^n .

- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$. Then Ax is also a vector in \mathbb{F}^n .


- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. $Ax = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$. Then Ax is also a vector in \mathbb{F}^n .

- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. $Ax = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$.
 $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $x \in \mathbb{F}^n$. Then Ax is also a vector in \mathbb{F}^n .

- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. $Ax = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$.
 $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$.



-
- An eigenvector of a matrix A is a nonzero vector that changes by a scalar factor when the matrix is multiplied with that vector.

-
- An eigenvector of a matrix A is a nonzero vector that changes by a scalar factor when the matrix is multiplied with that vector.
 - **[Definition:]** Let $A \in \mathbb{M}_n(\mathbb{F})$. A scalar λ is said to be an **eigenvalue** of A if there exists a non-zero vector $x \in \mathbb{F}^n$ such that $Ax = \lambda x$. Any such (non-zero) x is called an **eigenvector** of A corresponding to the eigenvalue λ .

-
- An eigenvector of a matrix A is a nonzero vector that changes by a scalar factor when the matrix is multiplied with that vector.
 - **[Definition:]** Let $A \in \mathbb{M}_n(\mathbb{F})$. A scalar λ is said to be an **eigenvalue** of A if there exists a non-zero vector $x \in \mathbb{F}^n$ such that $Ax = \lambda x$. Any such (non-zero) x is called an **eigenvector** of A corresponding to the eigenvalue λ .
 - $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

-
- An eigenvector of a matrix A is a nonzero vector that changes by a scalar factor when the matrix is multiplied with that vector.

- **[Definition:]** Let $A \in \mathbb{M}_n(\mathbb{F})$. A scalar λ is said to be an **eigenvalue** of A if there exists a non-zero vector $x \in \mathbb{F}^n$ such that $Ax = \lambda x$. Any such (non-zero) x is called an **eigenvector** of A corresponding to the eigenvalue λ .

- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. $Ax = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Here 3 is an eigenvalue of A and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is an eigenvector of A corresponding to 3.

-
- An eigenvector of a matrix A is a nonzero vector that changes by a scalar factor when the matrix is multiplied with that vector.

- **[Definition:]** Let $A \in \mathbb{M}_n(\mathbb{F})$. A scalar λ is said to be an **eigenvalue** of A if there exists a non-zero vector $x \in \mathbb{F}^n$ such that $Ax = \lambda x$. Any such (non-zero) x is called an **eigenvector** of A corresponding to the eigenvalue λ .

- $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. $Ax = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Here 3 is an eigenvalue of A and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is an eigenvector of A corresponding to 3.

- **[Geometrically]** An eigenvector, corresponding to a real nonzero eigenvalue, points in a direction in which it is stretched by the matrix and the eigenvalue is the factor by which it is stretched. If the eigenvalue is negative, the direction is reversed.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose coefficients are coming from \mathbb{F} .

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose co-efficients are coming from \mathbb{F} .

- For example consider the following matrix $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Then $\det(xI - A) = x^3 - 6x^2 + 11x - 6$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose co-efficients are coming from \mathbb{F} .
 - For example consider the following matrix $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Then $\det(xI - A) = x^3 - 6x^2 + 11x - 6$.
 - Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and $\lambda \in \mathbb{F}$. Then λ is an eigenvalue of A if and only if λ is a root of the polynomial $\det(xI - A)$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose co-efficients are coming from \mathbb{F} .

- For example consider the following matrix $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Then $\det(xI - A) = x^3 - 6x^2 + 11x - 6$.

- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and $\lambda \in \mathbb{F}$. Then λ is an eigenvalue of A if and only if λ is a root of the polynomial $\det(xI - A)$.

Proof. We first assume that λ is an eigenvalue of A .

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose co-efficients are coming from \mathbb{F} .

- For example consider the following matrix $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Then $\det(xI - A) = x^3 - 6x^2 + 11x - 6$.

- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and $\lambda \in \mathbb{F}$. Then λ is an eigenvalue of A if and only if λ is a root of the polynomial $\det(xI - A)$.

Proof. We first assume that λ is an eigenvalue of A . To show λ is a root of the polynomial $\det(xI - A)$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$. Then $\det(xI - A)$ is a polynomial of x whose co-efficients are coming from \mathbb{F} .

- For example consider the following matrix $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Then $\det(xI - A) = x^3 - 6x^2 + 11x - 6$.

- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and $\lambda \in \mathbb{F}$. Then λ is an eigenvalue of A if and only if λ is a root of the polynomial $\det(xI - A)$.

Proof. We first assume that λ is an eigenvalue of A . To show λ is a root of the polynomial $\det(xI - A)$.

There exists a non-zero vector $x \in \mathbb{F}^n$ such that $Ax = \lambda x$.

$$Ax - \lambda x = 0$$

$$(A - \lambda I)x = 0$$

$$Ax - \lambda x = 0$$

$$(A - \lambda I)x = 0$$

This says that the system of homogeneous equations $(A - \lambda I)y = 0$ has non-trivial solution. Hence $\text{rank}(A - \lambda I) < n$. Then $\det(A - \lambda I) = 0 = \det(\lambda I - A)$. This implies λ is a root of $\det(xI - A)$.

$$Ax - \lambda x = 0$$

$$(A - \lambda I)x = 0$$

This says that the system of homogeneous equations $(A - \lambda I)y = 0$ has non-trivial solution. Hence $\text{rank}(A - \lambda I) < n$. Then $\det(A - \lambda I) = 0 = \det(\lambda I - A)$. This implies λ is a root of $\det(xI - A)$.

We now assume that λ is a root of the polynomial $\det(xI - A)$. To show λ is an eigenvalue of A .

$$Ax - \lambda x = 0$$

$$(A - \lambda I)x = 0$$

This says that the system of homogeneous equations $(A - \lambda I)y = 0$ has non-trivial solution. Hence $\text{rank}(A - \lambda I) < n$. Then $\det(A - \lambda I) = 0 = \det(\lambda I - A)$. This implies λ is a root of $\det(xI - A)$.

We now assume that λ is a root of the polynomial $\det(xI - A)$. To show λ is an eigenvalue of A .

$$\det(\lambda I - A) = 0$$

$$\text{rank}(A - \lambda I) < n$$

$$Ax - \lambda x = 0$$

$$(A - \lambda I)x = 0$$

This says that the system of homogeneous equations $(A - \lambda I)y = 0$ has non-trivial solution. Hence $\text{rank}(A - \lambda I) < n$. Then $\det(A - \lambda I) = 0 = \det(\lambda I - A)$. This implies λ is a root of $\det(xI - A)$.

We now assume that λ is a root of the polynomial $\det(xI - A)$. To show λ is an eigenvalue of A .

$$\det(\lambda I - A) = 0$$

$$\text{rank}(A - \lambda I) < n$$

$(A - \lambda I)x = 0$ has non-trivial solution. There is a non-zero $y \in \mathbb{F}^n$ such that $Ay = \lambda y$. Hence λ is an eigenvalue.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$. Then the polynomial $\det(xI - A)$ is called characteristic polynomial and the equation $\det(xI - A) = 0$ is called characteristic equation.

- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. Calculate the eigenvalues of A .

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$. Then the polynomial $\det(xI - A)$ is called characteristic polynomial and the equation $\det(xI - A) = 0$ is called characteristic equation.

- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. Calculate the eigenvalues of A .

We have to calculate the roots of $\det(xI - A)$.

$$\det(xI - A) = \begin{vmatrix} x-1 & -2 & -1 \\ -2 & x-1 & -1 \\ -1 & -1 & x-2 \end{vmatrix} = x^3 - 4x^2 - x + 4.$$

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$. Then the polynomial $\det(xI - A)$ is called characteristic polynomial and the equation $\det(xI - A) = 0$ is called characteristic equation.

- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. Calculate the eigenvalues of A .

We have to calculate the roots of $\det(xI - A)$.

$$\det(xI - A) = \begin{vmatrix} x-1 & -2 & -1 \\ -2 & x-1 & -1 \\ -1 & -1 & x-2 \end{vmatrix} = x^3 - 4x^2 - x + 4.$$

The roots of $x^3 - 4x^2 - x + 4$ are $1, -1, 4$.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

Eigenvectors corresponding to $\lambda = 1$.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

Eigenvectors corresponding to $\lambda = 1$.

The eigenvectors of A corresponding to $\lambda = 1$ is the set of all non-zero solutions of $(A - I)x = 0$.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

Eigenvectors corresponding to $\lambda = 1$.

The eigenvectors of A corresponding to $\lambda = 1$ is the set of all non-zero solutions of $(A - I)x = 0$.

$$A - I = \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

Eigenvectors corresponding to $\lambda = 1$.

The eigenvectors of A corresponding to $\lambda = 1$ is the set of all non-zero solutions of $(A - I)x = 0$.

$$A - I = \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

Then REF of $A - I$ is $\begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. The set $\left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} - \{0\} \right\}$ is the set of all eigenvectors.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$.

Eigenvectors corresponding to $\lambda = 1$.

The eigenvectors of A corresponding to $\lambda = 1$ is the set of all non-zero solutions of $(A - I)x = 0$.

$$A - I = \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

Then REF of $A - I$ is $\begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. The set $\left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} - \{0\} \right\}$ is the set of all eigenvectors.

-
- The following question is quite natural. Does every matrix have eigenvalue?

-
- The following question is quite natural. Does every matrix have eigenvalue?
 - This question is similar to the following question. Let $P(x) \in \mathbb{P}(x, \mathbb{F})$, where $\mathbb{P}(x, \mathbb{F})$ set of all polynomials with coefficients are coming from \mathbb{F} . Does $P(x)$ have root in \mathbb{F} ?

-
- The following question is quite natural. Does every matrix have eigenvalue?
 - This question is similar to the following question. Let $P(x) \in \mathbb{P}(x, \mathbb{F})$, where $\mathbb{P}(x, \mathbb{F})$ set of all polynomials with coefficients are coming from \mathbb{F} . Does $P(x)$ have root in \mathbb{F} ?

The answer is no. For example, consider the polynomial $x^2 + 1$ in $\mathbb{P}(x, \mathbb{R})$, this polynomial does not have root in \mathbb{R} .

-
- The following question is quite natural. Does every matrix have eigenvalue?
 - This question is similar to the following question. Let $P(x) \in \mathbb{P}(x, \mathbb{F})$, where $\mathbb{P}(x, \mathbb{F})$ set of all polynomials with coefficients are coming from \mathbb{F} . Does $P(x)$ have root in \mathbb{F} ?

The answer is no. For example, consider the polynomial $x^2 + 1$ in $\mathbb{P}(x, \mathbb{R})$, this polynomial does not have root in \mathbb{R} .

Now we are able to answer our question. If $x^2 + 1$ is the characteristic polynomial of a matrix $A \in \mathbb{M}_2(\mathbb{R})$, then A does not have eigenvalues. Here is that $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

-
- **[Theorem]** Let $A \in \mathbb{M}_{n \times n}(\mathbb{C})$. Then A has at least one eigenvalue.

-
- **[Theorem]** Let $A \in \mathbb{M}_{n \times n}(\mathbb{C})$. Then A has at least one eigenvalue.

Proof: The characteristic polynomial of A is $P_A(z) = \det(A - zI)$. The Fundamental Theorem of Algebra says that $P_A(x)$ has at least one root. Hence A has at least one eigenvalue.

-
- **[Theorem]** Let $A \in \mathbb{M}_{n \times n}(\mathbb{C})$. Then A has at least one eigenvalue.

Proof: The characteristic polynomial of A is $P_A(z) = \det(A - zI)$. The Fundamental Theorem of Algebra says that $P_A(x)$ has at least one root. Hence A has at least one eigenvalue.

- **[Theorem]** Similar matrices have the same characteristic polynomial. But the converse is not true.

-
- **[Theorem]** Let $A \in \mathbb{M}_{n \times n}(\mathbb{C})$. Then A has at least one eigenvalue.

Proof: The characteristic polynomial of A is $P_A(z) = \det(A - zI)$. The Fundamental Theorem of Algebra says that $P_A(x)$ has at least one root. Hence A has at least one eigenvalue.

- **[Theorem]** Similar matrices have the same characteristic polynomial. But the converse is not true.

Proof: Let A and B be two similar matrices.

-
- **[Theorem]** Let $A \in \mathbb{M}_{n \times n}(\mathbb{C})$. Then A has at least one eigenvalue.

Proof: The characteristic polynomial of A is $P_A(z) = \det(A - zI)$. The Fundamental Theorem of Algebra says that $P_A(x)$ has at least one root. Hence A has at least one eigenvalue.

- **[Theorem]** Similar matrices have the same characteristic polynomial. But the converse is not true.

Proof: Let A and B be two similar matrices.

Then there exists a nonsingular matrix P such that $P^{-1}AP = B$.

Then $\det(B - xI)$

$$= \det(P^{-1}AP - xI)$$

$$= \det(P^{-1}AP - xP^{-1}P)$$

$$= \det(P^{-1}) \det(A - xI) \det(P) = \det(A - xI).$$

Converse is not true. Consider the following two matrices.

$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$. These two matrices have the same characteristic polynomial but they are not similar.

Converse is not true. Consider the following two matrices.

$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$. These two matrices have the same characteristic polynomial but they are not similar.

- Let $C = \begin{bmatrix} A_{n \times n} & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} \end{bmatrix}$. Then characteristic polynomial of C , $P_C(x) = P_A(x)P_B(x)$.

Converse is not true. Consider the following two matrices.

$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$. These two matrices have the same characteristic polynomial but they are not similar.

- Let $C = \begin{bmatrix} A_{n \times n} & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} \end{bmatrix}$. Then characteristic polynomial of C , $P_C(x) = P_A(x)P_B(x)$.

We have $C - xI_{n+m} = \begin{bmatrix} A_{n \times n} - xI_n & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} - xI_m \end{bmatrix}$.

Converse is not true. Consider the following two matrices.

$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$. These two matrices have the same characteristic polynomial but they are not similar.

- Let $C = \begin{bmatrix} A_{n \times n} & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} \end{bmatrix}$. Then characteristic polynomial of C , $P_C(x) = P_A(x)P_B(x)$.

We have $C - xI_{n+m} = \begin{bmatrix} A_{n \times n} - xI_n & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} - xI_m \end{bmatrix}$.

Then $\det(C) = \begin{vmatrix} A_{n \times n} - xI_n & D_{n \times m} \\ 0_{m \times n} & B_{m \times m} - xI_m \end{vmatrix} = \det(A - xI_n) \det(B - xI_m)$.

-
- [Theorem] Let $f(x)$ be a polynomial and let (λ, x) be an eigenpair of A . Then $(f(\lambda), x)$ is an eigenpair of $f(A)$. But the converse is not true.

-
- [Theorem] Let $f(x)$ be a polynomial and let (λ, x) be an eigenpair of A . Then $(f(\lambda), x)$ is an eigenpair of $f(A)$. But the converse is not true.

Proof: If (λ, x) is an eigenpair of A , then (λ^k, x) is an eigenpair of A^k where k is natural number. Using this we can easily prove that $(f(\lambda), x)$ is an eigenpair of $f(A)$.

-
- [Theorem] Let $f(x)$ be a polynomial and let (λ, x) be an eigenpair of A . Then $(f(\lambda), x)$ is an eigenpair of $f(A)$. But the converse is not true.

Proof: If (λ, x) is an eigenpair of A , then (λ^k, x) is an eigenpair of A^k where k is natural number. Using this we can easily prove that $(f(\lambda), x)$ is an eigenpair of $f(A)$.

Converse: Let λ be an eigenvalue of $f(A)$. Does there have an eigenvalue μ of A such that $f(\mu) = \lambda$

-
- [Theorem] Let $f(x)$ be a polynomial and let (λ, x) be an eigenpair of A . Then $(f(\lambda), x)$ is an eigenpair of $f(A)$. But the converse is not true.

Proof: If (λ, x) is an eigenpair of A , then (λ^k, x) is an eigenpair of A^k where k is natural number. Using this we can easily prove that $(f(\lambda), x)$ is an eigenpair of $f(A)$.

Converse: Let λ be an eigenvalue of $f(A)$. Does there have an eigenvalue μ of A such that $f(\mu) = \lambda$

Let $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \in \mathbb{M}_n(\mathbb{R})$ and let $f(x) = x^2$.

The characteristic polynomial of A is $x^2 + 1$ and this polynomial does not have any real root. Hence A does not have any eigenvalues. Then

$$f(A) = A^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}.$$

The eigenvalues of $f(A)$ are $-1, -1$. Then there is no eigenvalue μ in A such that $f(\mu) = -1$.

-
- [Theorem] Let $f(x)$ be a polynomial and let (λ, x) be an eigenpair of A . Then $(f(\lambda), x)$ is an eigenpair of $f(A)$. But the converse is not true.

Proof: If (λ, x) is an eigenpair of A , then (λ^k, x) is an eigenpair of A^k where k is natural number. Using this we can easily prove that $(f(\lambda), x)$ is an eigenpair of $f(A)$.

Converse: Let λ be an eigenvalue of $f(A)$. Does there have an eigenvalue μ of A such that $f(\mu) = \lambda$

Let $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \in \mathbb{M}_n(\mathbb{R})$ and let $f(x) = x^2$.

The characteristic polynomial of A is $x^2 + 1$ and this polynomial does not have any real root. Hence A does not have any eigenvalues. Then

$$f(A) = A^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}.$$

The eigenvalues of $f(A)$ are $-1, -1$. Then there is no eigenvalue μ in A such that $f(\mu) = -1$.

That means if (λ, x) is an eigenpair of $f(A)$, then there may not have

$$\bullet \begin{bmatrix} A_{n \times n} & B_{n \times m} \\ C_{m \times n} & D_{m \times m} \end{bmatrix} \begin{bmatrix} E_{n \times n} & F_{n \times m} \\ G_{m \times n} & H_{m \times m} \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}$$

- **[Result]** Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$ and $B \in \mathbb{M}_{n \times m}$, where $m \geq n$. Then $P_{AB}(x) = x^{m-n}P_{BA}(x)$.

- $$\begin{bmatrix} A_{n \times n} & B_{n \times m} \\ C_{m \times n} & D_{m \times m} \end{bmatrix} \begin{bmatrix} E_{n \times n} & F_{n \times m} \\ G_{m \times n} & H_{m \times m} \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}$$

- **[Result]** Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$ and $B \in \mathbb{M}_{n \times m}$, where $m \geq n$. Then $P_{AB}(x) = x^{m-n} P_{BA}(x)$.

We can write
$$\begin{bmatrix} I_m & -A \\ 0 & I_n \end{bmatrix} \begin{bmatrix} AB & 0 \\ B & 0_n \end{bmatrix} \begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix} = \begin{bmatrix} 0_m & 0 \\ B & BA \end{bmatrix}$$

- $$\begin{bmatrix} A_{n \times n} & B_{n \times m} \\ C_{m \times n} & D_{m \times m} \end{bmatrix} \begin{bmatrix} E_{n \times n} & F_{n \times m} \\ G_{m \times n} & H_{m \times m} \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}$$

- **[Result]** Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$ and $B \in \mathbb{M}_{n \times m}$, where $m \geq n$. Then $P_{AB}(x) = x^{m-n} P_{BA}(x)$.

We can write
$$\begin{bmatrix} I_m & -A \\ 0 & I_n \end{bmatrix} \begin{bmatrix} AB & 0 \\ B & 0_n \end{bmatrix} \begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix} = \begin{bmatrix} 0_m & 0 \\ B & BA \end{bmatrix}$$

The matrix $\begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix}$ is invertible and $\begin{bmatrix} I_m & -A \\ 0 & I_n \end{bmatrix}$ is the inverse.

$$\bullet \begin{bmatrix} A_{n \times n} & B_{n \times m} \\ C_{m \times n} & D_{m \times m} \end{bmatrix} \begin{bmatrix} E_{n \times n} & F_{n \times m} \\ G_{m \times n} & H_{m \times m} \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}$$

• **[Result]** Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$ and $B \in \mathbb{M}_{n \times m}$, where $m \geq n$. Then $P_{AB}(x) = x^{m-n} P_{BA}(x)$.

We can write $\begin{bmatrix} I_m & -A \\ 0 & I_n \end{bmatrix} \begin{bmatrix} AB & 0 \\ B & 0_n \end{bmatrix} \begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix} = \begin{bmatrix} 0_m & 0 \\ B & BA \end{bmatrix}$

The matrix $\begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix}$ is invertible and $\begin{bmatrix} I_m & -A \\ 0 & I_n \end{bmatrix}$ is the inverse.

$$P^{-1} \begin{bmatrix} AB & 0 \\ B & 0_n \end{bmatrix} P = \begin{bmatrix} 0_m & 0 \\ B & BA \end{bmatrix} \text{ where } P = \begin{bmatrix} I_m & A \\ 0 & I_n \end{bmatrix}.$$

-
- [Definition:] Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . The number of times λ appears as a root of the characteristic polynomial of A is called **algebraic multiplicity**.

-
- [Definition:] Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . The number of times λ appears as a root of the characteristic polynomial of A is called **algebraic multiplicity**.

Let $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$. The eigenvalues of A are 1, 1, 2. The algebraic multiplicity of 1 is 2 and 2 is 1.

-
- [Definition:] Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . The number of times λ appears as a root of the characteristic polynomial of A is called **algebraic multiplicity**.

Let $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$. The eigenvalues of A are 1, 1, 2. The algebraic multiplicity of 1 is 2 and 2 is 1.

Consider the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 4 & 2 & 0 \\ 6 & 0 & 2 \end{bmatrix}$. The eigenvalues of A are 2, 2, 2. The algebraic multiplicity of 2 is 3.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

Then we show that $k = p$.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

Then we show that $k = p$. Suppose that $k < p$. Then $S_{k+1} = \{x_1, \dots, x_k, x_{k+1}\}$ is linearly dependent.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

Then we show that $k = p$. Suppose that $k < p$. Then $S_{k+1} = \{x_1, \dots, x_k, x_{k+1}\}$ is linearly dependent.

In that case if $\sum_{i=1}^{k+1} \alpha_i x_i = 0$ $\alpha_1, \dots, \alpha_{k+1} \in \mathbb{F}$ not all zero.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

Then we show that $k = p$. Suppose that $k < p$. Then $S_{k+1} = \{x_1, \dots, x_k, x_{k+1}\}$ is linearly dependent.

In that case if $\sum_{i=1}^{k+1} \alpha_i x_i = 0$ $\alpha_1, \dots, \alpha_{k+1} \in \mathbb{F}$ not all zero. Then $\alpha_{k+1} \neq 0$, otherwise S_{k+1} will be linearly independent.

-
- **[Theorem:]** Let $\lambda_1, \dots, \lambda_p$ be distinct eigenvalues of A . Let x_1, \dots, x_p be eigenvectors of A corresponding to $\lambda_1, \dots, \lambda_p$. Then x_1, x_2, \dots, x_p are LI.

Proof: Consider $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_p\}$ one by one.

Take the largest k s.t. $S_k = \{x_1, \dots, x_k\}$ is linearly independent.

Then we show that $k = p$. Suppose that $k < p$. Then $S_{k+1} = \{x_1, \dots, x_k, x_{k+1}\}$ is linearly dependent.

In that case if $\sum_{i=1}^{k+1} \alpha_i x_i = 0$ $\alpha_1, \dots, \alpha_{k+1} \in \mathbb{F}$ not all zero. Then $\alpha_{k+1} \neq 0$, otherwise S_{k+1} will be linearly independent.

$$x_{k+1} = c_1 x_1 + c_2 x_2 + \dots + c_k x_k.$$

$$Ax_{k+1} = c_1 Ax_1 + c_2 Ax_2 + \dots + c_k Ax_k.$$

$$\lambda_{k+1} x_{k+1} = c_1 \lambda_1 x_1 + c_2 \lambda_2 x_2 + \dots + c_k \lambda_k x_k.$$

$$\lambda_{k+1}(c_1x_1 + c_2x_2 + \cdots + c_kx_k) = c_1\lambda_1x_1 + c_2\lambda_2x_2 + \cdots + c_k\lambda_kx_k.$$

$$(\lambda_{k+1} - \lambda_1)c_1x_1 + \cdots + (\lambda_{k+1} - \lambda_k)c_kx_k = 0$$

$$(\lambda_{k+1} - \lambda_1)c_1 = \cdots = (\lambda_{k+1} - \lambda_k)c_k =$$

$c_1 = \cdots = c_k = 0$ as $\lambda_{k+1} - \lambda_i \neq 0$ for $i = 1, \dots, k$.

Hence x_{k+1} is zero a contradiction. Then $k = p$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then $\prod_{i=1}^n \lambda_i = \det(A)$ and $\sum_{i=1}^n \lambda_i = \text{trace}(A)$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then $\prod_{i=1}^n \lambda_i = \det(A)$ and $\sum_{i=1}^n \lambda_i = \text{trace}(A)$.

Proof: The characteristic polynomial of A is $P_A(x) = \det(xI - A)$ and $\lambda_1, \dots, \lambda_n$ are the roots of $P_A(x)$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then $\prod_{i=1}^n \lambda_i = \det(A)$ and $\sum_{i=1}^n \lambda_i = \text{trace}(A)$.

Proof: The characteristic polynomial of A is $P_A(x) = \det(xI - A)$ and $\lambda_1, \dots, \lambda_n$ are the roots of $P_A(x)$.

Then $P_A(x) = (x - \lambda_1) \cdots (x - \lambda_n)$.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then $\prod_{i=1}^n \lambda_i = \det(A)$ and $\sum_{i=1}^n \lambda_i = \text{trace}(A)$.

Proof: The characteristic polynomial of A is $P_A(x) = \det(xI - A)$ and $\lambda_1, \dots, \lambda_n$ are the roots of $P_A(x)$.

Then $P_A(x) = (x - \lambda_1) \cdots (x - \lambda_n)$.

Then equating the co-efficient of x^{n-1} and constant term from both sides.

-
- Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then $\prod_{i=1}^n \lambda_i = \det(A)$ and $\sum_{i=1}^n \lambda_i = \text{trace}(A)$.

Proof: The characteristic polynomial of A is $P_A(x) = \det(xI - A)$ and $\lambda_1, \dots, \lambda_n$ are the roots of $P_A(x)$.

Then $P_A(x) = (x - \lambda_1) \cdots (x - \lambda_n)$.

Then equating the co-efficient of x^{n-1} and constant term from both sides.

Then we have $\sum_{i=1}^n \lambda_i = \text{trace}(A)$ and $\prod_{i=1}^n \lambda_i = \det(A)$.

Let $A \in \mathbb{M}_n(\mathbb{F})$. The $\det(A)$ and $\text{trace}(A)$ are known to you. But you cannot write $\det(A)$ is the product of the eigenvalues of A and $\text{trace}(A)$ is the sum of the eigenvalues of A . Because A may not have n number of eigenvalues.

For example $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & -2 & 0 \end{bmatrix}$ in $\mathbb{M}_n(\mathbb{R})$. It has exactly one eigenvalue which is 1. The $\det(A) = 4$ which is not the product of the eigenvalues of A .

If $A \in \mathbb{M}_n(\mathbb{F})$ and the $\det(A)$ and $\text{trace}(A)$ are known to you. Then it is always true $\det(A)$ is the product of the eigenvalues of A and $\text{trace}(A)$ is the sum of the eigenvalues of A .

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ be an eigenvalue of A . Let $E_\lambda(A) = \{x \in \mathbb{F}^n : (A - \lambda I)x = 0\}$ = set of all solutions of the homogeneous equations $(A - \lambda I)x = 0$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ be an eigenvalue of A . Let $E_\lambda(A) = \{x \in \mathbb{F}^n : (A - \lambda I)x = 0\}$ = set of all solutions of the homogeneous equations $(A - \lambda I)x = 0$.
 - $E_\lambda(A)$ is a subspace of \mathbb{F}^n .

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ be an eigenvalue of A . Let $E_\lambda(A) = \{x \in \mathbb{F}^n : (A - \lambda I)x = 0\}$ = set of all solutions of the homogeneous equations $(A - \lambda I)x = 0$.
 - $E_\lambda(A)$ is a subspace of \mathbb{F}^n .
 - Then $E_\lambda(A)$ is called the **eigenspace** of A corresponding to the eigenvalue of λ .

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ be an eigenvalue of A . Let $E_\lambda(A) = \{x \in \mathbb{F}^n : (A - \lambda I)x = 0\}$ = set of all solutions of the homogeneous equations $(A - \lambda I)x = 0$.
 - $E_\lambda(A)$ is a subspace of \mathbb{F}^n .
 - Then $E_\lambda(A)$ is called the **eigenspace** of A corresponding to the eigenvalue of λ .
 - The subspace E_λ is a finite dimensional for each eigenvalue λ of A . The dimension of E_λ is called the **geometric multiplicity** of λ with respect to A .

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$. Then

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$. Then

$$E_{(\lambda=1)} = \left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} \right\}$$

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$. Then

$$E_{(\lambda=1)} = \left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} \right\}$$

$\dim(E_{(\lambda=1)}) = 1$. Hence the geometric multiplicity of $\lambda = 1$ is 1.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$. Then

$$E_{(\lambda=1)} = \left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} \right\}$$

$\dim(E_{(\lambda=1)}) = 1$. Hence the geometric multiplicity of $\lambda = 1$ is 1.

- An eigenvalue λ of A is said to be **regular** if the algebraic and the geometric multiplicities of λ are equal.

-
- Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. We have already calculated eigenvalues of A which are $1, -1, 4$. Then

$$E_{(\lambda=1)} = \left\{ k \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} : k \in \mathbb{R} \right\}$$

$\dim(E_{(\lambda=1)}) = 1$. Hence the geometric multiplicity of $\lambda = 1$ is 1.

- An eigenvalue λ of A is said to be **regular** if the algebraic and the geometric multiplicities of λ are equal.
- λ is said to be a **Simple** eigenvalue of A if the algebraic multiplicity of λ with respect to A is 1.

-
- **[Theorem]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and λ be an eigenvalue of A . Then the geometric multiplicity of λ is $n - \text{rank}(A - \lambda I)$.

-
- **[Theorem]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and λ be an eigenvalue of A . Then the geometric multiplicity of λ is $n - \text{rank}(A - \lambda I)$.

Consider the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 4 & 2 & 0 \\ 6 & 0 & 2 \end{bmatrix}$. The eigenvalues of A are 2, 2, 2. The algebraic multiplicity of 2 is 3.

-
- **[Theorem]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and λ be an eigenvalue of A . Then the geometric multiplicity of λ is $n - \text{rank}(A - \lambda I)$.

Consider the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 4 & 2 & 0 \\ 6 & 0 & 2 \end{bmatrix}$. The eigenvalues of A are 2, 2, 2. The algebraic multiplicity of 2 is 3.

$A - 2I = \begin{bmatrix} 0 & 0 & 0 \\ 4 & 0 & 0 \\ 6 & 0 & 0 \end{bmatrix}$. The rank of $A - 2I$ is 1. Hence the geometric multiplicity of 2 is $3 - 1 = 2$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ_1 and λ_2 be two distinct eigenvalues of A . Then $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ_1 and λ_2 be two distinct eigenvalues of A . Then $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

Proof: Let $x \in E_{\lambda_1}(A) \cap E_{\lambda_2}(A)$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ_1 and λ_2 be two distinct eigenvalues of A . Then $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

Proof: Let $x \in E_{\lambda_1}(A) \cap E_{\lambda_2}(A)$.

Then $x \in E_{\lambda_1}(A)$ implies $Ax = \lambda_1 x$

and $x \in E_{\lambda_2}(A)$ implies $Ax = \lambda_2 x$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ_1 and λ_2 be two distinct eigenvalues of A . Then $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

Proof: Let $x \in E_{\lambda_1}(A) \cap E_{\lambda_2}(A)$.

Then $x \in E_{\lambda_1}(A)$ implies $Ax = \lambda_1 x$

and $x \in E_{\lambda_2}(A)$ implies $Ax = \lambda_2 x$.

This implies $\lambda_1 x = \lambda_2 x$.

-
- Let $A \in \mathbb{M}_{n \times n}(\mathbb{F})$ and let λ_1 and λ_2 be two distinct eigenvalues of A . Then $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

Proof: Let $x \in E_{\lambda_1}(A) \cap E_{\lambda_2}(A)$.

Then $x \in E_{\lambda_1}(A)$ implies $Ax = \lambda_1 x$

and $x \in E_{\lambda_2}(A)$ implies $Ax = \lambda_2 x$.

This implies $\lambda_1 x = \lambda_2 x$.

Since λ_1 and λ_2 are distinct, then $x = 0$. Hence $E_{\lambda_1}(A) \cap E_{\lambda_2}(A) = \{0\}$.

Discussions: Let $B \in \mathbb{M}_n(\mathbb{F})$. Then we can write B in the following way

$B = [b_1 : b_2 : \cdots : b_n]$ where b_i is the i th column of B for $i = 1, \dots, n$.

Discussions: Let $B \in \mathbb{M}_n(\mathbb{F})$. Then we can write B in the following way

$B = [b_1 : b_2 : \cdots : b_n]$ where b_i is the i th column of B for $i = 1, \dots, n$.

Let $A \in \mathbb{M}_n(\mathbb{F})$. Then

Discussions: Let $B \in \mathbb{M}_n(\mathbb{F})$. Then we can write B in the following way

$B = [b_1 : b_2 : \cdots : b_n]$ where b_i is the i th column of B for $i = 1, \dots, n$.

Let $A \in \mathbb{M}_n(\mathbb{F})$. Then

$$AB = A[b_1 : b_2 : \cdots : b_n]$$

$$= [Ab_1 : Ab_2 : \cdots : Ab_n] \text{ where } Ab_i \text{ is the } i\text{th column of } AB \text{ for } i = 1, \dots, n.$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.
Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

Since $E_\lambda(A)$ is a subspace of \mathbb{F}^n . We extend \mathcal{B}_1 to a basis for \mathbb{F}^n , say, $\mathcal{B} = \{x_1, \dots, x_m, x_{m+1}, \dots, x_n\}$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

Since $E_\lambda(A)$ is a subspace of \mathbb{F}^n . We extend \mathcal{B}_1 to a basis for \mathbb{F}^n , say, $\mathcal{B} = \{x_1, \dots, x_m, x_{m+1}, \dots, x_n\}$.

Thus $P = [x_1 : x_2 : \dots : x_n]$ is a non-singular matrix. Therefore

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

Since $E_\lambda(A)$ is a subspace of \mathbb{F}^n . We extend \mathcal{B}_1 to a basis for \mathbb{F}^n , say, $\mathcal{B} = \{x_1, \dots, x_m, x_{m+1}, \dots, x_n\}$.

Thus $P = [x_1 : x_2 : \dots : x_n]$ is a non-singular matrix. Therefore

$$P^{-1}AP = P^{-1}A[x_1 : x_2 : \dots : x_n].$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

Since $E_\lambda(A)$ is a subspace of \mathbb{F}^n . We extend \mathcal{B}_1 to a basis for \mathbb{F}^n , say, $\mathcal{B} = \{x_1, \dots, x_m, x_{m+1}, \dots, x_n\}$.

Thus $P = [x_1 : x_2 : \dots : x_n]$ is a non-singular matrix. Therefore

$$P^{-1}AP = P^{-1}A[x_1 : x_2 : \dots : x_n].$$

$$P^{-1}AP = P^{-1}[Ax_1 : Ax_2 : \dots : Ax_n].$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let λ be an eigenvalue of A . Then the algebraic multiplicity of λ is greater or equal to the geometric multiplicity of λ .

Proof: Let the geometric multiplicity of λ be m . That is $\dim(E_\lambda(A)) = m$.

Let $\mathcal{B}_1 = \{x_1, \dots, x_m\}$ be a basis of $E_\lambda(A)$.

Since $E_\lambda(A)$ is a subspace of \mathbb{F}^n . We extend \mathcal{B}_1 to a basis for \mathbb{F}^n , say, $\mathcal{B} = \{x_1, \dots, x_m, x_{m+1}, \dots, x_n\}$.

Thus $P = [x_1 : x_2 : \dots : x_n]$ is a non-singular matrix. Therefore

$$P^{-1}AP = P^{-1}A[x_1 : x_2 : \dots : x_n].$$

$$P^{-1}AP = P^{-1}[Ax_1 : Ax_2 : \dots : Ax_n].$$

$$= P^{-1}[\lambda x_1 : \dots : \lambda x_m : Ax_{m+1} : \dots : Ax_n].$$

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

$$= [\lambda P^{-1}x_1 : \dots : \lambda P^{-1}x_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

$$= [\lambda P^{-1}x_1 : \dots : \lambda P^{-1}x_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$= [\lambda e_1 : \dots : \lambda e_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

$$= [\lambda P^{-1}x_1 : \dots : \lambda P^{-1}x_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$= [\lambda e_1 : \dots : \lambda e_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$P^{-1}AP = \begin{bmatrix} \lambda I_m & B \\ 0 & C \end{bmatrix} \text{ for some matrices } B \text{ and } C.$$

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

$$= [\lambda P^{-1}x_1 : \dots : \lambda P^{-1}x_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$= [\lambda e_1 : \dots : \lambda e_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$P^{-1}AP = \begin{bmatrix} \lambda I_m & B \\ 0 & C \end{bmatrix} \text{ for some matrices } B \text{ and } C.$$

$$\text{Hence } P_A(x) = P_{P^{-1}AP}(x) = (x - \lambda)^m P_C(x).$$

We can show that $P^{-1}(\lambda x_j) = \lambda P^{-1}x_j = \lambda e_j$ (Here x_j is the j th column of P and P^{-1} is the inverse of P) $j = 1, \dots, m$. Then

$$= [\lambda P^{-1}x_1 : \dots : \lambda P^{-1}x_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$= [\lambda e_1 : \dots : \lambda e_m : P^{-1}Ax_{m+1} : \dots : P^{-1}Ax_n].$$

$$P^{-1}AP = \begin{bmatrix} \lambda I_m & B \\ 0 & C \end{bmatrix} \text{ for some matrices } B \text{ and } C.$$

$$\text{Hence } P_A(x) = P_{P^{-1}AP}(x) = (x - \lambda)^m P_C(x).$$

So the algebraic multiplicity of λ with respect to A is at least m and the theorem follows.

-
- Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

-
- Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.

• Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.
2. You can easily find the determinant.

• Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.
2. You can easily find the determinant.
3. You can easily find the power of the matrix.

• Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.
2. You can easily find the determinant.
3. You can easily find the power of the matrix.
4. You can easily find the solutions of the equations $Ax = 0$.

• Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.
2. You can easily find the determinant.
3. You can easily find the power of the matrix.
4. You can easily find the solutions of the equations $Ax = 0$.
6. You can easily find the eigenvalues and eigenvectors.

• Suppose you have a diagonal matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$.

1. You can easily find the rank.
2. You can easily find the determinant.
3. You can easily find the power of the matrix.
4. You can easily find the solutions of the equations $Ax = 0$.
6. You can easily find the eigenvalues and eigenvectors.

• Suppose we have a non-diagonal matrix A and if we are able to show that A is similar to a diagonal matrix, then we can easily find the above information for A .

-
- **[Definition:]** A matrix is said to be diagonalizable

-
- **[Definition:]** A matrix is said to be diagonalizable if there exists a nonsingular matrix P such that

-
- **[Definition:]** A matrix is said to be diagonalizable if there exists a nonsingular matrix P such that $P^{-1}AP$ is a diagonal matrix. That is, A is similar to a diagonal matrix.
 - **[Example:]** Every diagonal matrix is diagonalizable. Use the argument every matrix is similar to itself.

-
- **[Definition:]** A matrix is said to be diagonalizable if there exists a nonsingular matrix P such that $P^{-1}AP$ is a diagonal matrix. That is, A is similar to a diagonal matrix.
 - **[Example:]** Every diagonal matrix is diagonalizable. Use the argument every matrix is similar to itself.

Consider the matrix $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.

-
- **[Definition:]** A matrix is said to be diagonalizable if there exists a nonsingular matrix P such that $P^{-1}AP$ is a diagonal matrix. That is, A is similar to a diagonal matrix.
 - **[Example:]** Every diagonal matrix is diagonalizable. Use the argument every matrix is similar to itself.

Consider the matrix $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.

Take $P = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$.

-
- **[Definition:]** A matrix is said to be diagonalizable if there exists a nonsingular matrix P such that $P^{-1}AP$ is a diagonal matrix. That is, A is similar to a diagonal matrix.
 - **[Example:]** Every diagonal matrix is diagonalizable. Use the argument every matrix is similar to itself.

Consider the matrix $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.

Take $P = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$.

Then you can check that $P^{-1}AP = \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix}$. Hence A is diagonalizable matrix.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

Then $PD = P[d_1e_1 : \cdots : d_ne_n]$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

Then $PD = P[d_1e_1 : \cdots : d_ne_n]$

$$PD = [d_1Pe_1 : \cdots : d_nPe_n]$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

Then $PD = P[d_1e_1 : \cdots : d_ne_n]$

$$PD = [d_1Pe_1 : \cdots : d_nPe_n]$$

$$PD = [d_1P_1 : \cdots : d_nP_n]$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

Then $PD = P[d_1 e_1 : \cdots : d_n e_n]$

$$PD = [d_1 P e_1 : \cdots : d_n P e_n]$$

$$PD = [d_1 P_1 : \cdots : d_n P_n]$$

and $AP = [AP_1 : \cdots : AP_n]$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{F})$ and let $P^{-1}AP = D$ where $D = (d_i)$ is a diagonal matrix. Then d_i is an eigenvalue of A and i th column of P is an eigenvector corresponding to d_i .

Proof: We have $P^{-1}AP = D$. That is $AP = PD$.

Let $P = [P_1 : \cdots : P_n]$ where P_i is the i th column of P .

Then $PD = P[d_1 e_1 : \cdots : d_n e_n]$

$$PD = [d_1 P e_1 : \cdots : d_n P e_n]$$

$$PD = [d_1 P_1 : \cdots : d_n P_n]$$

and $AP = [AP_1 : \cdots : AP_n]$.

Therefore $AP_i = d_i P_i$ for $i = 1, \dots, n$.

Since P is non-singular, we have each P_i is non-zero for each $i = 1, \dots, n$.

Since P is non-singular, we have each P_i is non-zero for each $i = 1, \dots, n$.

This implies that d_i is an eigenvalue of A and corresponding eigenvector P_i for $i = 1, \dots, n$

-
- There is a square matrix which is not diagonalizable.

-
- There is a square matrix which is not diagonalizable.

For example, consider the matrix $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.

-
- There is a square matrix which is not diagonalizable.

For example, consider the matrix $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.

It is not diagonalizable.

-
- There is a square matrix which is not diagonalizable.

For example, consider the matrix $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.

It is not diagonalizable.

If A is diagonalizable then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix and each diagonal entry is the eigenvalue of A .

-
- There is a square matrix which is not diagonalizable.

For example, consider the matrix $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.

It is not diagonalizable.

If A is diagonalizable then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix and each diagonal entry is the eigenvalue of A .

Hence $D = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. Therefore $P^{-1}AP = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. This implies that $A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ which is not possible.

-
- We have seen that all the matrices are not diagonalizable. Then the following questions are natural.

-
- We have seen that all the matrices are not diagonalizable. Then the following questions are natural.

Q1. How do we check whether a matrix is diagonalizable or not?

-
- We have seen that all the matrices are not diagonalizable. Then the following questions are natural.

Q1. How do we check whether a matrix is diagonalizable or not?

Q2. If A is diagonalizable, then how do I calculate such P matrix?

Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively.

Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively.

Then the characteristic polynomial of A is $\prod_{i=1}^k (x - \lambda_i)^{m_i}$. This implies that $m_1 + \dots + m_k = n$.

Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively.

Then the characteristic polynomial of A is $\prod_{i=1}^k (x - \lambda_i)^{m_i}$. This implies that $m_1 + \dots + m_k = n$.

Consider the following matrix $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$ in $\mathbb{M}_n(\mathbb{R})$. Then A has exactly one eigenvalue which is 1 with algebraic multiplicity 1.

So the above argument is not true if we consider some other field \mathbb{F} instead of \mathbb{C} .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

We first assume that A is diagonalizable.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

We first assume that A is diagonalizable.

Then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

We first assume that A is diagonalizable.

Then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix.

Using previous theorem, each column of the matrix P is an eigenvector of A .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

We first assume that A is diagonalizable.

Then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix.

Using previous theorem, each column of the matrix P is an eigenvector of A .

Since P is non-singular, the columns vectors of P are linearly independent.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

We first assume that A is diagonalizable.

Then there exists a non-singular matrix P such that $P^{-1}AP = D$ where D is a diagonal matrix.

Using previous theorem, each column of the matrix P is an eigenvector of A .

Since P is non-singular, the columns vectors of P are linearly independent.

Hence A has n linearly independent eigenvectors.

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

$$AP = [Ax_1 : Ax_2 : \dots : Ax_n]$$

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

$$AP = [Ax_1 : Ax_2 : \dots : Ax_n]$$

$$AP = [\lambda_1 x_1 : \dots : \lambda_n x_n]$$

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

$$AP = [Ax_1 : Ax_2 : \dots : Ax_n]$$

$$AP = [\lambda_1 x_1 : \dots : \lambda_n x_n]$$

$$AP = [\lambda_1 P e_1 : \dots : \lambda_n P e_n]$$

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

$$AP = [Ax_1 : Ax_2 : \dots : Ax_n]$$

$$AP = [\lambda_1 x_1 : \dots : \lambda_n x_n]$$

$$AP = [\lambda_1 P e_1 : \dots : \lambda_n P e_n]$$

$$AP = P[\lambda_1 e_1 : \dots : \lambda_n e_n].$$

We now assume that A has n linearly independent eigenvectors those are x_1, \dots, x_n .

Take the matrix $P = [x_1 : x_2 : \dots : x_n]$. Then P is non-singular. Therefore

$$AP = A[x_1 : x_2 : \dots : x_n]$$

$$AP = [Ax_1 : Ax_2 : \dots : Ax_n]$$

$$AP = [\lambda_1 x_1 : \dots : \lambda_n x_n]$$

$$AP = [\lambda_1 P e_1 : \dots : \lambda_n P e_n]$$

$$AP = P[\lambda_1 e_1 : \dots : \lambda_n e_n]. \quad P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & 0 & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_n \end{bmatrix}. \quad \text{Hence } A$$

is diagonalizable.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Suppose that there exists $1 \leq t \leq k$ such that the geometric multiplicity of λ_t is strictly less than m_t .

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Suppose that there exists $1 \leq t \leq k$ such that the geometric multiplicity of λ_t is strictly less than m_t .

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$ and $i \neq t$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Suppose that there exists $1 \leq t \leq k$ such that the geometric multiplicity of λ_t is strictly less than m_t .

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$ and $i \neq t$.

The eigenspace $E_{\lambda_t}(A)$ gives us at most $m_t - 1$ linearly independent eigenvectors.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Suppose that there exists $1 \leq t \leq k$ such that the geometric multiplicity of λ_t is strictly less than m_t .

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$ and $i \neq t$.

The eigenspace $E_{\lambda_t}(A)$ gives us at most $m_t - 1$ linearly independent eigenvectors.

So we have at most $n - 1$ eigenvector. A contradiction that a diagonalizable matrix must have n linearly eigenvectors.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k , respectively. Then A is diagonalizable if and only if the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Proof: We first assume that A is diagonalizable. To show the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

Suppose that there exists $1 \leq t \leq k$ such that the geometric multiplicity of λ_t is strictly less than m_t .

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$ and $i \neq t$.

The eigenspace $E_{\lambda_t}(A)$ gives us at most $m_t - 1$ linearly independent eigenvectors.

So we have at most $n - 1$ eigenvectors. A contradiction that a diagonalizable matrix must have n linearly independent eigenvectors.

Contradiction because we assume that the geometric multiplicity of λ_t is strictly less than m_t . Hence the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

We now assume that the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

We now assume that the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$.

We now assume that the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

The eigenspace $E_{\lambda_i}(A)$ gives us at most m_i linearly independent eigenvectors for $i = 1, \dots, k$.

So we have n linearly independent eigenvectors. Hence A is diagonalizable.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Proof: We first assume that A is diagonalizable. Then the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Proof: We first assume that A is diagonalizable. Then the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

We have already proved that $E_{\lambda_i}(A) \cap E_{\lambda_j}(A) = \{0\}$ for $1 \leq i, j \leq k$ and $i \neq j$.

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Proof: We first assume that A is diagonalizable. Then the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

We have already proved that $E_{\lambda_i}(A) \cap E_{\lambda_j}(A) = \{0\}$ for $1 \leq i, j \leq k$ and $i \neq j$.

Then $\dim(E_{\lambda_1}(A) + E_{\lambda_2}(A) + E_{\lambda_3}(A) + \dots + E_{\lambda_k}(A)) = \sum_{i=1}^k \dim(E_{\lambda_i}(A)) =$

$$\sum_{i=1}^k m_i = n.$$

-
- **[Theorem:]** Let $A \in \mathbb{M}_n(\mathbb{C})$ and let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with algebraic multiplicities m_1, \dots, m_k respectively. Then A is diagonalizable if and only if $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Proof: We first assume that A is diagonalizable. Then the geometric multiplicity of λ_i is m_i for $i = 1, \dots, k$.

We have already proved that $E_{\lambda_i}(A) \cap E_{\lambda_j}(A) = \{0\}$ for $1 \leq i, j \leq k$ and $i \neq j$.

Then $\dim(E_{\lambda_1}(A) + E_{\lambda_2}(A) + E_{\lambda_3}(A) + \dots + E_{\lambda_k}(A)) = \sum_{i=1}^k \dim(E_{\lambda_i}(A)) = \sum_{i=1}^k m_i = n$.

Hence $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \dots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Assume that $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Assume that $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Let B_i be a basis of $E_{\lambda_i}(A)$ for $i = 1, \dots, k$.

Assume that $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Let B_i be a basis of $E_{\lambda_i}(A)$ for $i = 1, \dots, k$.

It can be proved that $\cup_{i=1}^k B_i$ is a basis of $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Assume that $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

Let B_i be a basis of $E_{\lambda_i}(A)$ for $i = 1, \dots, k$.

It can be proved that $\cup_{i=1}^k B_i$ is a basis of $E_{\lambda_1}(A) \oplus E_{\lambda_2}(A) \oplus E_{\lambda_3}(A) \oplus \cdots \oplus E_{\lambda_k}(A) = \mathbb{C}^n$.

The cardinality of $\cup_{i=1}^k B_i$ is n . Hence we have n linearly independent vectors. Thus A is diagonalizable.

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}.$$

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}.$$

Eigenvalues of A are 1 and 2 with algebraic multiplicity 1 and 2, respectively.

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}.$$

Eigenvalues of A are 1 and 2 with algebraic multiplicity 1 and 2, respectively.

Eigenspace corresponding to $\lambda = 1$ is

$$\left\{ k \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} : k \in \mathbb{R} \right\}. \text{ Hence the geometric multiplicity of } \lambda = 1 \text{ is } 1.$$

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}.$$

Eigenvalues of A are 1 and 2 with algebraic multiplicity 1 and 2, respectively.

Eigenspace corresponding to $\lambda = 1$ is

$$\left\{ k \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} : k \in \mathbb{R} \right\}. \text{ Hence the geometric multiplicity of } \lambda = 1 \text{ is } 1.$$

Eigenspace corresponding to $\lambda = 2$ is

$$\left\{ k_1 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + k_2 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} : k_1, k_2 \in \mathbb{R} \right\}. \text{ Hence the geometric multiplicity of } \lambda = 2 \text{ is } 2.$$

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}.$$

Eigenvalues of A are 1 and 2 with algebraic multiplicity 1 and 2, respectively.

Eigenspace corresponding to $\lambda = 1$ is

$$\left\{ k \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} : k \in \mathbb{R} \right\}. \text{ Hence the geometric multiplicity of } \lambda = 1 \text{ is } 1.$$

Eigenspace corresponding to $\lambda = 2$ is

$$\left\{ k_1 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + k_2 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} : k_1, k_2 \in \mathbb{R} \right\}. \text{ Hence the geometric multiplicity of } \lambda = 2 \text{ is } 2.$$

We have seen that the algebraic multiplicity equal to the geometric multiplicity for each eigenvalue. Hence A is diagonalizable.