

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the same field \mathbb{F} .

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.
2. Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$, let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be defined by $T(x) = Ax$, $x \in \mathbb{R}^n$. Then T is a linear transformation.

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the **same field** \mathbb{F} .

A mapping $T : \mathbb{V} \rightarrow \mathbb{W}$ is said to be a linear transformation if

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \text{ for all } x, y \in \mathbb{V} \text{ and for all } \alpha, \beta \in \mathbb{F}.$$

- **[Example:]**

1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T(x, y) = x$. Then T is a linear transformation.
2. Let $A \in \mathbb{M}_{m \times n}(\mathbb{R})$, let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be defined by $T(x) = Ax$, $x \in \mathbb{R}^n$. Then T is a linear transformation.
3. Let $T : C[a, b] \rightarrow \mathbb{R}$ be defined by $T(f) = \int_a^b f(x)dx$. Then T is a linear transformation.

-
1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.

-
1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
 2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.
3. Let \mathbb{V} be a vector space over the field \mathbb{F} and let $\lambda \in \mathbb{F}$. Let

1. Let \mathbb{V} and \mathbb{W} be vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be defined by $T(x) = 0, x \in \mathbb{V}$. This transformation is called **Zero transformation**.
2. Let \mathbb{V} be vector space over the field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = x, x \in \mathbb{V}$. This transformation is called **identity transformation**.
3. Let \mathbb{V} be a vector space over the field \mathbb{F} and let $\lambda \in \mathbb{F}$. Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = \lambda x, x \in \mathbb{V}$. This transformation is called **scalar transformation**.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

$T : \mathbb{R} \rightarrow \mathbb{R}$ be a map defined by $T(x) = x + 1$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then $T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$

Proof: We know that $0_{\mathbb{V}} + 0_{\mathbb{V}} = 0_{\mathbb{V}}$.

$$T(0_{\mathbb{V}} + 0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) = T(0_{\mathbb{V}})$$

$$\implies T(0_{\mathbb{V}}) + T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}})) = T(0_{\mathbb{V}}) + (-T(0_{\mathbb{V}}))$$

$$\implies T(0_{\mathbb{V}}) + 0_{\mathbb{W}} = 0_{\mathbb{W}}$$

$$\implies T(0_{\mathbb{V}}) = 0_{\mathbb{W}}$$

$T : \mathbb{R} \rightarrow \mathbb{R}$ be a map defined by $T(x) = x + 1$. Using above theorem you can say that T is not linear.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^k \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}$ be a LT. Then T has the following form.

$$T(x_1, \dots, x_n) = \sum_{i=1}^n \alpha_i x_i \text{ for some } \alpha_i \in \mathbb{R} \text{ for } i = 1, \dots, n \text{ and for all}$$

$$(x_1, \dots, x_n) \in \mathbb{R}^n.$$

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = \alpha_i$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

$$T(x) = \sum_{i=1}^n \alpha_i x_i$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

- **[Theorem:]** Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Then there exist linear transformations $T_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, m$ such that $T(x) = (T_1(x), \dots, T_m(x))$ for all $x \in \mathbb{R}^n$.

Proof: Let $\{e_1, \dots, e_n\}$ be a basis of \mathbb{R}^n .

Let $T(e_i) = (\alpha_1^i, \dots, \alpha_m^i)$ for $i = 1, \dots, n$.

Let $x = [x_1, x_2, \dots, x_n] \in \mathbb{V}$. Then $x = \sum_{i=1}^n x_i e_i$.

$$T(x) = T\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i T(e_i)$$

$$T(x) = \sum_{i=1}^n (\alpha_1^i, \dots, \alpha_m^i) x_i$$

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{i=1}^n x_i \alpha_i^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

Could it be possible to get the linear map explicitly?

$$T(x) = \left(\sum_{i=1}^n x_i \alpha_i^1, \dots, \sum_{i=1}^n x_i \alpha_i^m \right)$$

$$T(x) = (T_1(x), \dots, T_n(x)), \text{ where } T_i(x) = \sum_{j=1}^n x_j \alpha_j^i$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a map defined by $T(x_1, x_2) = (x_2 - x_1, x_1^2, x_2)$.

T is linear or not?

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(e_1) = (1, 1)$ and $T(e_2) = (-1, 1)$.

Could it be possible to get the linear map explicitly?

Answer: Yes. Let $(x_1, x_2) \in \mathbb{R}^2$. Then $(x_1, x_2) = x_1 e_1 + x_2 e_2$.

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

$$\text{Then } T(x_1, x_2) = x_1 T(e_1) + x_2 T(e_2)$$

$$= x_1(1, 1) + x_2(-1, 1)$$

$$= (x_1 - x_2, x_1 + x_2)$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

Could it be possible to get the linear map explicitly?

$$\begin{aligned}\text{Then } T(x_1, x_2) &= x_1 T(e_1) + x_2 T(e_2) \\ &= x_1(1, 1) + x_2(-1, 1) \\ &= (x_1 - x_2, x_1 + x_2)\end{aligned}$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(1, 0) = (1, 1)$.

Could it be possible to get the linear map explicitly?

Answer: No it is not possible.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} .

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} .

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$.

- **[Theorem:]** Let \mathbb{V} be a finite-dimensional vector space over the field \mathbb{F} and let $\{u_1, \dots, u_n\}$ be an **ordered basis** for \mathbb{V} . Let \mathbb{W} be a vector space over the same field \mathbb{F} and let w_1, \dots, w_n be any vectors in \mathbb{W} . Then there is precisely one linear transformation T from \mathbb{V} into \mathbb{W} such that $T(u_j) = w_j$, for $j = 1, \dots, n$.

Proof:

Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$.

Define $T(x) = \sum_{i=1}^n c_i w_i$. It is clear that T is well defined because $x = \sum_{i=1}^n c_i u_i$, this expression unique.

We first show that T is a linear transformation. Take $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$ and $y = \sum_{i=1}^n d_i u_i$.

We first show that T is a linear transformation. Take $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^n c_i u_i$ and $y = \sum_{i=1}^n d_i u_i$.

Let $\alpha, \beta \in \mathbb{F}$. $T(\alpha x + \beta y) = T(\sum_{i=1}^n (\alpha c_i + \beta d_i) u_i)$.

$$\begin{aligned} T(\alpha x + \beta y) &= \sum_{i=1}^n (\alpha c_i + \beta d_i) w_i. \\ &= \alpha \sum_{i=1}^n c_i w_i + \beta \sum_{i=1}^n d_i w_i. \\ &= \alpha T(x) + \beta T(y). \end{aligned}$$

Hence T is linear.

Uniqueness: Suppose that there is another linear transformation U such that $U(u_i) = w_i$.

Uniqueness: Suppose that there is another linear transformation U such that $U(u_i) = w_i$.

To show that $U = T$. Let $x \in \mathbb{V}$. Then $x = \sum_{i=1}^n a_i u_i$. Using definition of T

$$\text{we have } T(x) = T\left(\sum_{i=1}^n a_i u_i\right) = \sum_{i=1}^n a_i w_i.$$

$$U(x) = U\left(\sum_{i=1}^n a_i u_i\right)$$

$$= \sum_{i=1}^n a_i U(u_i) \text{ (applying the definition of linear transformation)}$$

$$= \sum_{i=1}^n a_i w_i.$$

Then $U(x) = T(x)$ for all $x \in \mathbb{V}$. Hence $U = T$.

- [Example]

Take the basis $\{e_1, e_2, e_3\}$ in \mathbb{R}^3 . Take $1, 2, 3 \in \mathbb{R}$. Then using previous theorem we have a unique linear transformation T from \mathbb{R}^3 to \mathbb{R} such that $T(e_1) = 1$, $T(e_2) = 2$, $T(e_3) = 3$ and $T(x_1, x_2, x_3) = x_1 + 2x_2 + 3x_3$.

Take the basis $\{e_1, e_2, e_3\}$ in \mathbb{R}^3 . Take $1, 2, 3 \in \mathbb{R}$. Then using previous theorem we have a unique linear transformation T from \mathbb{R}^3 to \mathbb{R} such that $T(e_1) = 2$, $T(e_2) = 1$, $T(e_3) = 3$ and $T(x_1, x_2, x_3) = 2x_1 + x_2 + 3x_3$. This transformation is different between the previous transformation.

The previous theorem gives a technique to construct a linear transformation from a finite dimensional vector space to another dimensional vector space over the same field \mathbb{F} .

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspaces $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$.

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$. you can easily check that $R(T)$ is a subspace of \mathbb{W} .

• [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

1. $\text{Ker}(T) := \{x \in \mathbb{V} : T(x) = 0\}$. This is called $\text{ker}(T)$. You can easily check that $\text{Ker}(T)$ is a subspace of \mathbb{V} .

The subspace $\text{ker}(T)$ is called the **null space** of T and sometimes it is denoted by $N(T)$.

2. $R(T) := \{T(x) : x \in \mathbb{V}\}$. you can easily check that $R(T)$ is a subspace of \mathbb{W} .

The subspace $R(T)$ is called the **range space** of T .

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

$$\implies N(T) := \text{ls}(\{(1, 1, 1)\})$$

- [Example:] $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a map defined by

$$T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_3)$$

$$N(T) := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : T(x_1, x_2, x_3) = 0\}.$$

$$T(x_1, x_2, x_3) = 0$$

$$\implies (x_1 - x_2, x_1 - x_3) = (0, 0)$$

$$\implies x_1 - x_2 = 0, x_1 - x_3 = 0$$

$$\implies x_1 = x_2, x_1 = x_3$$

$$\implies x_1 = x_2 = x_3 = k, k \in \mathbb{R}$$

$$\implies (x_1, x_2, x_3) = k(1, 1, 1)$$

$$\implies N(T) := \text{ls}(\{(1, 1, 1)\})$$

$$\dim(N(T)) = 1.$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \text{ls}(\{(1, 0), (0, 1)\})$$

$$R(T) := \{T(x) : x \in \mathbb{R}^3\}.$$

Let $y = (y_1, y_2) \in R(T)$. Then there exists $(x_1, x_2, x_3) \in \mathbb{R}^3$ such that $(y_1, y_2) = T(x_1, x_2, x_3)$.

$$(y_1, y_2) = (x_1 - x_2, x_1 - x_3).$$

$$(y_1, y_2) = (x_1 - x_2)(1, 0) + (x_1 - x_3)(0, 1)$$

$$(y_1, y_2) = k_1(1, 0) + k_2(0, 1)$$

$$R(T) := \text{ls}(\{(1, 0), (0, 1)\})$$

$$\dim(R(T)) = 2$$

-
- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the field \mathbb{F} .

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the field \mathbb{F} .

The $\dim(R(T))$ is called the **rank** of T and $\dim(N(T))$ is called the **nullity** of T .

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$. This implies $T(x_1 - x_2) = 0$. Hence $x_1 - x_2 \in \text{Ker}(T)$.

- [Definition:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.
 1. T is called one-one (injective) if $T(x_1) = T(x_2) \implies x_1 = x_2$ for all $x_1, x_2 \in \mathbb{V}$.
 2. T is called onto (surjective) if $T(\mathbb{V}) = \mathbb{W}$, that is $R(T) = \mathbb{W}$.
- [Theorem:] Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is one-one if and only if $\text{Ker}(T) = \{0\}$.

Proof: We first assume that T is one-one.

To show $\text{Ker}(T) = \{0\}$. Let $x \in \text{ker}(T)$. Then $T(x) = 0$ and we know that $T(0) = 0$. Since T is one-one, then $x = 0$. Hence $\text{Ker}(T) = \{0\}$.

We now assume that $\text{ker}(T) = \{0\}$.

To show T is one-one. Let $T(x_1) = T(x_2)$. This implies $T(x_1 - x_2) = 0$. Hence $x_1 - x_2 \in \text{Ker}(T)$. Therefore $x_1 - x_2 = 0$. This implies $x_1 = x_2$.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

- **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } T(u_1), \dots, T(u_n) \text{ are LI.}$$

• **[Theorem:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then

If u_1, \dots, u_n are in \mathbb{V} such that $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} , then u_1, \dots, u_n are linearly independent in \mathbb{V} .

Given that $T(u_1), \dots, T(u_n)$ are LI. To show u_1, \dots, u_n are LI.

$$c_1 u_1 + \dots + c_n u_n = 0_{\mathbb{V}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}$$

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } T(u_1), \dots, T(u_n) \text{ are LI.}$$

Converse is not true in general. That is if u_1, \dots, u_n are LI, then $T(u_1), \dots, T(u_n)$ may or may not be LI.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$$c_i u_1 + \dots + c_n u_n = 0_{\mathbb{V}} \text{ as } T \text{ is one-one.}$$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a LT defined by $T(x_1, x_2) = (x_1 - x_2, x_2 - x_1)$. Take $u_1 = (1, 0)$ and $u_2 = (1, 1)$. Notice that u_1, u_2 are LI. But $T(u_1) = (1, -1)$ and $T(u_2) = (0, 0)$ are LD.

- **[Theorem:]** If T is one-one and u_1, \dots, u_n are linearly independent in \mathbb{V} , then $T(u_1), \dots, T(u_n)$ are linearly independent in \mathbb{W} .

Proof: Given u_1, \dots, u_n are LI and T is one-one. To show $T(u_1), \dots, T(u_n)$ are LI.

$$c_1 T(u_1) + \dots + c_n T(u_n) = 0_{\mathbb{W}}.$$

$$T(c_1 u_1 + \dots + c_n u_n) = 0_{\mathbb{W}}.$$

$$c_i u_1 + \dots + c_n u_n = 0_{\mathbb{V}} \text{ as } T \text{ is one-one.}$$

$$c_1 = c_2 = \dots = c_n = 0 \text{ as } u_1, \dots, u_n \text{ are LI}$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$T(x) = T(c_1 u_1 + \dots + c_n u_n)$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$\begin{aligned} T(x) &= T(c_1 u_1 + \dots + c_n u_n) \\ &= c_1 T(u_1) + \dots + c_n T(u_n) \end{aligned}$$

- **[Rank-Nullity Theorem]** Let \mathbb{V} be a finite dimensional vector space. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then $\dim(\mathbb{V}) = \text{nullity}(T) + \text{rank}(T)$.

Proof: Since \mathbb{V} is finite dimensional, then $\text{Ker}(T)$ is finite dimensional.

Let $\dim(\mathbb{V}) = n$ and let $\text{Ker}(T) = k$.

Let $\{u_1, \dots, u_k\}$ be a basis of $\dim(\text{Ker}(T))$.

Using extension theorem we extend $\{u_1, \dots, u_k\}$ to a basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$.

Let $y \in R(T)$. Then there exists $x \in \mathbb{V}$ such that $T(x) = y$.

Since $x \in \mathbb{V}$, then there exist unique $c_1, \dots, c_n \in \mathbb{F}$ such that $x = c_1 u_1 + \dots + c_n u_n$.

$$T(x) = T(c_1 u_1 + \dots + c_n u_n)$$

$$= c_1 T(u_1) + \dots + c_n T(u_n) = c_{k+1} T(u_{k+1}) + \dots + c_n T(u_n)$$

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Each vector of $T(X)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Since $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$ is basis of \mathbb{V} . Then $a_1 = \dots = a_{n-k} = 0$.

Each vector of $T(x)$ is a linear combination of $T(u_{k+1}), \dots, T(u_n)$ and $T(u_{k+1}), \dots, T(u_n) \in R(T)$. Hence $\text{ls}(\{T(u_{k+1}), \dots, T(u_n)\}) = R(T)$

To show that $T(u_{k+1}), \dots, T(u_n)$ are LI. Take $a_1 T(u_{k+1}) + \dots + a_{n-k} T(u_n) = 0$.

$$T(a_1 u_{k+1} + \dots + a_{n-k} u_n) = 0.$$

Then $a_1 u_{k+1} + \dots + a_{n-k} u_n \in \text{Ker}(T)$.

$$a_1 u_{k+1} + \dots + a_{n-k} u_n = b_1 u_1 + \dots + b_k u_k$$

$$a_1 u_{k+1} + \dots + a_{n-k} u_n - b_1 u_1 - \dots - b_k u_k = 0$$

Since $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$ is basis of \mathbb{V} . Then $a_1 = \dots = a_{n-k} = 0$.

Therefore $T(u_{k+1}), \dots, T(u_n)$ are LI. Hence $\{T(u_{k+1}), \dots, T(u_n)\}$ is a basis of $R(T)$. Then $\dim(R(T)) = n - k$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Let $\mathbb{S} = \text{ls}(\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\})$. So \mathbb{S} is a subspace of \mathbb{V} and $\dim(\mathbb{S}) = m + k + 1$.

Note: Rank-nullity theorem is true only when \mathbb{V} is finite dimensional. Otherwise there is no meaning of rank nullity theorem.

Question: Is there a linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ are finite?

Answer: Not possible to have such type of LT. Let $nullity(T) = k$ and $rank(T) = m$.

Let $\{u_1, \dots, u_k\}$ be a basis of $Ker(T)$. We extend $\{u_1, \dots, u_k\}$ to a LI set of $m + k$ vectors (this is possible as \mathbb{V} is infinite dimensional) which is $\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\}$.

Let $\mathbb{S} = \text{ls}(\{u_1, \dots, u_k, \dots, u_{k+1}, \dots, u_{m+k+1}\})$. So \mathbb{S} is a subspace of \mathbb{V} and $\dim(\mathbb{S}) = m + k + 1$.

$T_{\mathbb{S}}$ is a LT from \mathbb{S} to \mathbb{W} . Then $Ker(T_{\mathbb{S}}) \subseteq Ker(T)$ and $R(T_{\mathbb{S}}) \subseteq R(T)$.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

1. T is one-one.

$nullity(T_{\mathbb{S}}) \leq k$ and $rank(T_{\mathbb{S}}) \leq m$. Then $nullity(T_{\mathbb{S}}) + rank(T_{\mathbb{S}}) \leq k + m$ and $\dim(\mathbb{S}) = m + k + 1$. Then rank nullity theorem is not true on \mathbb{S} .

Therefore there is no linear transformation T from an infinite dimensional vector space \mathbb{V} to another vectors space \mathbb{W} such that $rank(T)$ and $nullity(T)$ is finite.

Applications of Rank-Nullity Theorem: Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a LT. Then the following are equivalent.

1. T is one-one.
2. T is onto.

-
- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) < \dim(\mathbb{W})$. Then there is no onto linear transformation from \mathbb{V} to \mathbb{W} .

- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) < \dim(\mathbb{W})$. Then there is no onto linear transformation from \mathbb{V} to \mathbb{W} .
- Let \mathbb{V} and \mathbb{W} be two vector spaces over the field \mathbb{F} such that $\dim(\mathbb{V}) > \dim(\mathbb{W})$. Then there is no one-one linear transformation from \mathbb{V} to \mathbb{W} .

- **[Definition:]** Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is said to be isomorphism if T is bijective (one-one+onto).

- **[Example:]**

1. Let \mathbb{V} be a vector space over \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = \alpha x$, $\alpha \neq 0$. Then T is an isomorphism.

2. Let $\mathbb{V} = M_{n \times m}(\mathbb{R})$ be the set of $n \times m$ matrices with real entries and let $\mathbb{W} = \mathbb{R}^{mn}$. Define $T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$T(A) = (a_{11}, \dots, a_{1m}, a_{21}, \dots, a_{2m}, \dots, a_{n1}, \dots, a_{nm}).$$

Here $A = (a_{ij}) \in \mathbb{V}$. Then T is an isomorphism.

4. Let $\mathbb{V} = M_{n \times n}(\mathbb{R})$ be the set of $n \times n$ matrices with real entries. Define $T : \mathbb{V} \rightarrow \mathbb{R}$ by $T(A) = \text{trace}(A)$. Then T is not an isomorphism as T is not one-one.

5. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation define by $T(x_1, x_2) = (x_1, x_1 - x_2)$. Then T is an isomorphism.

- **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are said to be isomorphic if there is an isomorphism from \mathbb{V} to \mathbb{W} .

- **[Example:]**

1. \mathbb{R}^n and \mathbb{R}^m are isomorphic if and only if $m = n$.
2. $\mathbb{R}^{m \times n}$ are isomorphic to $\mathbb{V} = M_{n \times m}(\mathbb{R})$, the set of $n \times m$ matrices with real entries.
3. \mathbb{R}^n is isomorphic to $\mathbb{P}_n(x, \mathbb{R})$, set of all real polynomials of degree at most n .

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic.

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic. Let T be an isomorphism from \mathbb{V} to \mathbb{W} .

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic. Let T be an isomorphism from \mathbb{V} to \mathbb{W} . Let $\{u_1, \dots, u_n\}$ be a basis of \mathbb{V} .

Since T is one-one, then $\{T(u_1), \dots, T(u_n)\}$ is linearly independent.

$$\mathbb{W} = R(T) = \text{ls}(\{T(u_1), \dots, T(u_n)\}).$$

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic. Let T be an isomorphism from \mathbb{V} to \mathbb{W} . Let $\{u_1, \dots, u_n\}$ be a basis of \mathbb{V} .

Since T is one-one, then $\{T(u_1), \dots, T(u_n)\}$ is linearly independent.

$\mathbb{W} = R(T) = \text{ls}(\{T(u_1), \dots, T(u_n)\})$. Hence $\{T(u_1), \dots, T(u_n)\}$ is a basis of \mathbb{W} .

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic. Let T be an isomorphism from \mathbb{V} to \mathbb{W} . Let $\{u_1, \dots, u_n\}$ be a basis of \mathbb{V} .

Since T is one-one, then $\{T(u_1), \dots, T(u_n)\}$ is linearly independent.

$\mathbb{W} = R(T) = \text{ls}(\{T(u_1), \dots, T(u_n)\})$. Hence $\{T(u_1), \dots, T(u_n)\}$ is a basis of \mathbb{W} . Then $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Converse: We now assume that $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces over the same field \mathbb{F} . Then \mathbb{V} and \mathbb{W} are isomorphic if and only if $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Proof: We first assume that \mathbb{V} and \mathbb{W} are isomorphic. Let T be an isomorphism from \mathbb{V} to \mathbb{W} . Let $\{u_1, \dots, u_n\}$ be a basis of \mathbb{V} .

Since T is one-one, then $\{T(u_1), \dots, T(u_n)\}$ is linearly independent.

$\mathbb{W} = R(T) = \text{ls}(\{T(u_1), \dots, T(u_n)\})$. Hence $\{T(u_1), \dots, T(u_n)\}$ is a basis of \mathbb{W} . Then $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Converse: We now assume that $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Let $\{u_1, \dots, u_n\}$ and $\{v_1, \dots, v_n\}$ be two bases of \mathbb{V} and \mathbb{W} , respectively.

We have a linear transformation T such that $T(u_i) = v_i$ for $i = 1, \dots, n$. We can easily check that T is bijective.

• **[Definition:]** Let \mathbb{V} and \mathbb{W} be two vector spaces over the same field \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Then T is called invertible if T is bijective (one-one+onto).

• **[Example:]**

1. Let \mathbb{V} be a vector space over \mathbb{F} . Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be defined by $T(x) = \alpha x$, $\alpha \neq 0$. Then T is invertible.

2. Let $\mathbb{V} = M_{n \times m}(\mathbb{R})$ be the set of $n \times m$ matrices with real entries and let $\mathbb{W} = \mathbb{R}^{mn}$. Define $T : \mathbb{V} \rightarrow \mathbb{W}$ by

$T(A) = (a_{11}, \dots, a_{1m}, a_{21}, \dots, a_{2m}, \dots, a_{n1}, \dots, a_{nm})$. Here $A = (a_{ij}) \in \mathbb{V}$. Then T is invertible.

4. Let $\mathbb{V} = M_{n \times n}(\mathbb{R})$ be the set of $n \times n$ matrices with real entries. Define $T : \mathbb{V} \rightarrow \mathbb{R}$ by $T(A) = \text{trace}(A)$. Then T is not invertible.

- We use $\mathcal{L}(\mathbb{V}, \mathbb{W})$ to denote set of all linear transformation from \mathbb{V} to \mathbb{W} .
- Let $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{R}$. Then $S + T$ and αS defined by $(S + T)(x) = S(x) + T(x)$ (vector addition) and $(\alpha S)x = \alpha S(x)$ (scalar multiplication).
- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be two VS over \mathbb{F} . Then $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is also a vector space with respect to above two operations over \mathbb{F} .

Proof It is trivial.

- [Definition:]

1. A linear transformation T from \mathbb{V} to \mathbb{V} is called a **linear operator**.
2. A linear transformation T from \mathbb{V} to \mathbb{F} is called a **linear functional**.

- [Example]

1. Let $T : \mathbb{M}_{n \times n} \rightarrow \mathbb{R}$ defined as $T(A) = \text{trace}(A)$, $A \in \mathbb{M}_{n \times n}$. T is linear functional.
2. Let $T : C[0, 1] \rightarrow \mathbb{R}$ defined as $T(f) = \int_0^1 f(x) dx$. T is linear functional.
3. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined as $T(x_1, x_2) = (x_1 + x_2, x_1 - x_2)$. T is linear operator.

- **[Definition:]** The space $\mathcal{L}(\mathbb{V}, \mathbb{F})$ is called the **dual space** of \mathbb{V} and it is denoted by \mathbb{V}^* . Elements of \mathbb{V}^* are usually denoted by lower case letters f , g , etc.
- **[Theorem:]** Let \mathbb{V} be a finite dimensional space and $B = \{v_1, \dots, v_n\}$ be an ordered basis of \mathbb{V} .

For each $j \in \{1, \dots, n\}$, let $f_j : \mathbb{V} \rightarrow \mathbb{F}$ be defined by $f_j(x) = \alpha_j$ for $x = \sum_{j=1}^n \alpha_j v_j$.

Then the following are true.

1. f_1, \dots, f_n are in \mathbb{V}^* and they satisfy $f_i(v_j) = \delta_{ij}$ for $i, j \in \{1, \dots, n\}$.
2. $\{f_1, \dots, f_n\}$ is a basis of \mathbb{V}^* .

Proof: We first show that $f_i(v_j) = \delta_{ij}$ for $i, j \in \{1, \dots, n\}$.

$v_j = 0v_1 + \dots + v_j + \dots + 0v_n$. Using the definition of f_i , we have

$$f_i(v_j) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

That is $f_i(v_j) = \delta_{ij}$.

We now show that $f_i \in \mathbb{V}^*$, that is f_i is a linear functional for $i = 1, \dots, n$.

Let $x, y \in \mathbb{V}$. Then $x = a_1v_1 + \dots + a_nv_n$ and $y = b_1v_1 + \dots + b_nv_n$.

Using definition of f_i we have, $f_i(x) = a_i$ and $f_i(y) = b_i$.

Let $\alpha, \beta \in \mathbb{F}$. Then $\alpha x + \beta y = (\alpha a_1 + \beta b_1)v_1 + \cdots + (\alpha a_n + \beta b_n)v_n$.

Using definition of f_i we have

$$f_i(\alpha x + \beta y)$$

$$= \alpha a_i + \beta b_i$$

$$= \alpha f_i(x) + \beta f_i(y).$$

Hence f_i is linear transformation from \mathbb{V} to \mathbb{F} for $i = 1, \dots, n$. We have proved that $f_i \in \mathbb{V}^*$ for $i = 1, \dots, n$.

We now show that $\{f_1, \dots, f_n\}$ is a basis of \mathbb{V}^* . We first show that $\{f_1, \dots, f_n\}$ is linearly independent.

$$c_1 f_1 + c_2 f_2 + \cdots + c_n f_n = 0.$$

$$(c_1 f_1 + c_2 f_2 + \cdots + c_n f_n)v_1 = 0(v_1).$$

$$c_1 f_1(v_1) + \cdots + c_n f_n(v_1) = 0.$$

$$c_1 = 0.$$

Similarly you can show that $c_2 = \cdots = c_n = 0$. Hence $\{f_1, \dots, f_n\}$ is linearly independent.

We now show that $\text{ls}(\{f_1, \dots, f_n\}) = \mathbb{V}^*$.

Let $f \in \mathbb{V}^*$. Let $f(v_i) = c_i$ for $i = 1, \dots, n$ where $c_1, \dots, c_n \in \mathbb{F}$. We have to show that $f = a_1 f_1 + \dots + a_n f_n$ where $a_1, \dots, a_n \in \mathbb{F}$.

Let $x \in \mathbb{V}$. Then $x = b_1 v_1 + \dots + b_n v_n$.

$$f(x) = f(b_1 v_1 + \dots + b_n v_n)$$

$$= b_1 f(v_1) + \dots + b_n f(v_n)$$

$$= c_1 b_1 + \dots + c_n b_n$$

$$= c_1 f_1(x) + \dots + c_n f_n(x)$$

$f(x) = (c_1 f_1 + \dots + c_n f_n)(x)$ for all $x \in \mathbb{V}$. Therefore $f = c_1 f_1 + \dots + c_n f_n$.
Hence $\text{ls}(\{f_1, \dots, f_n\}) = \mathbb{V}^*$.

- **[Definition:]** Let \mathbb{V} be a finite dimensional space and $B = \{v_1, \dots, v_n\}$ be an order basis of \mathbb{V} . A basis $\{f_1, \dots, f_n\}$ of \mathbb{V}^* such that $f_i(v_j) = \delta_{ij}$ for $i, j \in \{1, \dots, n\}$. Then $\{f_1, \dots, f_j\}$ is called **dual basis** of \mathbb{V}^* .
- **[Example:** How to compute dual basis:]

Let $\mathbb{V} = \mathbb{R}^2$. Let $B = \{(1, 0), (0, 1)\}$ be a basis of \mathbb{V} . Find the dual basis of \mathbb{V}^* corresponding B .

Let $\{f_1, f_2\}$ be the dual basis of \mathbb{V}^* corresponding B .

$f_1(x_1, x_2) = \alpha_1 x_1 + \alpha_2 x_2$ and $f_2 = \beta_1 x_1 + \beta_2 x_2$ where $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}$.

$f_1(1, 0) = 1 \implies \alpha_1 = 1$ and $f_1(0, 1) = 0 \implies \alpha_2 = 0$

Similarly $f_2(1, 0) = 0 \implies \beta_1 = 0$ and $f_2(0, 1) = 1 \implies \beta_2 = 1$.

$f_1(x_1, x_2) = x_1$ and $f_2(x_1, x_2) = x_2$

- **[Theorem:]** If V is finite dimensional, then V and V^* are isomorphic.

Proof: We have seen that $\dim(V) = \dim(V^*)$. Then they are isomorphic.

Let \mathbb{V} and \mathbb{W} be two FDVS over \mathbb{F} . Let $B_1 = \{u_1, \dots, u_n\}$ and $B_2 = \{v_1, \dots, v_m\}$ be two ordered bases of \mathbb{V} and \mathbb{W} , respectively.

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

$T(u_j) \in \mathbb{W}$. Then there exist $a_{ij} \in \mathbb{F}$ for $i = 1, \dots, m$ such that

$$T(u_j) = a_{1j}v_1 + a_{2j}v_2 + \cdots + a_{mj}v_m \text{ for } j = 1, \dots, n.$$

Let $x \in \mathbb{V}$. There exist $\alpha_1, \dots, \alpha_n \in \mathbb{F}$ such that $x = \sum_{j=1}^n \alpha_j u_j$. That is

$$[x]_{B_1} = \begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{bmatrix}.$$

$$\begin{aligned}T(x) &= T\left(\sum_{j=1}^n \alpha_j u_j\right) \\&= \sum_{j=1}^n \alpha_j T(u_j) \\&= \sum_{j=1}^n \alpha_j \sum_{i=1}^m a_{ij} v_i \\&= \sum_{i=1}^m \left(\sum_{j=1}^n \alpha_j a_{ij} \right) v_i.\end{aligned}$$

$$\text{Let } [T(x)]_{B_2} = \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_m \end{bmatrix}.$$

$$\text{Then } \beta_i = \sum_{j=1}^n \alpha_j a_{ij} \text{ for } j = 1, \dots, m$$

$$\begin{bmatrix} \beta_1 \\ \vdots \\ \beta_m \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{bmatrix}.$$

$[T(x)]_{B_2} = A[x]_{B_1}$ where $A = [a_{ij}]_{m \times n}$. That is co-ordinate of $T(x)$ with respect to the basis B_2 is $[T(x)]_{B_2}$ which can be calculated using the co-ordinate of x with respect to basis B_1 .

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by $T\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} x + y \\ x - z \end{bmatrix}$. Let

$B_1 = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ and $B_2 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ be bases of \mathbb{R}^3

and \mathbb{R}^2 , respectively.

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} - 2 \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ -1 \end{bmatrix} = 0 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} - 1 \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & -2 & -1 \end{bmatrix}.$$

$$\text{Let } \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \in \mathbb{R}^3. \text{ Then } \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 1 \cdot \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + 0 \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

$$\text{The co-ordinate of } \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \text{ with respect to } B_1 \text{ is } \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

$$[T \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}]_{B_2} = A[\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}]_{B_1} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} - 2 \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

The matrix $A = (a_{ij})$ in the above discussion is called the **matrix representation** of T with respect to the ordered bases B_1 and B_2 of \mathbb{V} and \mathbb{W} , respectively. This matrix is usually denoted by $[T]_{B_1 B_2}$, that is, $[T]_{B_1 B_2} = (a_{ij})$.

Let \mathbb{V} and \mathbb{W} be finite dimensional vector spaces over the same field \mathbb{F} . Let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$. Assume that $B = \{u_1, \dots, u_n\}$ and $B' = \{v_1, \dots, v_m\}$ are ordered basis of \mathbb{V} and \mathbb{W} respectively.

1. We have seen that for each linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$, we have a matrix $A \in \mathbb{M}_{m \times n}(\mathbb{F})$ such that $[T]_{BB'} = A$.
2. Let $A \in \mathbb{M}_{m \times n}(\mathbb{F})$. Then there exists a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ such that $A = [T]_{BB'}$ and such linear transformation is $T(u_j) = \sum_{i=1}^m a_{ij}v_i$ for $j = 1 \dots, n$.
3. Let $T, S : \mathbb{V} \rightarrow \mathbb{W}$ be two linear transformation. Let B_1 and B_2 be two bases of \mathbb{V} and \mathbb{W} , respectively. Then $[T + S]_{B_1B_2} = [T]_{B_1B_2} + [S]_{B_1B_2}$ and $[\alpha T]_{B_1B_2} = \alpha[T]_{B_1B_2}$.

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be finite dimensional vector spaces over the same field \mathbb{F} . Let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$. Then $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is isomorphic to $\mathbb{M}_{m \times n}(\mathbb{F})$.

Proof: Let $B = \{u_1, \dots, u_n\}$ and $B' = \{v_1, \dots, v_m\}$ be bases of \mathbb{V} and \mathbb{W} , respectively.

Define $\zeta : \mathcal{L}(\mathbb{V}, \mathbb{W}) \rightarrow \mathbb{M}_{m \times n}(\mathbb{F})$ such that $\zeta(T) = [T]_{BB'}$.

Using previous remark it is cleared that ζ is linear from $\mathcal{L}(\mathbb{V}, \mathbb{W})$ to $\mathbb{M}_{m \times n}(\mathbb{F})$. We now show that ζ is bijective.

Let $T \in \text{Ker}(\zeta)$. Then $\zeta(T) = 0_{m \times n}$.

This implies that $[T]_{BB'} = 0_{m \times n}$.

This implies $[T(x)]_{B'} = 0_{n \times 1}$, co-ordinate of $T(x)$ for each $x \in \mathbb{V}$ with respect to B' is zero. Hence $T(x) = 0_{\mathbb{W}}$ for each $x \in \mathbb{V}$. Then $T = 0$. Therefore $\text{Ker}(\zeta) = \{0\}$.

We now show that ζ is onto. Let $A \in \mathbb{M}_{m \times n}$. Define $T(u_j) = \sum_{i=1}^m a_{ij} v_i$ for $j = 1 \dots, n$. It is clear that $[T]_{BB'} = A$. Hence ζ is onto.

Therefore $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is isomorphic to $\mathbb{M}_{m \times n}(\mathbb{F})$.

- **[Theorem:]** Let \mathbb{V} and \mathbb{W} be finite dimensional vector spaces over the same field \mathbb{F} . Let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$. Then dimension of $\mathcal{L}(\mathbb{V}, \mathbb{W}) = mn$.

- **[Theorem:]** Let \mathbb{V} be a finite dimensional vector space over the same field \mathbb{F} . Let S and T be two linear transformations from \mathbb{V} and to \mathbb{V} . Let B be an ordered basis of \mathbb{V} . Then $[S \circ T]_{BB} = [S]_{BB}[T]_{B,B}$.

Proof: Let $B = \{v_1, \dots, v_n\}$. Let $[T]_{BB} = A$ and $[S]_{BB} = C$.

Then $T(v_i) = a_{1i}v_1 + a_{2i}v_2 + \dots + a_{ni}v_n$ for $i = 1, \dots, n$.

$S(v_i) = b_{1i}v_1 + b_{2i}v_2 + \dots + b_{ni}v_n$ for $i = 1, \dots, n$.

$$(S \circ T)(v_1) = S(T(v_1))$$

$$= S(a_{11}v_1 + a_{21}v_2 + \dots + a_{n1}v_n)$$

$$= a_{11}S(v_1) + a_{21}S(v_2) + \dots + a_{n1}S(v_n)$$

$$= a_{11}(b_{11}v_1 + b_{21}v_2 + \dots + b_{n1}v_n) + a_{21}(b_{12}v_1 + b_{22}v_2 + \dots + b_{n2}v_n) + \dots + a_{n1}(b_{1n}v_1 + b_{2n}v_2 + \dots + b_{nn}v_n)$$

$$= (a_{11}b_{11} + a_{21}b_{12} + \cdots a_{n1}b_{1n})v_1 + \cdots + (a_{11}b_{n1} + a_{21}b_{n2} + \cdots a_{n1}b_{nn})v_n$$

$$= \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{bmatrix} \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{bmatrix}$$

$$[(S \circ T)(u_i)] = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{bmatrix} \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{ni} \end{bmatrix}$$

$$[S \circ T]_{BB} = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$$

- **[Remark:]** Let $\dim V = n$ and $\dim W = m$. Let $T : V \rightarrow W$ and $S : W \rightarrow V$ be two linear transformation. Let B and B' be bases of V and W , respectively. Then $[S \circ T]_{BB'} = [S]_{BB'}[T]_{BB'}$.

- **[Theorem:]** Let V be a finite dimensional vector space over the same field \mathbb{F} . Let T be an invertible linear transformation from V and to V . Let B be an ordered basis of V . Then $[T^{-1}]_{BB} = [T]_{BB}^{-1}$.

- **[Theorem]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces and let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$.

- **[Theorem]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces and let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$.

Let B and C be two ordered bases of \mathbb{V} and let B' and C' be two bases of \mathbb{W} .

- **[Theorem]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces and let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$.

Let B and C be two ordered bases of \mathbb{V} and let B' and C' be two bases of \mathbb{W} .

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

- **[Theorem]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces and let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$.

Let B and C be two ordered bases of \mathbb{V} and let B' and C' be two bases of \mathbb{W} .

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

Question: Is there any relation between $[T]_{BB'}$ and $[T]_{CC'}$?

- **[Theorem]** Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces and let $\dim(\mathbb{V}) = n$ and $\dim(\mathbb{W}) = m$.

Let B and C be two ordered bases of \mathbb{V} and let B' and C' be two bases of \mathbb{W} .

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation.

Question: Is there any relation between $[T]_{BB'}$ and $[T]_{CC'}$?

Then there exist two non-singular matrix $P \in \mathbb{M}_m(\mathbb{F})$ and $Q \in \mathbb{M}_n(\mathbb{F})$ such that $[T]_{BB'} = P^{-1}[T]_{CC'}Q$.

- **[Theorem]** Let V be an n -dimensional vector space over \mathbb{F} .

- **[Theorem]** Let V be an n -dimensional vector space over \mathbb{F} .

Let T be a linear transformation from V and to V .

- **[Theorem]** Let \mathbb{V} be an n -dimensional vector space over \mathbb{F} .

Let T be a linear transformation from \mathbb{V} and to \mathbb{V} .

Let B and B' be two ordered basis of \mathbb{V} .

- **[Theorem]** Let \mathbb{V} be an n -dimensional vector space over \mathbb{F} .

Let T be a linear transformation from \mathbb{V} and to \mathbb{V} .

Let B and B' be two ordered basis of \mathbb{V} .

Then there exists a non-singular matrix P such that $[T]_{B'B'} = P^{-1}[T]_{BB}P$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Therefore $S \circ T \circ S^{-1}(v_j) = S \circ T(u_j) = S(T(u_j)) = S(\sum_{i=1}^n a_{ji} u_i) = \sum_{i=1}^n a_{ji} v_i$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Therefore $S \circ T \circ S^{-1}(v_j) = S \circ T(u_j) = S(T(u_j)) = S(\sum_{i=1}^n a_{ji} u_i) = \sum_{i=1}^n a_{ji} v_i$.

$[STS^{-1}]_{B'B'} = (a_{ij}) = [T]_{BB}$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Therefore $S \circ T \circ S^{-1}(v_j) = S \circ T(u_j) = S(T(u_j)) = S(\sum_{i=1}^n a_{ji} u_i) = \sum_{i=1}^n a_{ji} v_i$.

$[STS^{-1}]_{B'B'} = (a_{ij}) = [T]_{BB}$.

$\implies [S]_{B'B'}[T]_{B'B'}[S^{-1}]_{B'B'} = [T]_{BB}$.

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Therefore $S \circ T \circ S^{-1}(v_j) = S \circ T(u_j) = S(T(u_j)) = S(\sum_{i=1}^n a_{ji} u_i) = \sum_{i=1}^n a_{ji} v_i$.

$[STS^{-1}]_{B'B'} = (a_{ij}) = [T]_{BB}$.

$$\implies [S]_{B'B'} [T]_{B'B'} [S^{-1}]_{B'B'} = [T]_{BB}.$$

$$\implies [S]_{B'B'} [T]_{B'B'} [S]_{B'B'}^{-1} = [T]_{BB}.$$

Let $B = \{u_1, \dots, u_n\}$ and let $B' = \{v_1, \dots, v_n\}$.

Let $S : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation such that $S(u_i) = v_i$ for $i = 1, \dots, n$.

It is clear that S is bijective. Let $[T]_{BB} = A = (a_{ij})$.

Then $T(u_j) = \sum_{i=1}^n a_{ji} u_i$.

Therefore $S \circ T \circ S^{-1}(v_j) = S \circ T(u_j) = S(T(u_j)) = S(\sum_{i=1}^n a_{ji} u_i) = \sum_{i=1}^n a_{ji} v_i$.

$[STS^{-1}]_{B'B'} = (a_{ij}) = [T]_{BB}$.

$$\implies [S]_{B'B'} [T]_{B'B'} [S^{-1}]_{B'B'} = [T]_{BB}.$$

$$\implies [S]_{B'B'} [T]_{B'B'} [S]_{B'B'}^{-1} = [T]_{BB}.$$

$$\implies [T]_{B'B'} = [S]_{B'B'}^{-1} [T]_{BB} [S]_{B'B'} = P^{-1} [T]_{BB} P \text{ where } P = [S]_{B'B'}.$$

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x + y) = f(x) + f(y)$.

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x + y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x + y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x+y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

Can we construct a basis B' from B which contains 1 and $\sqrt{2}$?

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x+y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

Can we construct a basis B' from B which contains 1 and $\sqrt{2}$?

Yes, we can construct a basis B' from B which contains 1 and $\sqrt{2}$. This basis contains exactly one rational number which is 1. If B' contains two rationals 1 and q , then they are linearly dependent which is not possible.

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x+y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

Can we construct a basis B' from B which contains 1 and $\sqrt{2}$?

Yes, we can construct a basis B' from B which contains 1 and $\sqrt{2}$. This basis contains exactly one rational number which is 1. If B' contains two rationals 1 and q , then they are linearly dependent which is not possible.

We can easily construct a linear transformation $f: \mathbb{R}(\mathbb{Q}) \rightarrow \mathbb{R}(\mathbb{Q})$ such that

$$f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } x \in B' \text{ and irrational} \end{cases}$$

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x+y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

Can we construct a basis B' from B which contains 1 and $\sqrt{2}$?

Yes, we can construct a basis B' from B which contains 1 and $\sqrt{2}$. This basis contains exactly one rational number which is 1. If B' contains two rationals 1 and q , then they are linearly dependent which is not possible.

We can easily construct a linear transformation $f: \mathbb{R}(\mathbb{Q}) \rightarrow \mathbb{R}(\mathbb{Q})$ such that

$$f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } x \in B' \text{ and irrational} \end{cases}$$

Since f is a map from \mathbb{R} to \mathbb{R} and it satisfies $f(x+y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$ as f is a linear transformation from $\mathbb{R}(\mathbb{Q})$ to $\mathbb{R}(\mathbb{Q})$.

There is a non-linear map from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ such that $f(x+y) = f(x) + f(y)$.

We know that $\mathbb{R}(\mathbb{Q})$ is an infinite dimensional space.

Let B be a basis of $\mathbb{R}(\mathbb{Q})$.

Can we construct a basis B' from B which contains 1 and $\sqrt{2}$?

Yes, we can construct a basis B' from B which contains 1 and $\sqrt{2}$. This basis contains exactly one rational number which is 1. If B' contains two rationals 1 and q , then they are linearly dependent which is not possible.

We can easily construct a linear transformation $f: \mathbb{R}(\mathbb{Q}) \rightarrow \mathbb{R}(\mathbb{Q})$ such that

$$f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } x \in B' \text{ and irrational} \end{cases}$$

Since f is a map from \mathbb{R} to \mathbb{R} and it satisfies $f(x+y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$ as f is a linear transformation from $\mathbb{R}(\mathbb{Q})$ to $\mathbb{R}(\mathbb{Q})$.

$$f(\sqrt{2}) = \sqrt{2}f(1).$$

$$f(\sqrt{2}) = \sqrt{2}f(1).$$

$0 = \sqrt{2}$ a contradiction. Hence f is non-linear from $\mathbb{R}(\mathbb{R})$ to $\mathbb{R}(\mathbb{R})$ satisfying $f(x+y) = f(x) + f(y)$.