

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

- The inner product can be defined on the vector space over the field either \mathbb{R} or \mathbb{C} . We use the following notation of the field \mathbb{K} where \mathbb{K} is either \mathbb{R} or \mathbb{C} .

Definition: Let \mathbb{V} be a vector space over the field \mathbb{K} .

An inner product is a mapping $\langle ., . \rangle : \mathbb{V} \times \mathbb{V} \rightarrow \mathbb{K}$ which satisfies the following conditions.

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

1. $\langle x, x \rangle \geq 0$ for all $x \in \mathbb{V}$ (**positivity**)

and $\langle x, x \rangle = 0$ iff $x = 0$ (**definiteness**).

2. $\langle x, y \rangle = \overline{\langle y, x \rangle}$ for all $x, y \in \mathbb{V}$ (**conjugate symmetry**).

3. $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ for all $x, y \in \mathbb{V}$ and for all $\alpha \in \mathbb{K}$ (**homogeneity**).

4. $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$ for all $x, y, z \in \mathbb{V}$ (**additivity**).

Examples:

- Dot product on \mathbb{R}^n is an inner product.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_i^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Examples:

- Dot product on \mathbb{R}^n is an inner product.
- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$ for all $x, y \in \mathbb{V}$.

Ans: This is not an inner product. Take $x = (i, i, \dots, i)$. Then $\langle x, x \rangle = \sum_{i=1}^n i \cdot i = -n$ which is negative. Therefore fails to satisfy property 1.

- Let $\mathbb{V} = \mathbb{C}^n(\mathbb{C})$. Let $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$ for all $x, y \in \mathbb{V}$.

Ans: This an inner product.

- $\mathbb{V} = \mathbb{M}_n(\mathbb{R})(\mathbb{R})$. Let $\langle A, B \rangle = \text{trace}(AB^t)$ for all $A, B \in \mathbb{M}_n(\mathbb{R})$.

Ans: This is an inner product.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

- $\mathbb{V} = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

- $V = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

- $\mathbb{V} = \mathbb{C}[a, b]$ over the field \mathbb{R} . Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}[a, b]$.

1. Let $f \in \mathbb{C}[a, b]$. Take $\langle f, f \rangle = \int_a^b f(x)f(x)dx = \int_a^b f^2(x)dx \geq 0$.

$$\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0 \text{ (why?)}$$

2. Let $f, g \in \mathbb{C}[a, b]$.

$$\text{Take } \langle f, g \rangle = \int_a^b f(x)g(x)dx = \int_a^b g(x)f(x)dx = \langle g, f \rangle.$$

3. Let $f, g \in \mathbb{C}[a, b]$ and $\alpha \in \mathbb{R}$.

$$\text{Then } \langle \alpha f, g \rangle = \int_a^b (\alpha f(x))g(x)dx = \alpha \int_a^b f(x)g(x)dx = \alpha \langle f, g \rangle.$$

4. Let $f, g, h \in \mathbb{C}[a, b]$.

4. Let $f, g, h \in \mathbb{C}[a, b]$.

$$\begin{aligned}\text{Then } \langle f + g, h \rangle &= \int_a^b (f(x) + g(x)) h(x) dx \\ &= \int_a^b (f(x)h(x) + g(x)h(x)) dx \\ &= \langle f, h \rangle + \langle g, h \rangle.\end{aligned}$$

- Let $f : [a, b]$ be integrable and $f(x) \geq 0$. Let f be continuous at $x = c$ and $f(c) > 0$ (resp. $f(c) < 0$). Then $\int_a^b f(x)dx > 0$ (resp. $\int_a^b f(x)dx < 0$).

Sol: Case I. Let $a < c < b$.

f is continuous at $x = c$. Take $\epsilon = \frac{f(c)}{2}$ then there exists $\delta > 0$ such that

for each $x \in (c - \delta, c + \delta) \implies |f(x) - f(c)| < \frac{f(c)}{2}$.

Then $-\frac{f(c)}{2} < f(x) - f(c) < \frac{f(c)}{2}$ for all $x \in (c - \delta, c + \delta)$.

Therefore $f(x) > \frac{f(c)}{2} > 0$ for all $x \in (c - \delta, c + \delta)$.

$$\int_a^b f^2(x)dx = \int_a^{c-\delta} f(x)dx + \int_{c-\delta}^{c+\delta} f(x)dx + \int_{c+\delta}^b f(x)dx > 0$$

as $\int_{c-\delta}^{c+\delta} f^2(x)dx > 0$.

Case II. $c = a$

When $c = a$, we have a neighborhood $(a, a + \delta)$ where f is positive.

$$\int_a^b f^2(x) dx = \int_a^{a+\delta} f(x) dx + \int_b^{a+\delta} f(x) dx > 0$$

Case-III. $c = b$. Same as Case II.

To show $\langle f, f \rangle = \int_a^b f(x)f(x)dx = 0 \implies f \equiv 0$.

Here f^2 is nonnegative function and continuous. If f is not identically zero, then there exists $c \in [a, b]$ such that $f(c) \neq 0$. Therefore $f^2(c) > 0$. Using above result we have $\int_a^b f(x)f(x)dx > 0$, a contradiction. Hence $f \equiv 0$.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

- $V = \mathbb{C}(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in \mathbb{C}(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in \mathbb{C}(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

- $V = C(\mathbb{R})$ and $\mathbb{K} = \mathbb{R}$. Let $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ for all $f, g \in C(\mathbb{R})$ where a and b are fixed real numbers.

Ans: This is not an inner product space.

Take $f \in C(\mathbb{R})$, defined by

$$f(x) = \begin{cases} a - x & \text{if } x \leq a \\ 0 & \text{if } x \in [a, b] \\ x - b & \text{if } x \geq b \end{cases}$$

f is continuous on \mathbb{R} .

$$\langle f, f \rangle = \int_a^b f^2(x)dx = 0 \text{ but } f \not\equiv 0.$$

Hence $\langle f, g \rangle = \int_a^b f(x)g(x)dx$ is not an inner product on $C(\mathbb{R})$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let V be a vector space over \mathbb{K} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, k\}$ be a basis of \mathbb{V} .

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

Theorem: There exists an inner product on every non-trivial vector space over \mathbb{K} .

Proof: Let \mathbb{V} be a vector space over \mathbb{K} .

There are two cases.

Case I. \mathbb{V} is finite dimensional. Let $B = \{u_1, \dots, u_k\}$ be a basis of \mathbb{V} .

Let $x, y \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$ for some $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Define

$$\langle x, y \rangle = \sum_{i=1}^k a_i \bar{b}_i .$$

We now show that \langle, \rangle is an IP on \mathbb{V} .

-
1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

1. Let $x \in \mathbb{V}$. The $x = \sum_{i=1}^k a_i u_i$ for some $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, x \rangle = \sum_{i=1}^k a_i \bar{a}_i = \sum_{i=1}^k |a_i|^2 \geq 0.$$

if $\langle x, x \rangle = 0$, then $\sum_{i=1}^k |a_i|^2 = 0 \implies a_i = 0$ for $i = 1, \dots, k$.

Therefore $x = \sum_{i=1}^k a_i u_i = 0$.

2. Let $x, y \in \mathbb{V}$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

2. Let $x, y \in \mathbb{V}$.

Then $x = \sum_{i=1}^k a_i u_i$ and $y = \sum_{i=1}^k b_i u_i$, $a_i, b_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$$\langle x, y \rangle = \sum_{i=1}^k a_i \overline{b_i} = \overline{\sum_{i=1}^k \overline{a_i} b_i} = \overline{\langle y, x \rangle}.$$

3. This part is same as part 2.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

3. This part is same as part 2.

4. Let $x, y, z \in \mathbb{V}$. Then $x = \sum_{i=1}^k a_i u_i$, $y = \sum_{i=1}^k b_i u_i$ and $z = \sum_{i=1}^k c_i u_i$ where $a_i, b_i, c_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Then $x + y = \sum_{i=1}^k (a_i + b_i) u_i$. Therefore,

$$\begin{aligned}\langle x + y, z \rangle &= \sum_{i=1}^k (a_i + b_i) \overline{c_i} \\ &= \sum_{i=1}^k (a_i \overline{c_i} + b_i \overline{c_i}) \\ &= \langle x, z \rangle + \langle y, z \rangle.\end{aligned}$$

Case II. \mathbb{V} is infinite dimensional.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Case II. \mathbb{V} is infinite dimensional.

Let $B = \{u_\alpha : \alpha \in I\}$ be a basis of \mathbb{V} where I is an index set.

Take $f : B \times B \rightarrow \mathbb{K}$ define by

$$f(u_\alpha, u_\beta) = \begin{cases} 1 & \text{if } \alpha = \beta \\ 0 & \text{if } \alpha \neq \beta \end{cases}$$

Let $x, y \in \mathbb{V}$. Then there exist two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ and $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Define $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{j=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_i} f(u_{\alpha_i}, u_{\alpha_i}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_i} f(u_{\alpha_i}, u_{\alpha_i}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k}$ where $a_i \in \mathbb{K}$ for $i = 1, \dots, k$.

$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m}$ where $b_j \in \mathbb{K}$ for $j = 1, \dots, m$.

Check $\langle x, y \rangle = \sum_{j=1}^m \sum_{i=1}^k a_i \overline{b_j} f(u_{\alpha_i}, u_{\beta_j})$ is an inner product on \mathbb{V} .

$$\begin{aligned} 1. \quad \langle x, x \rangle &= \sum_{i=1}^k \sum_{i=1}^k a_i \overline{a_j} f(u_{\alpha_i}, u_{\alpha_j}). \\ &= \sum_{i=1}^k |a_i|^2 \text{ (using the definition of } f) \end{aligned}$$

It is easy to show that $\langle x, x \rangle = 0$ if and only if $x = 0$.

2. It is easy.

3. It is easy

4. Let $x, y, z \in \mathbb{V}$. Then there exists two finite subsets $\{u_{\alpha_1}, \dots, u_{\alpha_k}\}$ $\{u_{\beta_1}, \dots, u_{\beta_m}\}$ and $\{u_{\gamma_1}, \dots, u_{\gamma_n}\}$ of B such that

$$x = a_1 u_{\alpha_1} + \dots + a_k u_{\alpha_k} \text{ where } a_i \in \mathbb{K} \text{ for } i = 1, \dots, k.$$

$$y = b_1 u_{\beta_1} + \dots + b_m u_{\beta_m} \text{ where } b_j \in \mathbb{K} \text{ for } j = 1, \dots, m.$$

$$z = c_1 u_{\gamma_1} + \dots + c_n u_{\gamma_n} \text{ where } c_l \in \mathbb{K} \text{ for } l = 1, \dots, n.$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle.$$

$$x + y = \sum_{i=1}^k a_i u_{\alpha_i} + \sum_{j=1}^m b_j u_{\beta_j}$$

$$\langle x + y, z \rangle = \sum_{i=1}^k \sum_{l=1}^n a_i \overline{c_l} f(u_{\alpha_i}, u_{\gamma_l}) + \sum_{j=1}^m \sum_{l=1}^n b_j \overline{c_l} f(u_{\beta_j}, u_{\gamma_l}).$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle.$$

- A vectors space together with an inner product is called an inner product space.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
 3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.

-
1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
 2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
 3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.
 4. $\langle u, \lambda v \rangle = \overline{\lambda} \langle u, v \rangle$ for all $u, v \in \mathbb{V}$ and $\lambda \in \mathbb{K}$.

1. $\langle 0, u \rangle = 0$ for every $u \in \mathbb{V}$.
2. $\langle u, 0 \rangle = 0$ for every $u \in \mathbb{V}$.
3. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$ for every $u, v, w \in \mathbb{V}$.
4. $\langle u, \lambda v \rangle = \overline{\lambda} \langle u, v \rangle$ for all $u, v \in \mathbb{V}$ and $\lambda \in \mathbb{K}$.

Proof: 1. $\langle 0, u \rangle = \langle u - u, u \rangle$

$$= \langle u, u \rangle + \langle -u, u \rangle$$

$$= \langle u, u \rangle - \langle u, u \rangle$$

$$= 0.$$

$$2. \langle 0, u \rangle = \overline{\langle 0, u \rangle}$$

$$= \overline{\langle u - u, u \rangle}$$

$$= \overline{\langle u, u \rangle + \langle -u, u \rangle}$$

$$= \overline{\langle u, u \rangle - \langle u, u \rangle}$$

$$= \overline{0}$$

3. Suppose $u, v, w \in \mathbb{V}$. Then

$$\begin{aligned}\langle u, v + w \rangle &= \overline{\langle v + w, u \rangle} \\ &= \overline{\langle v, u \rangle + \langle w, u \rangle} \\ &= \overline{\langle v, u \rangle} + \overline{\langle w, u \rangle} \\ &= \langle u, v \rangle + \langle u, w \rangle.\end{aligned}$$

4. Suppose $\lambda \in \mathbb{K}$ and $u, v \in \mathbb{V}$. Then

$$\langle u, \lambda v \rangle = \overline{\langle \lambda v, u \rangle}$$

$$= \overline{\lambda \langle v, u \rangle}$$

$$= \overline{\lambda} \overline{\langle v, u \rangle}$$

$$= \overline{\lambda} \langle u, v \rangle.$$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

$\langle e_1, e_2 \rangle_1 = -1$. These two vectors are not orthogonal with respect to $\langle \cdot, \cdot \rangle_1$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u, v \in \mathbb{V}$. Then u and v is called **orthogonal** if $\langle u, v \rangle = 0$.

Note: Orthogonality of two vectors depends on the inner product.

Consider $\mathbb{V} = \mathbb{R}^2$. Take $\langle x, y \rangle = x_1y_1 + x_2y_2$ where $x = (x_1, x_2)$, $y = (y_1, y_2)$. This an inner product on \mathbb{R}^2 .

$e_1 = (1, 0)$ and $e_2 = (0, 1)$ are orthogonal with respect to the above inner product.

Consider $\langle x, y \rangle_1 = x_1y_1 - x_2y_1 - x_1y_2 + 4x_2y_2$. This is an inner product on \mathbb{R}^2 (check!).

$\langle e_1, e_2 \rangle_1 = -1$. These two vectors are not orthogonal with respect to $\langle \cdot, \cdot \rangle_1$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. A subset S of \mathbb{V} is said to be **orthogonal** if $\langle u, v \rangle = 0$ for all $u, v \in S$ and $u \neq v$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$. Take inner product with α_i both side.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S = \{\alpha_1, \dots, \alpha_k\}$ be an orthogonal subset of \mathbb{V} and let $\alpha_i \neq 0$ for $i = 1, \dots, k$. Then S is linearly independent.

Proof: To show $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

Take $c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k = 0$. Take inner product with α_i both side.

$$\langle c_1\alpha_1 + c_2\alpha_2 + \dots + c_k\alpha_k, \alpha_i \rangle = \langle 0, \alpha_i \rangle$$

$$\sum_{j=1}^k c_j \langle \alpha_j, \alpha_i \rangle = 0.$$

$$c_i \langle \alpha_i, \alpha_i \rangle = 0$$

$c_i = 0$. This is true for $i = 1, \dots, k$.

Hence $S = \{\alpha_1, \dots, \alpha_k\}$ is LI.

-
- The converse of the above theorem is not true. The following is an example.

- The converse of the above theorem is not true. The following is an example.

Example: Let $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$ be an inner product space where $\langle x, y \rangle = \sum_{i=1}^2 x_i y_i$.

- The converse of the above theorem is not true. The following is an example.

Example: Let $(\mathbb{R}^2, \langle \cdot, \cdot \rangle)$ be an inner product space where $\langle x, y \rangle = \sum_{i=1}^2 x_i y_i$.

Take $u = (1, 0)$ and $v = (1, 1)$. These two vectors are linearly independent but not orthogonal.

- **[Definition:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space.

- **[Definition:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq \mathbb{V}$.

- **[Definition:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq \mathbb{V}$. Then $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is called **orthonormal** if $\langle \alpha_i, \alpha_j \rangle = 0$ for $i \neq j$ and $\langle \alpha_i, \alpha_i \rangle = 1$.

- **[Definition:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq \mathbb{V}$. Then $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is called **orthonormal** if $\langle \alpha_i, \alpha_j \rangle = 0$ for $i \neq j$ and $\langle \alpha_i, \alpha_i \rangle = 1$.

The next immediate question is that can we construct an orthogonal set from a finite linearly independent set?

- **[Definition:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subseteq \mathbb{V}$. Then $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is called **orthonormal** if $\langle \alpha_i, \alpha_j \rangle = 0$ for $i \neq j$ and $\langle \alpha_i, \alpha_i \rangle = 1$.

The next immediate question is that can we construct an orthogonal set from a finite linearly independent set?

The answer is yes. Gram Schimdt supplied a process to construct an orthogonal set from a linearly independent finite set.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $u_1, u_2, \dots, u_k \in \mathbb{V}$ be LI.

How to construct orthogonal vectors v_1, \dots, v_k using u_1, u_2, \dots, u_k ?

Step 1. Take $v_1 = u_1$.

Step 2. We now construct v_2 using v_1 and u_2 . Take $v_2 = u_2 + cv_1$ where $c \in \mathbb{K}$. We have to calculate the value of c such that v_2 is orthogonal to v_1 . That means

$$\langle v_2, v_1 \rangle = \langle u_2, v_1 \rangle + c \langle v_1, v_1 \rangle$$

$$0 = \langle u_2, v_1 \rangle + c \langle v_1, v_1 \rangle.$$

$$c = -\frac{\langle u_2, v_1 \rangle}{\langle v_1, v_1 \rangle}.$$

Therefore $v_2 = u_2 - \frac{\langle u_2, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1$ is a vector which is perpendicular to v_1 . You can easily check that $v_2 \neq 0$, otherwise u_2 is scalar multiple of u_1 which is not possible.

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Take $v_3 = u_3 + c_1 v_2 + c_2 v_1$ is an element in \mathbb{V} where $c_1, c_2 \in \mathbb{K}$. We have to calculate the values of c_1, c_2 such that v_3 is orthogonal to v_1 and v_2 . That means

$$\langle v_3, v_1 \rangle = \langle u_3, v_1 \rangle + c_1 \langle v_2, v_1 \rangle + c_2 \langle v_1, v_1 \rangle$$

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Take $v_3 = u_3 + c_1 v_2 + c_2 v_1$ is an element in \mathbb{V} where $c_1, c_2 \in \mathbb{K}$. We have to calculate the values of c_1, c_2 such that v_3 is orthogonal to v_1 and v_2 . That means

$$\langle v_3, v_1 \rangle = \langle u_3, v_1 \rangle + c_1 \langle v_2, v_1 \rangle + c_2 \langle v_1, v_1 \rangle$$

$$0 = \langle u_3, v_1 \rangle + c_1 \times 0 + c_2 \langle v_1, v_1 \rangle.$$

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Take $v_3 = u_3 + c_1 v_2 + c_2 v_1$ is an element in \mathbb{V} where $c_1, c_2 \in \mathbb{K}$. We have to calculate the values of c_1, c_2 such that v_3 is orthogonal to v_1 and v_2 . That means

$$\langle v_3, v_1 \rangle = \langle u_3, v_1 \rangle + c_1 \langle v_2, v_1 \rangle + c_2 \langle v_1, v_1 \rangle$$

$$0 = \langle u_3, v_1 \rangle + c_1 \times 0 + c_2 \langle v_1, v_1 \rangle.$$

$$c_2 = -\frac{\langle u_3, v_1 \rangle}{\langle v_1, v_1 \rangle}.$$

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Take $v_3 = u_3 + c_1 v_2 + c_2 v_1$ is an element in \mathbb{V} where $c_1, c_2 \in \mathbb{K}$. We have to calculate the values of c_1, c_2 such that v_3 is orthogonal to v_1 and v_2 . That means

$$\langle v_3, v_1 \rangle = \langle u_3, v_1 \rangle + c_1 \langle v_2, v_1 \rangle + c_2 \langle v_1, v_1 \rangle$$

$$0 = \langle u_3, v_1 \rangle + c_1 \times 0 + c_2 \langle v_1, v_1 \rangle.$$

$$c_2 = -\frac{\langle u_3, v_1 \rangle}{\langle v_1, v_1 \rangle}.$$

$$\text{Similarly, we have } c_1 = -\frac{\langle u_3, v_2 \rangle}{\langle v_2, v_2 \rangle}.$$

Step 3. We now construct v_3 using v_1, v_2, u_3 .

Take $v_3 = u_3 + c_1 v_2 + c_2 v_1$ is an element in \mathbb{V} where $c_1, c_2 \in \mathbb{K}$. We have to calculate the values of c_1, c_2 such that v_3 is orthogonal to v_1 and v_2 . That means

$$\langle v_3, v_1 \rangle = \langle u_3, v_1 \rangle + c_1 \langle v_2, v_1 \rangle + c_2 \langle v_1, v_1 \rangle$$

$$0 = \langle u_3, v_1 \rangle + c_1 \times 0 + c_2 \langle v_1, v_1 \rangle.$$

$$c_2 = -\frac{\langle u_3, v_1 \rangle}{\langle v_1, v_1 \rangle}.$$

$$\text{Similarly, we have } c_1 = -\frac{\langle u_3, v_2 \rangle}{\langle v_2, v_2 \rangle}.$$

Therefore $v_3 = u_3 - \frac{\langle u_3, v_2 \rangle}{\langle v_2, v_2 \rangle} v_2 - \frac{\langle u_3, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1$ is a vector which is perpendicular to v_1 and v_2 . You can easily check that $v_3 \neq 0$, otherwise u_3 is a linear combination of u_1 and u_2 .

Step 4. Similar way you can calculate v_4 using u_4, v_1, v_2 and v_3 .

Step k. $v_k = u_k - \frac{\langle u_k, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \cdots - \frac{\langle u_k, v_{k-1} \rangle}{\langle v_{k-1}, v_{k-1} \rangle} v_{k-1}$. You can easily check that $v_k \neq 0$.

It is very easy to check that v_1, \dots, v_k are orthogonal.

Step 4. Similar way you can calculate v_4 using u_4, v_1, v_2 and v_3 .

Step k. $v_k = u_k - \frac{\langle u_k, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \cdots - \frac{\langle u_k, v_{k-1} \rangle}{\langle v_{k-1}, v_{k-1} \rangle} v_{k-1}$. You can easily check that $v_k \neq 0$.

It is very easy to check that v_1, \dots, v_k are orthogonal.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space.

Step 4. Similar way you can calculate v_4 using u_4, v_1, v_2 and v_3 .

Step k. $v_k = u_k - \frac{\langle u_k, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \dots - \frac{\langle u_k, v_{k-1} \rangle}{\langle v_{k-1}, v_{k-1} \rangle} v_{k-1}$. You can easily check that $v_k \neq 0$.

It is very easy to check that v_1, \dots, v_k are orthogonal.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{u_1, u_2, \dots, u_n\}$ be a linearly independent set. Assume that the set $\{u_1, \dots, u_k\}$ is orthogonal. Then

$v_1 = u_1, v_2 = u_2, \dots, v_k = u_k$. To calculate v_{k+1} apply above technique on v_1, \dots, v_k and u_{k+1} .

- **Theorem:**[Gram Schmidt Orthogonalization] Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space.

Step 4. Similar way you can calculate v_4 using u_4, v_1, v_2 and v_3 .

Step k. $v_k = u_k - \frac{\langle u_k, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \dots - \frac{\langle u_k, v_{k-1} \rangle}{\langle v_{k-1}, v_{k-1} \rangle} v_{k-1}$. You can easily check that $v_k \neq 0$.

It is very easy to check that v_1, \dots, v_k are orthogonal.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{u_1, u_2, \dots, u_n\}$ be a linearly independent set. Assume that the set $\{u_1, \dots, u_k\}$ is orthogonal. Then

$v_1 = u_1, v_2 = u_2, \dots, v_k = u_k$. To calculate v_{k+1} apply above technique on v_1, \dots, v_k and u_{k+1} .

- **Theorem:**[Gram Schmidt Orthogonalization] Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set.

Step 4. Similar way you can calculate v_4 using u_4, v_1, v_2 and v_3 .

Step k. $v_k = u_k - \frac{\langle u_k, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \dots - \frac{\langle u_k, v_{k-1} \rangle}{\langle v_{k-1}, v_{k-1} \rangle} v_{k-1}$. You can easily check that $v_k \neq 0$.

It is very easy to check that v_1, \dots, v_k are orthogonal.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{u_1, u_2, \dots, u_n\}$ be a linearly independent set. Assume that the set $\{u_1, \dots, u_k\}$ is orthogonal. Then

$v_1 = u_1, v_2 = u_2, \dots, v_k = u_k$. To calculate v_{k+1} apply above technique on v_1, \dots, v_k and u_{k+1} .

- **Theorem:**[Gram Schmidt Orthogonalization] Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an orthogonal set $\{\beta_1, \beta_2, \dots, \beta_n\}$ such that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Proof: Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set, $\alpha_i \neq 0$ for $i = 1, \dots, n$.

Consider.

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$$\vdots$$

$$\beta_n = \alpha_n - \frac{\langle \alpha_n, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1 - \dots - \frac{\langle \alpha_n, \beta_{n-1} \rangle}{\langle \beta_{n-1}, \beta_{n-1} \rangle} \beta_{n-1}.$$

It is clear from the above discussion that $\beta_1, \beta_2, \dots, \beta_n$ are orthogonal.

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$\beta_2 = \alpha_2 + a_1 \alpha_1$ where $a_1 = -\frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}$. Therefore β_2 is a linear combination of α_1 and α_2 .

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$\beta_2 = \alpha_2 + a_1 \alpha_1$ where $a_1 = -\frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}$. Therefore β_2 is a linear combination of α_1 and α_2 .

$$\beta_3 = \alpha_3 - \frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \beta_2 - \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1.$$

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$\beta_2 = \alpha_2 + a_1 \alpha_1$ where $a_1 = -\frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}$. Therefore β_2 is a linear combination of α_1 and α_2 .

$$\beta_3 = \alpha_3 - \frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \beta_2 - \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1.$$

$$\beta_3 = \alpha_3 + b_1 \beta_2 + b_2 \beta_1 \text{ where } b_2 = -\frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \text{ and } b_1 = \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}.$$

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$\beta_2 = \alpha_2 + a_1 \alpha_1$ where $a_1 = -\frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}$. Therefore β_2 is a linear combination of α_1 and α_2 .

$$\beta_3 = \alpha_3 - \frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \beta_2 - \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1.$$

$$\beta_3 = \alpha_3 + b_1 \beta_2 + b_2 \beta_1 \text{ where } b_2 = -\frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \text{ and } b_1 = \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}.$$

$$\beta_3 = \alpha_3 + b_1(\alpha_2 + a_1 \alpha_1) + b_2 \alpha_1$$

Explain: I want to write β_3 is a linear combination of $\alpha_1, \alpha_2, \alpha_3$.

We know that:

$$\beta_1 = \alpha_1.$$

$$\beta_2 = \alpha_2 - \frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1$$

$\beta_2 = \alpha_2 + a_1 \alpha_1$ where $a_1 = -\frac{\langle \alpha_2, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}$. Therefore β_2 is a linear combination of α_1 and α_2 .

$$\beta_3 = \alpha_3 - \frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \beta_2 - \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle} \beta_1.$$

$$\beta_3 = \alpha_3 + b_1 \beta_2 + b_2 \beta_1 \text{ where } b_2 = -\frac{\langle \alpha_3, \beta_2 \rangle}{\langle \beta_2, \beta_2 \rangle} \text{ and } b_1 = \frac{\langle \alpha_3, \beta_1 \rangle}{\langle \beta_1, \beta_1 \rangle}.$$

$$\beta_3 = \alpha_3 + b_1(\alpha_2 + a_1 \alpha_1) + b_2 \alpha_1$$

Therefore β_k is a linear combination of $\alpha_1, \dots, \alpha_{k-1}$ a contradiction. Hence $\beta_i \neq 0$ for $i = 1, \dots, n$.

To show $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Therefore β_k is a linear combination of $\alpha_1, \dots, \alpha_{k-1}$ a contradiction. Hence $\beta_i \neq 0$ for $i = 1, \dots, n$.

To show $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

We have already seen that each β_i as a linear combination of $\alpha_1, \dots, \alpha_i$ for $i = 1, \dots, n$. Using construction of β_i it is clear that each α_i is a linear combination β_1, \dots, β_i for $i = 1, \dots, n$. Therefore $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Therefore β_k is a linear combination of $\alpha_1, \dots, \alpha_{k-1}$ a contradiction. Hence $\beta_i \neq 0$ for $i = 1, \dots, n$.

To show $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

We have already seen that each β_i as a linear combination of $\alpha_1, \dots, \alpha_i$ for $i = 1, \dots, n$. Using construction of β_i it is clear that each α_i is a linear combination β_1, \dots, β_i for $i = 1, \dots, n$. Therefore $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

This follows that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

- **[Example:]** Let $(\mathbb{R}^3, \langle \cdot, \cdot \rangle)$ be an inner product space and let $\langle x, y \rangle = \sum_{i=1}^3 x_i y_i$. Let $u_1 = (0, 1, 2)$, $u_2 = (1, 1, 2)$ and $u_3 = (1, 0, 1)$. Then find orthogonal vectors v_1, v_2, v_3 using u_1, u_2, u_3 .

- **[Example:]** Let $(\mathbb{R}^3, \langle \cdot, \cdot \rangle)$ be an inner product space and let $\langle x, y \rangle = \sum_{i=1}^3 x_i y_i$. Let $u_1 = (0, 1, 2)$, $u_2 = (1, 1, 2)$ and $u_3 = (1, 0, 1)$. Then find orthogonal vectors v_1, v_2, v_3 using u_1, u_2, u_3 .

Sol: $v_1 = u_1 = (0, 1, 2)$.

$$v_2 = u_2 - \frac{\langle u_2, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1.$$

$$v_2 = (1, 1, 2) - \frac{\langle (1, 1, 2), (0, 1, 2) \rangle}{\langle (0, 1, 1), (0, 1, 2) \rangle} (0, 1, 2)$$

$$v_2 = (1, 1, 2) - \frac{5}{5} (0, 1, 2) = (1, 0, 0).$$

$$v_3 = u_3 - \frac{\langle u_3, v_2 \rangle}{\langle v_2, v_2 \rangle} v_2 - \frac{\langle u_3, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1.$$

$$v_3 = (1, 0, 1) - \frac{\langle (1, 0, 1), (1, 0, 0) \rangle}{\langle (1, 0, 0), (1, 0, 0) \rangle} (1, 0, 0) - \frac{\langle (1, 0, 1), (0, 1, 2) \rangle}{\langle (0, 1, 2), (0, 1, 2) \rangle} (0, 1, 2).$$

$$v_3 = (1, 0, 1) - (1, 0, 0) - \frac{2}{5} (0, 1, 2) = (0, -\frac{2}{5}, \frac{1}{5}).$$

-
- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set.

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Proof: Using Gram Schimdt process we have an orthogonal set $\{\beta_1, \beta_2, \dots, \beta_n\}$ such that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Proof: Using Gram Schimdt process we have an orthogonal set $\{\beta_1, \beta_2, \dots, \beta_n\}$ such that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Then put $\gamma_i = \frac{1}{\sqrt{\langle \beta_i, \beta_i \rangle}} \beta_i$ for $i = 1, \dots, n$.

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Proof: Using Gram Schimdt process we have an orthogonal set $\{\beta_1, \beta_2, \dots, \beta_n\}$ such that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Then put $\gamma_i = \frac{1}{\sqrt{\langle \beta_i, \beta_i \rangle}} \beta_i$ for $i = 1, \dots, n$.

Then $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ is an orthonormal set. You can easily check that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

- Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a linearly independent set. Then there exists an **orthonormal** set $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ such that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Proof: Using Gram Schimdt process we have an orthogonal set $\{\beta_1, \beta_2, \dots, \beta_n\}$ such that $\text{ls}(\{\beta_1, \beta_2, \dots, \beta_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

Then put $\gamma_i = \frac{1}{\sqrt{\langle \beta_i, \beta_i \rangle}} \beta_i$ for $i = 1, \dots, n$.

Then $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ is an orthonormal set. You can easily check that $\text{ls}(\{\gamma_1, \gamma_2, \dots, \gamma_n\}) = \text{ls}(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$.

- [Theorem:]** Every non-trivial finite dimensional inner product space has an orthonormal basis.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis. Then for each $x \in \mathbb{V}$ we have

$$x = \sum_{i=1}^n \langle x, \alpha_i \rangle \alpha_i$$

.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis. Then for each $x \in \mathbb{V}$ we have

$$x = \sum_{i=1}^n \langle x, \alpha_i \rangle \alpha_i$$

Proof: Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is an orthonormal basis. Let $x \in \mathbb{V}$. Then there exist c_1, \dots, c_n such that $x = c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis. Then for each $x \in \mathbb{V}$ we have

$$x = \sum_{i=1}^n \langle x, \alpha_i \rangle \alpha_i$$

Proof: Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is an orthonormal basis. Let $x \in \mathbb{V}$. Then there exist c_1, \dots, c_n such that $x = c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n$.

Take $\langle x, \alpha_i \rangle = \langle c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n, \alpha_i \rangle$

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis. Then for each $x \in \mathbb{V}$ we have

$$x = \sum_{i=1}^n \langle x, \alpha_i \rangle \alpha_i$$

Proof: Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is an orthonormal basis. Let $x \in \mathbb{V}$. Then there exist c_1, \dots, c_n such that $x = c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n$.

$$\begin{aligned} \text{Take } \langle x, \alpha_i \rangle &= \langle c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n, \alpha_i \rangle \\ &= c_1\langle \alpha_1, \alpha_i \rangle + \dots + c_i\langle \alpha_i, \alpha_i \rangle + \dots + c_n\langle \alpha_n, \alpha_i \rangle \end{aligned}$$

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis. Then for each $x \in \mathbb{V}$ we have

$$x = \sum_{i=1}^n \langle x, \alpha_i \rangle \alpha_i$$

Proof: Since $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is an orthonormal basis. Let $x \in \mathbb{V}$. Then there exist c_1, \dots, c_n such that $x = c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n$.

$$\begin{aligned} \text{Take } \langle x, \alpha_i \rangle &= \langle c_1\alpha_1 + c_2\alpha_2 + \dots + c_n\alpha_n, \alpha_i \rangle \\ &= c_1\langle \alpha_1, \alpha_i \rangle + \dots + c_i\langle \alpha_i, \alpha_i \rangle + \dots + c_n\langle \alpha_n, \alpha_i \rangle \\ &= c_i \text{ for } i = 1, \dots, n. \end{aligned}$$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space.

-
- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .
- S^\perp is a subspace of \mathbb{V} for any $S \subseteq \mathbb{V}$.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .
- S^\perp is a subspace of \mathbb{V} for any $S \subseteq \mathbb{V}$.

Proof: Let $x, y \in S^\perp$ and $\alpha, \beta \in \mathbb{K}$. To show that $\alpha x + \beta y \in S^\perp$.

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .

- S^\perp is a subspace of \mathbb{V} for any $S \subseteq \mathbb{V}$.

Proof: Let $x, y \in S^\perp$ and $\alpha, \beta \in \mathbb{K}$. To show that $\alpha x + \beta y \in S^\perp$.

$$\langle \alpha x + \beta y, v \rangle = \langle \alpha x, v \rangle + \langle \beta y, v \rangle$$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .
- S^\perp is a subspace of \mathbb{V} for any $S \subseteq \mathbb{V}$.

Proof: Let $x, y \in S^\perp$ and $\alpha, \beta \in \mathbb{K}$. To show that $\alpha x + \beta y \in S^\perp$.

$$\begin{aligned}\langle \alpha x + \beta y, v \rangle &= \langle \alpha x, v \rangle + \langle \beta y, v \rangle \\ &= \alpha \langle x, v \rangle + \beta \langle y, v \rangle\end{aligned}$$

- **[Definition]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $S \subseteq \mathbb{V}$. Then the set $\{u \in \mathbb{V} \mid \langle u, v \rangle = 0 \text{ for all } v \in S\}$ is called **S perpendicular** of S and it is denoted by S^\perp .

- S^\perp is a subspace of \mathbb{V} for any $S \subseteq \mathbb{V}$.

Proof: Let $x, y \in S^\perp$ and $\alpha, \beta \in \mathbb{K}$. To show that $\alpha x + \beta y \in S^\perp$.

$$\begin{aligned}\langle \alpha x + \beta y, v \rangle &= \langle \alpha x, v \rangle + \langle \beta y, v \rangle \\ &= \alpha \langle x, v \rangle + \beta \langle y, v \rangle \\ &= 0 + 0 = 0 \text{ for all } v \in S.\end{aligned}$$

Hence S^\perp is a subspace of \mathbb{V} .

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

Using extension theorem and Gram Schmidt process we extend B to an orthonormal basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$. It is clear that $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

Using extension theorem and Gram Schmidt process we extend B to an orthonormal basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$. It is clear that $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$.

We now show that $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

Using extension theorem and Gram Schmidt process we extend B to an orthonormal basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$. It is clear that $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$.

We now show that $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$.

It is clear that $\text{ls}\{u_{k+1}, \dots, u_n\} \subseteq \mathbb{W}^\perp$ as $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$ and \mathbb{W}^\perp is a subspace.

Let $x \in \mathbb{W}^\perp$.

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

Using extension theorem and Gram Schmidt process we extend B to an orthonormal basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$. It is clear that $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$.

We now show that $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$.

It is clear that $\text{ls}\{u_{k+1}, \dots, u_n\} \subseteq \mathbb{W}^\perp$ as $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$ and \mathbb{W}^\perp is a subspace.

Let $x \in \mathbb{W}^\perp$.

$$\text{Then } x = \sum_{i=1}^n \langle x, u_i \rangle u_i$$

- **[Theorem:]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Then $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Proof: Let $\dim(\mathbb{V}) = n$ and let $\dim(\mathbb{W}) = k$.

Let $B = \{u_1, \dots, u_k\}$ be an orthonormal basis of \mathbb{W} .

Using extension theorem and Gram Schmidt process we extend B to an orthonormal basis of \mathbb{V} which is $\{u_1, \dots, u_k, u_{k+1}, \dots, u_n\}$. It is clear that $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$.

We now show that $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$.

It is clear that $\text{ls}\{u_{k+1}, \dots, u_n\} \subseteq \mathbb{W}^\perp$ as $u_{k+1}, \dots, u_n \in \mathbb{W}^\perp$ and \mathbb{W}^\perp is a subspace.

Let $x \in \mathbb{W}^\perp$.

$$\text{Then } x = \sum_{i=1}^n \langle x, u_i \rangle u_i$$

$$= \sum_{i=k+1}^n \langle x, u_i \rangle u_i; \text{ as } u_1, \dots, u_k \in \mathbb{W} \text{ and } u_{k+1}, \dots, u_n \in \mathbb{W}^\perp.$$

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let S is subspace of \mathbb{V} . Then $(S^\perp)^\perp = S$.

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let S is subspace of \mathbb{V} . Then $(S^\perp)^\perp = S$.

Proof: This can be proved using above theorem.

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let S is subspace of \mathbb{V} . Then $(S^\perp)^\perp = S$.

Proof: This can be proved using above theorem.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional inner product space. Let S and T be two subspaces of \mathbb{V} . Then the following are true.

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let S is subspace of \mathbb{V} . Then $(S^\perp)^\perp = S$.

Proof: This can be proved using above theorem.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional inner product space. Let S and T be two subspaces of \mathbb{V} . Then the following are true.

1. If $S \subseteq T \implies T^\perp \subseteq S^\perp$.

Hence $x \in \text{ls}\{u_{k+1}, \dots, u_n\}$. Therefore $\mathbb{W}^\perp = \text{ls}\{u_{k+1}, \dots, u_n\}$. It is clear that $\mathbb{W} \cap \mathbb{W}^\perp = \{0\}$. Hence $\mathbb{V} = \mathbb{W} \oplus \mathbb{W}^\perp$.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a **finite dimensional** inner product space. Let S is subspace of \mathbb{V} . Then $(S^\perp)^\perp = S$.

Proof: This can be proved using above theorem.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional inner product space. Let S and T be two subspaces of \mathbb{V} . Then the following are true.

1. If $S \subseteq T \implies T^\perp \subseteq S^\perp$.

2. $(S + T)^\perp = S^\perp \cap T^\perp$ and $(S \cap T)^\perp = S^\perp + T^\perp$

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional IPS. Let $\mathbb{W} \subseteq \mathbb{V}$ be a subspace. For each $x \in \mathbb{V}$ there exists unique $x_1 \in \mathbb{W}$ and $x_2 \in \mathbb{W}^\perp$ such that $x = x_1 + x_2$. The vector x_1 is called the **orthogonal projection** of x into W .

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional IPS. Let $\mathbb{W} \subseteq \mathbb{V}$ be a subspace. For each $x \in \mathbb{V}$ there exists unique $x_1 \in \mathbb{W}$ and $x_2 \in \mathbb{W}^\perp$ such that $x = x_1 + x_2$. The vector x_1 is called the **orthogonal projection** of x into W .
- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be a finite dimensional inner product space. Let \mathbb{W} be a subspace of \mathbb{V} . Let $\{\alpha_1, \dots, \alpha_k\}$ be an orthogonal basis of \mathbb{W} . Then the orthogonal projection of any vector $x \in \mathbb{V}$ into \mathbb{W} is
$$\sum_{i=1}^k \frac{\langle x, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle} \alpha_i.$$

Let V be a vector space over the field \mathbb{K} .

Let \mathbb{V} be a vector space over the field \mathbb{K} .

A map $\|\cdot\| : \mathbb{V} \rightarrow \mathbb{R}$ is said to be a norm on \mathbb{V} if it satisfies the following condition.

Let \mathbb{V} be a vector space over the field \mathbb{K} .

A map $\|\cdot\| : \mathbb{V} \rightarrow \mathbb{R}$ is said to be a norm on \mathbb{V} if it satisfies the following condition.

1. $\|x\| \geq 0$ for all $x \in \mathbb{V}$ and $\|x\| = 0$ if and only if $x = 0$.

Let \mathbb{V} be a vector space over the field \mathbb{K} .

A map $\|\cdot\| : \mathbb{V} \rightarrow \mathbb{R}$ is said to be a norm on \mathbb{V} if it satisfies the following condition.

1. $\|x\| \geq 0$ for all $x \in \mathbb{V}$ and $\|x\| = 0$ if and only if $x = 0$.
2. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{K}$ and for all $x \in \mathbb{V}$.

Let \mathbb{V} be a vector space over the field \mathbb{K} .

A map $||, || : \mathbb{V} \rightarrow \mathbb{R}$ is said to be a norm on \mathbb{V} if it satisfies the following condition.

1. $||x|| \geq 0$ for all $x \in \mathbb{V}$ and $||x|| = 0$ if and only if $x = 0$.
2. $||\alpha x|| = |\alpha| ||x||$ for all $\alpha \in \mathbb{K}$ and for all $x \in \mathbb{V}$.
3. $||x + y|| \leq ||x|| + ||y||$ for all $x, y \in \mathbb{V}$.

Let \mathbb{V} be a vector space over the field \mathbb{K} .

A map $||, || : \mathbb{V} \rightarrow \mathbb{R}$ is said to be a norm on \mathbb{V} if it satisfies the following condition.

1. $||x|| \geq 0$ for all $x \in \mathbb{V}$ and $||x|| = 0$ if and only if $x = 0$.
2. $||\alpha x|| = |\alpha| ||x||$ for all $\alpha \in \mathbb{K}$ and for all $x \in \mathbb{V}$.
3. $||x + y|| \leq ||x|| + ||y||$ for all $x, y \in \mathbb{V}$.

A vector space together with a norm $||, ||$ is called a **normed linear space**.

The following are examples of norm.

1. Let $x \in \mathbb{R}^n$. Then $\|x\| = \sum_{i=1}^n |x_i|$ is a norm on \mathbb{R}^n .

The following are examples of norm.

1. Let $x \in \mathbb{R}^n$. Then $\|x\| = \sum_{i=1}^n |x_i|$ is a norm on \mathbb{R}^n .

1. Each $|x_i| \geq 0$ for $i = 1, \dots, n$, then $\sum_{i=1}^n |x_i| \geq 0$. Hence $\|x\| \geq 0$ for all $x \in \mathbb{R}^n$.

The following are examples of norm.

1. Let $x \in \mathbb{R}^n$. Then $\|x\| = \sum_{i=1}^n |x_i|$ is a norm on \mathbb{R}^n .

1. Each $|x_i| \geq 0$ for $i = 1, \dots, n$, then $\sum_{i=1}^n |x_i| \geq 0$. Hence $\|x\| \geq 0$ for all $x \in \mathbb{R}^n$.

If $\|x\| = 0 \implies \sum_{i=1}^n |x_i| = 0 \implies |x_i| = 0$ for $i = 1, \dots, n$

$x_i = 0$ for $i = 1, \dots, n$. Then $x = 0$. If $x = 0$, then $\|x\| = \sum_{i=1}^n 0 = 0$.

The following are examples of norm.

1. Let $x \in \mathbb{R}^n$. Then $\|x\| = \sum_{i=1}^n |x_i|$ is a norm on \mathbb{R}^n .

1. Each $|x_i| \geq 0$ for $i = 1, \dots, n$, then $\sum_{i=1}^n |x_i| \geq 0$. Hence $\|x\| \geq 0$ for all $x \in \mathbb{R}^n$.

If $\|x\| = 0 \implies \sum_{i=1}^n |x_i| = 0 \implies |x_i| = 0$ for $i = 1, \dots, n$

$x_i = 0$ for $i = 1, \dots, n$. Then $x = 0$. If $x = 0$, then $\|x\| = \sum_{i=1}^n 0 = 0$.

2. $\|\alpha x\| = \sum_{i=1}^n |\alpha x_i| = \sum_{i=1}^n |\alpha| |x_i| = |\alpha| \sum_{i=1}^n |x_i| = |\alpha| \|x\|$ for all $x \in \mathbb{R}^n$ and for $\alpha \in \mathbb{K}$.

The following are examples of norm.

1. Let $x \in \mathbb{R}^n$. Then $\|x\| = \sum_{i=1}^n |x_i|$ is a norm on \mathbb{R}^n .

1. Each $|x_i| \geq 0$ for $i = 1, \dots, n$, then $\sum_{i=1}^n |x_i| \geq 0$. Hence $\|x\| \geq 0$ for all $x \in \mathbb{R}^n$.

If $\|x\| = 0 \implies \sum_{i=1}^n |x_i| = 0 \implies |x_i| = 0$ for $i = 1, \dots, n$

$x_i = 0$ for $i = 1, \dots, n$. Then $x = 0$. If $x = 0$, then $\|x\| = \sum_{i=1}^n 0 = 0$.

2. $\|\alpha x\| = \sum_{i=1}^n |\alpha x_i| = \sum_{i=1}^n |\alpha| |x_i| = |\alpha| \sum_{i=1}^n |x_i| = |\alpha| \|x\|$ for all $x \in \mathbb{R}^n$ and for $\alpha \in \mathbb{K}$.

3.

$\|x+y\| = \sum_{i=1}^n |x_i + y_i| \leq \sum_{i=1}^n (|x_i| + |y_i|) = \sum_{i=1}^n |x_i| + \sum_{i=1}^n |y_i| = \|x\| + \|y\|$
for all $x, y \in \mathbb{R}^n$.

- **[Cauchy Schwarz Inequality]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an IPS. Let $x, y \in \mathcal{V}$. Then $|\langle x, y \rangle|^2 \leq \langle x, x \rangle \langle y, y \rangle$.

Equality holds if and only if x and y are linearly dependent.

- **[Cauchy Schwarz Inequality]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an IPS. Let $x, y \in \mathbb{V}$. Then $|\langle x, y \rangle|^2 \leq \langle x, x \rangle \langle y, y \rangle$.

Equality holds if and only if x and y are linearly dependent.

Proof: If $y = 0$, then it is trivial.

- **[Cauchy Schwarz Inequality]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an IPS. Let $x, y \in \mathbb{V}$. Then $|\langle x, y \rangle|^2 \leq \langle x, x \rangle \langle y, y \rangle$.

Equality holds if and only if x and y are linearly dependent.

Proof: If $y = 0$, then it is trivial.

If $y \neq 0$, take $x - ty$ where $t \in \mathbb{C}$. Then

- **[Cauchy Schwarz Inequality]** Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an IPS. Let $x, y \in \mathbb{V}$. Then $|\langle x, y \rangle|^2 \leq \langle x, x \rangle \langle y, y \rangle$.

Equality holds if and only if x and y are linearly dependent.

Proof: If $y = 0$, then it is trivial.

If $y \neq 0$, take $x - ty$ where $t \in \mathbb{C}$. Then

$$0 \leq \langle x - ty, x - ty \rangle$$

$$= \langle x, x \rangle + \langle x, -ty \rangle + \langle -ty, x \rangle + \langle -ty, -ty \rangle$$

$$= \langle x, x \rangle - \bar{t} \langle x, y \rangle - t \langle y, x \rangle + |t|^2 \langle y, y \rangle$$

$$\text{Put } t = \frac{\langle x, y \rangle}{\langle y, y \rangle}.$$

$$\langle x, x \rangle - \frac{\overline{\langle x, y \rangle}}{\langle y, y \rangle} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle$$

$$\begin{aligned} \langle x, x \rangle &= \overline{\frac{\langle x, y \rangle}{\langle y, y \rangle}} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle \\ &= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{(\langle y, y \rangle)^2} \langle y, y \rangle \end{aligned}$$

$$\begin{aligned} \langle x, x \rangle &= \overline{\langle x, y \rangle} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle \\ &= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{(\langle y, y \rangle)^2} \langle y, y \rangle \\ &= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \end{aligned}$$

$$\langle x, x \rangle - \frac{\overline{\langle x, y \rangle}}{\langle y, y \rangle} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle$$

$$= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{(\langle y, y \rangle)^2} \langle y, y \rangle$$

$$= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle}$$

$$\langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \geq 0$$

$$\begin{aligned} \langle x, x \rangle - \frac{\overline{\langle x, y \rangle}}{\langle y, y \rangle} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle \\ = \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{(\langle y, y \rangle)^2} \langle y, y \rangle \\ = \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \\ \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \geq 0 \end{aligned}$$

Then $|\langle x, y \rangle| \leq (\langle x, x \rangle)^{1/2} (\langle y, y \rangle)^{1/2}$.

$$\begin{aligned} \langle x, x \rangle - \frac{\overline{\langle x, y \rangle}}{\langle y, y \rangle} \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \langle y, x \rangle + |t|^2 \langle y, y \rangle \\ = \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{(\langle y, y \rangle)^2} \langle y, y \rangle \\ = \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \\ \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \geq 0 \end{aligned}$$

Then $|\langle x, y \rangle| \leq (\langle x, x \rangle)^{1/2} (\langle y, y \rangle)^{1/2}$.

The first inequality which we have used in this proof is $0 \geq \langle x - ty, x - ty \rangle$. If $|\langle x, y \rangle|^2 = \langle x, x \rangle \langle y, y \rangle$ hold, then $\langle x - ty, x - ty \rangle = 0$. This says that $x = ty$. Then x and y are linearly dependent.

- Let $(\mathbb{V}, \langle \cdot, \cdot \rangle)$ be an inner product space. Let $x \in \mathbb{V}$. Then we can easily check that $\|x\| = (\langle x, x \rangle)^{1/2}$ is a norm on \mathbb{V}