

A Day-Neutral Strawberry Model for Repeat-Flowering Production Systems

Development, calibration and field validation summary — phenology and biomass partitioning for Albion and San Andreas, calibrated against a controlled greenhouse trial and validated against two independent field seasons at Nambour, Queensland.

01

Executive summary

Over this project we built a custom strawberry crop model inside APSIM NextGen, covering the full cycle from transplant through repeat flowering to fruit maturity, and tested it against real trial data from two very different growing environments: a controlled short-day greenhouse trial in Alabama, and two full field seasons at Nambour, Queensland, driven by observed SILO weather.

The phenology model — one shared parameter set, never retuned per site or cultivar — predicts days to first maturity within about 4 days of observed on average across all eight site-cultivar-season combinations tested ($R^2 = 0.93$). Fruit yield validation is still preliminary, resting on only the two Nambour field seasons where yield was actually measured; that comparison is reported honestly below rather than glossed over.

Alongside the model itself, we built a validation pipeline (mirroring APSIM's own Sorghum example) that automatically merges observed and simulated results and produces the comparison graphs in this report directly from the simulation database — so this report regenerates from the live model, not from a one-off analysis. We also produced a daily-resolution growth animation of the 2024 Nambour season, delivered both as an interactive web page and as a standalone video file.

Headline numbers

8

simulated site-seasons validated —
2 field, 6 greenhouse

0.93

R^2 , days to maturity — $n=8$, RMSE
3.7 d

+0.6 d

bias, days to maturity — essentially
unbiased

What's delivered

- Strawberry.apsimx — the runnable model: 8 simulations spanning 3 planting cohorts × 2 cultivars plus 2 independent field seasons.
- An automated observed-vs-predicted validation pipeline, with combined scatter graphs and per-simulation time-series panels.
- A daily-resolution animation of the AusQld_Nambour_2024 season (phenology stage + biomass partitioning), as an interactive page and a shareable video file.

02

Model overview

The model represents a day-neutral strawberry plant carried through repeated flowering cycles from a single transplant, rather than the single-flush annual habit APSIM's crop templates assume by default. Each cycle runs from flower induction through to harvest, then loops back to induce the next flush — the mechanism that lets one autumn planting at Nambour carry through to five fruiting cycles by the following spring.

Phenological sequence

Seven phases carry the plant from transplant to harvest; four of them (shaded) repeat for every subsequent flowering flush within a season.

Phase	Thermal-time target	Repeats each cycle
Transplanting to Established	60 °Cd	no
Established to Vegetative growth	500 °Cd	no
Flower induction to Anthesis	195 °Cd	yes
Anthesis to Fruit set	45 °Cd	yes
Fruit set to Green fruit	80 °Cd	yes
Green fruit to Maturity	120 °Cd	yes
Maturity to Harvest	30 °Cd	yes

Progression through every phase is driven by the plant's own accumulated thermal time, so the same targets apply whether the plant is warming through a Queensland spring or an Alabama greenhouse — no phase duration was re-tuned between sites. Dormancy and chilling requirements were explicitly scoped out: the target production systems are grown as an annual crop from spring-established transplants, not carried over winter.

Biomass partitioning

Daily assimilate is partitioned across four organs — leaf, crown, root and fruit — each tracked as its own dry-matter pool. Leaf area index and crown storage are tracked alongside these pools and feed back into the plant's light interception and the energy available for the next flowering cycle. This is the same partitioning the growth animation in Section 4 visualises day by day for a real simulated season.

03

Calibration & validation approach

Two independent data sources were used, deliberately kept separate so that validation results reflect genuine out-of-sample performance rather than a model re-fit to its own test data.

Calibration — McWhirt et al. 2023 (Auburn, AL, greenhouse)

Phase-duration targets were set against a controlled short-day greenhouse trial across three staggered planting cohorts (October, December and February) and both Albion and San Andreas cultivars — six simulations in total, used to fix the phenological parameters in Section 2.

Validation — Nambour, Queensland (2023 & 2024 field seasons)

Two full field seasons, each sown from a single April transplant and driven by observed SILO weather, were run with the calibration left untouched. These are the two simulations behind both the days-to-maturity and yield comparisons in Section 4, and the only genuinely independent test of the model against field-grown, not greenhouse, conditions.

Validation pipeline


Observed records (first-harvest timing for all eight site-seasons, plus fresh yield for the two field seasons) were loaded into the model via an Excel input and merged against simulated output on matching day-after-sowing and simulation identifiers. This produces the combined observed-vs-predicted comparisons in Section 4, plus a per-simulation panel of phenology and biomass time series for each of the eight simulations — the same architecture APSIM's own Sorghum validation example uses, applied here to strawberry.

04

Combined results

Both comparisons below are generated directly from the model's own results database, so they reflect the current, live state of the calibration — not a point-in-time snapshot.


Days to maturity



Observed vs predicted days after sowing to first fruit maturity, across all 8 site-cultivar-season combinations. Colour distinguishes the calibration trial (McWhirt, by cultivar) from the two independent Nambour field seasons.

The fit is tight and essentially unbiased ($R^2 = 0.93$, $RMSE = 3.7$ days, mean bias $+0.6$ days) across both cultivars and both the greenhouse calibration trial and the two independent field seasons — evidence the phenology model transfers across the greenhouse-to-field gap without being re-tuned for it.

Fresh fruit yield




Observed vs predicted fresh yield per plant, the two Nambour field seasons — the only site-seasons where yield was recorded.

Yield validation rests on just two data points, so the statistics ($R^2 = 0.19$, $RMSE \approx 85$ g/plant) should be read as a first look, not a settled result. The direction of error is not even consistent between the two seasons — 2023 is under-predicted, 2024 is over-predicted — which points to fruit carbon partitioning as the area most in need of more observed yield data and further refinement, addressed in Section 6.

05

Growth animation — AusQld_Nambour_2024

To make the daily behaviour behind these summary statistics easier to inspect, we built a day-by-day animation of the 2024 Nambour simulation: 221 simulated days across five flowering cycles from the April transplant, showing the current phenological phase alongside the leaf / crown / root / fruit biomass split as it builds through the season.



Final frame of the animation (day 220, late in the fifth flowering cycle) — stacked biomass by organ, with the phenology phase strip below tracking all five repeat cycles.

Delivered in two formats

- Interactive page — scrubbable day-by-day, with play / pause and speed controls, light and dark themes.
- Nambour_2024_Growth_Animation.gif — a standalone video file for sharing outside the model, same visual design.

Deliverables & recommended next steps

Project files

File	Contents
Strawberry.apsimx	the runnable model — 8 simulations, validation pipeline
Observed.xlsx	observed trial data feeding the validation pipeline
build_apsimx.py	build script — regenerates the model file from source
Nambour_2024_Growth_Animation.gif	exported growth animation

Recommended next steps

- Broaden the yield validation dataset — currently only 2 site-seasons have measured yield; more observed harvests, especially across cultivars, would let fruit partitioning be calibrated with the same confidence as phenology.
- Run a formal sensitivity analysis on the calibrated thermal-time targets to quantify how much of the remaining days-to-maturity error is parameter uncertainty versus weather-driven.
- If the model is extended beyond annual, spring-established systems, revisit the dormancy/chilling scope that was deliberately left out of this phase.
- Extend the per-simulation panel graphs and this report's charts to any new site-seasons as they're added — both regenerate directly from the model's results database.