

Problem set 5: Cosmic Microwave Background

1)

```
In [ ]: import numpy as np
import camb
from matplotlib import pyplot as plt
import time

plt.rc('text', usetex=True)
plt.rc('font', family='serif')
plt.rcParams['figure.dpi'] = 100

# pset 5
# 1)

# calculate model power spectra from parameters
def get_spectrum(pars, lmax=3000):
    H0=pars[0]
    ombh2=pars[1]
    omch2=pars[2]
    tau=pars[3]
    As=pars[4]
    ns=pars[5]
    pars=camb.CAMBparams()

    pars.set_cosmology(H0=H0, ombh2=ombh2, omch2=omch2, mnu=0.06, omk=0, tau=tau)
    pars.InitPower.set_params(As=As, ns=ns, r=0)
    pars.set_for_lmax(lmax, lens_potential_accuracy=0)
    results=camb.get_results(pars)
    powers=results.get_cmb_power_spectra(pars, CMB_unit='muK')
    cmb=powers['total']
    tt=cmb[:,0]
    return tt[2:]

# parameters dialed into script
pars=np.asarray([60,0.02,0.1,0.05,2.00e-9,1.0])
```

```


# CMB data from Planck satellite
planck=np.loadtxt('COM_PowerSpect_CMB-TT-full_R3.01.txt',skiprows=1)
ell=planck[:,0] # multipole, starting with l=2
spec=planck[:,1] # variance of sky as a function of multipole
errs=0.5*(planck[:,2]+planck[:,3]) # error: average of upper/lower 1 sigma
uncertainty
model=get_spectrum(pars)
model=model[:len(spec)]
resid=spec-model
chisq=np.sum( (resid/errs)**2)
print("Chi squared for initial values of parameters is ",chisq," for
",len(resid)-len(pars)," degrees of freedom.")
#read in a binned version of the Planck PS for plotting purposes
planck_binned=np.loadtxt('COM_PowerSpect_CMB-TT-
binned_R3.01.txt',skiprows=1)
errs_binned=0.5*(planck_binned[:,2]+planck_binned[:,3]);
plt.clf()
plt.plot(ell,model,label='Power spectrum')
plt.title('Power spectrum for initial parameters')
plt.xlabel('$l$')
plt.ylabel('spectrum')
plt.errorbar(planck_binned[:,0],planck_binned[:,1],errs_binned,fmt='.',label
data')
plt.legend()
plt.show()

# new parameters
pars=np.asarray([69,0.022,0.12,0.06,2.1e-9,0.95])
model=get_spectrum(pars)
model=model[:len(spec)]
resid=spec-model
chisq=np.sum( (resid/errs)**2)
print("Chi squared for new parameter is ",chisq," for ",len(resid)-
len(pars)," degrees of freedom.")
plt.clf()
plt.plot(ell,model, label='Power spectrum')
plt.title('Power spectrum for new parameters')
plt.xlabel('$l$')
plt.ylabel('spectrum')


```

```
plt.errorbar(planck_binned[:,0],planck_binned[:,1],errs_binned,fmt='.',label
            = 'Planck data')
plt.legend()
plt.show()
```

Chi squared for initial values of parameters is 15267.937150261656 for 2501 degrees of freedom.



Chi squared for new parameter is 3272.205355920218 for 2501 degrees of freedom.



The mean and variance for χ^2 is n and $2n$ respectively, where n is the degree of freedom. Since

we have 2501 degrees of freedom, we expect a chi squared with mean $\mu = 2501$ ($\mu = n$) and standard deviation $\sigma = 70.72$ ($\sigma = \sqrt{2n}$). The χ^2 value for the parameters dialed into the test script is 15267.937150261656. This is incredibly high, it is absolutely not within 1 σ of the mean (it's within 181 σ though!).

The χ^2 for the new parameters is 3272.205355920218. This much better than before, but still not within 1 σ of the mean (it's within 11 σ). None of these fits are acceptable.

2) Levenberg-Marquardt to find best fit parameters

```
In [ ]: # function to take the derivative
def centered_derivative(params,i):
    # some arbitrary dx; will not really be optimal
    dx=1e-9
    params_left=params.copy();params_right=params.copy()
    params_right[i]+=dx
    params_left[i]-=dx
    # two points centered derivative
    d=0.5*(get_spectrum(params_right)-get_spectrum(params_left))/(dx)
    return d

# function to take derivative with respect to all parameters
def derivative(params):
    grad = np.zeros([model.size, params.size])
    for i in range(params.size):
        temp=centered_derivative(params,i)
        temp=temp[:len(planck[:,0])]
        grad[:,i]=temp
    return grad

# initial guess are the params from 1)
params=np.asarray([69,0.022,0.12,0.06,2.1e-9,0.95])
error=0.5*(planck[:,2]+planck[:,3])

# Levenberg-marquardt method to find best fit parameters
t=2 # Levenberg-marquardt parameter
d=get_spectrum(params)
d=d[:len(planck[:,0])]
r=planck[:,1]-model
chisq=np.sum((r/error)**2)
```

```

# Let's do 70 iterations for fun
for i in range(70):
    grad=derivative(params)
    grad=grad[:len(planck[:,0])]
    r=planck[:,1]-d
    lhs = grad.T@grad
    lhs = lhs + t*np.diag(np.diag(lhs))
    rhs = grad.T@r
    # shift in parameters
    dp = np.linalg.inv(lhs)@rhs
    params_new = params+dp

    # calculate new chi squared with these new 'trial' parameters
    d=get_spectrum(params_new)
    d=d[:len(planck[:,0])]
    r=planck[:,1]-d
    chisq_trial=np.sum((r/error)**2)

    # keep the trial parameters if new chi squared is better
    # reduce Levenberg-marquardt parameter
    if chisq_trial<chisq:
        t=t/2
        if t<0.2:
            t=0
        params=params_new
        chisq = chisq_trial

    # if new chi squared is worse, don't take new parameters
    # make Levenberg-marquardt parameter bigger
    if chisq_trial>chisq:
        if t==0:
            t=2
        else:
            t=t*3

pred=get_spectrum(params)
pred=pred[:len(planck[:,0])]
grad=derivative(params)

```

```

N=np.mean((planck[:,1]-pred)**2)
Ninv=1/N
covariance=np.linalg.inv(Ninv*grad.T@grad) # covariance matrix
params_err=np.sqrt(np.diag(covariance)) # error on parameters

```


```

In [ ]: # writing best fit parameters and errors in 'planck_fit_params.txt'
lines = ['Best fit parameters with error are:',
'H0= '+str(params[0])+' with error of '+str(params_err[0]),
'omb= '+str(params[1])+' with error of '+str(params_err[1]),
'omc= '+str(params[2])+' with error of '+str(params_err[2]),
'tau= '+str(params[3])+' with error of '+str(params_err[3]),
'As= '+str(params[4])+' with error of '+str(params_err[4]),
'ns= '+str(params[5])+' with error of '+str(params_err[5])]
with open('planck_fit_params.txt','w') as f:
    for line in lines:
        f.write(line)
        f.write('\n')

# plotting stuff, look at the txt file for the values!
print("Chi squared for best fit parameters is ",chisq," for ",len(resid)-
len(pars)," degrees of freedom.")
plt.plot(planck[:,0],pred, label='Power spectrum with best fit parameters
from Levenberg-Marquardt')
plt.xlabel('$l$')
plt.ylabel('spectrum')
plt.errorbar(planck_binned[:,0],planck_binned[:,1],errs_binned,fmt='.',label
= 'Planck data')
plt.legend()
plt.show()

```

Chi squared for best fit parameters is 2582.15066327397 for 2501 degrees of freedom.



χ^2 for fit with best parameters obtained with Levenberg-Marquardt is 2582.15066327397. This is pretty good, as it is within 2σ of the mean χ^2

3) Markov Chain Monte Carlo to find best fit parameters

```
In [ ]: # 3) MCMC

# function to calculate the chi squared during the chain
def get_chisq(params):
    d=get_spectrum(params)
    d=d[:len(planck[:,0])]
    r=planck[:,1]-d
    chisq=np.sum((r/error)**2)
    return chisq

# Markov Chain Monte Carlo
def run_mcmc(params,n=5000):
    chain=np.zeros([n,len(params)])
    chivec=np.zeros(n)

    # chi squared at current location
    chi=get_chisq(params)
    for i in range(n):
```

```

        # trial parameter, use covariance matrix from 2) to get new trial
        parameters
        params_trial=np.random.multivariate_normal(params,covariance)
        # chi squared with trial parameters
        chi_trial=get_chisq(params_trial)
        # probability of accepting step
        accept_prob=np.exp(-0.5*(chi_trial-chi))
        if np.random.rand(1)<accept_prob: # accept step with appropriate
        probability
            params=params_trial
            chi=chi_trial
            chain[i,:]=params
            chivec[i]=chi
        return chain,chivec


# same guess for params as 2)
params=np.asarray([69,0.022,0.12,0.06,2.1e-9,0.95])
# Let's do 5000 steps!
chain,chivec=run_mcmc(params,5000)

```

```

In [ ]: # check if chain converged
plt.plot(chivec)
plt.xlabel('Number of steps')
plt.ylabel('$\chi^2$')
plt.title('Evolution of $\chi^2$ in the MCMC chain')
plt.show()

```

Based on the plot above, the chain clearly converged to some value of χ^2 after a very little amount of steps, after which χ^2 just oscillates a little around that value. Because of this oscillation, what I'll do is pick out the smallest χ^2 the chain outputted and get the parameters associated to that step. These will be my best fit parameters for the chain.

```
In [ ]: # chi squared oscillates after chain converged
# so just pick the smallest chi squared and the parameters associated to
it
a=np.where(chivec==np.min(chivec))
chisq_mcmc=chivec[a[0][0]] # best fit chisq
params_mcmc=chain[a[0][0]] # best fit params

# getting covariance matrix and error on params from mcmc chain
pred=get_spectrum(params_mcmc)
pred=pred[:len(planck[:,0])]
grad=derivative(params_mcmc)
N=np.mean((planck[:,1]-pred)**2)
Ninv=1/N
covariance=np.linalg.inv(Ninv*grad.T@grad) # covariance matrix
params_err=np.sqrt(np.diag(covariance)) # error on parameters

# saving chain in 'planck_chain.txt', first column is the chisquared
# other columns are parameters in order
```


```

lines=np.empty([len(chain),7])
lines[:,0]=chivec
lines[:,1]=chain[:,0]
lines[:,2]=chain[:,1]
lines[:,3]=chain[:,2]
lines[:,4]=chain[:,3]
lines[:,5]=chain[:,4]
lines[:,6]=chain[:,5]
np.savetxt('planck_chain.txt',lines)
print('Best fit parameters using MCMC are: ',params_mcmc)
print('best fit chisq using MCMC is: ', chisq_mcmc)
# plotting stuff
plt.plot(planck[:,0],pred, label='Power spectrum with best fit parameters
from MCMC')
plt.xlabel('$l$')
plt.ylabel('spectrum')
plt.errorbar(planck_binned[:,0],planck_binned[:,1],errs_binned,fmt='.',label
= 'Planck data')
plt.legend()
plt.show()

```

Best fit parameters using MCMC are: [6.89661801e+01 2.25618488e-02 1.16354494e-01 9.69329436e-02 2.26412716e-09 9.76234731e-01]

best fit chisq using MCMC is: 2576.9639845303773



χ^2 is now 2576.9639845303773. This is better than what I got from running the Levenberg-Marquardt fit (I should hope so because the MCMC chain took several hours to run). However, this is still not within 1σ of the mean value of χ^2 (2510 ± 70.72). It is within 2σ though, not bad.

Estimating the mean value of the dark energy Ω_Λ , using $\Omega_b + \Omega_c + \Omega_\Lambda = 1$.

```
In [ ]: # finding the mean value of the dark energy using parameters found with
mcmc

omb=(100**2*params_mcmc[1])/(params_mcmc[0])**2
omc=(100**2*params_mcmc[2])/(params_mcmc[0])**2
oml=1-omb-omc

# do some error propagation to find error on mean dark energy
omb_error=omb*np.sqrt((params_err[0]/params_mcmc[0])**2+
(params_err[1]/params_mcmc[1])**2)
omc_error=omc*np.sqrt((params_err[0]/params_mcmc[0])**2+
(params_err[2]/params_mcmc[2])**2)
oml_error=oml*np.sqrt((omb_error/omb)**2+(omc_error/omc)**2)

print('My estimate of the mean value of the dark energy is: ',oml, 'with
uncertainty of: ',oml_error)
```

My estimate of the mean value of the dark energy is: 0.7079340035099035 with uncertainty of: 0.011418849435390723

4) Markov Chain Monte Carlo with fixed tau

Use the same mcmc as in 3) but this time we have a fixed tau with fixed value of 0.0540 ± 0.0074 . This means that every step of the chain, I need to fix the value of tau to some value taken from a univariate gaussian distribution with mean 0.0540 and standard deviation 0.0074. This can be done using `np.random.normal(mean, std)`.

```
In [ ]: # Markov Chain Monte Carlo with fixed tau

def run_mcmc_fixed_tau(params,n=5000):
    chain=np.zeros([n,len(params)])
    chivec=np.zeros(n)

    # chi squared at current location
    chi=get_chisq(params)
    for i in range(n):
```

```

        # trial parameter, use covariance matrix from 2) to get new trial
        parameters
        params_trial=np.random.multivariate_normal(params,covariance)
        # fixed tau
        params_trial[3]=np.random.normal(0.0540,0.0074)
        # chi squared with trial parameters
        chi_trial=get_chisq(params_trial)
        # probability of accepting step
        accept_prob=np.exp(-0.5*(chi_trial-chi))
        if np.random.rand(1)<accept_prob: # accept step with appropriate
        probability
            params=params_trial
            chi=chi_trial
            chain[i,:]=params
            chivec[i]=chi
        return chain,chivec

# same guess for params as 3) but with the appropriate value of tau
params=np.asarray([69,0.022,0.12,0.0540,2.1e-9,0.95])


# Let's do 5000 steps again
chain_fixed,chivec_fixed=run_mcmc_fixed_tau(params,n=5000)

```

```

In [ ]: # check if chain converged
plt.plot(chivec_fixed)
plt.xlabel('Number of steps')
plt.ylabel('$\chi^2$')
plt.title('Evolution of $\chi^2$ in the MCMC chain')
plt.show()

```



Chain converged pretty fast again. Same thing here, I'll pick the smallest χ^2 and the parameters associated to it as the best fit parameters.

```
In [ ]: # chi squared oscillates after chain converged
# so just pick the smallest chi squared and the parameters associated to
it
a=np.where(chivec_fixed==np.min(chivec_fixed))
chisq_mcmc_fixed=chivec_fixed[a[0][0]] # best fit chisq
params_mcmc_fixed=chain_fixed[a[0][0]] # best fit params
# saving chain in 'planck_chain_tauprior.txt', first column is the
chisquared
# other columns are parameters in order
lines=np.empty([len(chain),7])
lines[:,0]=chivec_fixed
lines[:,1]=chain_fixed[:,0]
lines[:,2]=chain_fixed[:,1]
lines[:,3]=chain_fixed[:,2]
lines[:,4]=chain_fixed[:,3]
lines[:,5]=chain_fixed[:,4]
lines[:,6]=chain_fixed[:,5]
np.savetxt('planck_chain_tauprior.txt',lines)
print('Best fit parameters using MCMC with tau prior are:
',params_mcmc_fixed)
```

```

print('best fit chisq using MCMC with tau prior is: ', chisq_mcmc_fixed)


temp=get_spectrum(params_mcmc_fixed)
temp=temp[:len(planck[:,0])]

plt.plot(planck[:,0],temp, label='Power spectrum with best fit parameters
from MCMC with fixed tau')
plt.xlabel('$l$')
plt.ylabel('spectrum')
plt.errorbar(planck_binned[:,0],planck_binned[:,1],errs_binned,fmt='.',label
= 'Planck data')
plt.legend()
plt.show()

```

Best fit parameters using MCMC with tau prior are: [6.78827422e+01 2.23749836e-02 1.18708926e-01 5.71353128e-02 2.10281184e-09 9.70503772e-01]

best fit chisq using MCMC with tau prior is: 2577.270677067728



In []: *# Let's compare those results to what I get from importance sampling my chain from problem 3*

```

def fixed_chisq(params,params_fixed,errs):
    return np.sum(((params-params_fixed)/errs)**2)

expected_params=np.zeros(params.size)

```

```

expected_params[3]=0.0540
expected_errs=1e20*np.ones(params.size)
expected_errs[3]=0.0074

# chi squared vector from importance sampling
importance_chivec=np.zeros(np.size(chivec))
for i in range(np.size(chivec)):
    chi=fixed_chisq(chain[i,:],expected_params,expected_errs)
importance_chivec=importance_chivec-importance_chivec.mean()
# weights
w=np.exp(0.5*importance_chivec)

# parameters from importance sampling
importance_params=np.zeros(params.size)
for i in range(params.size):
    importance_params[i]=np.sum(w*chain[:,i])/np.sum(w)

print('Chi squared from importance sampling my MCMC chain from problem 3
is: ', get_chisq(importance_params))
print('Chi squared from running a MCMC with fixed tau is (problem 4) is:
', get_chisq(params_mcmc_fixed))

```

Chi squared from importance sampling my MCMC chain from problem 3 is: 2577.9705341706585

Chi squared from running a MCMC with fixed tau is (problem 4) is: 2577.270677067728

χ^2 is similar but slightly higher when importance sampling my chain from problem 3). However neither of those χ^2 fall within 1σ of 2501.