



Biomass Coverage in Other Marine Percoidids Taxonomy Group




Biomass Coverage Over Time in Other Marine Percoidids Taxonomy Group

Stocks with highest average biomass




Biomass Coverage Over Time in Other Marine Percoidids Taxonomy Group

Stocks with lowest average biomass




Red snapper Gulf of Mexico – Other Marine Percoidids Taxonomy Group

Stock represents 5% of average absolute surplus production




Bluefish Atlantic Coast – Other Marine Percoidids Taxonomy Group

Stock represents 30% of average absolute surplus production




Red grouper Gulf of Mexico – Other Marine Percoidids Taxonomy Group

Stock represents 13% of average absolute surplus production




Australian salmon New Zealand – Other Marine Percoidids Taxonomy Group

Stock represents 4% of average absolute surplus production




Red snapper Gulf of Mexico in Other Marine Percoidids Taxonomy Group




First and last years with recorded surplus production are labeled

Bluefish Atlantic Coast
in Other Marine Percoidids Taxonomy Group




First and last years with recorded surplus production are labeled

Red grouper Gulf of Mexico in Other Marine Percoidids Taxonomy Group



Australian salmon New Zealand in Other Marine Percoidids Taxonomy Group



First and last years with recorded surplus production are labeled