
TCIA CT Colonography
825 patients, 3,451 CT series;
prone and supine
CTSpine1K, COLONOG subset
784 vertebral masks,
VerSe ids (L6 = 25)
CTPelvic1K, dataset2
714 sacrum and hip masks;
tags in the filenames
place each mask on the series it was drawn on
Neither source records a series UID, so among one patient's series the one with the most bone (HU > 200) under the mask wins: spine by world-space resampling, pelvis by bone z-profile cross-correlation.
804 patients: 342 fused (same series), 352 separate (different series), 90 spine only, 20 pelvis only
orientation and pairing review of the QC renders (21 April)
9 patients looked antero-posteriorly inverted: CT and masks mirrored front to back. The two checked since (tokens 75, 480) are left–right mirror images of the patient (liver on the left).
On the 6 whose pelvis was mirrored, a TotalSegmentator audit (26–27 April) disagreed with the left and right hip labels, which were then swapped.
1 separate patient excluded: its spine mask did not lie on the bone of its chosen series. 803 patients
export (25 April; last rewritten 1 May)
CT and label share one affine (PIR); every CT is resampled into its label's grid (fix of 27 April: a CT with its mask's shape but not its affine had skipped this and was written with scrambled axes).
Ten classes: 0 background, 1–6 L1–L6, 7 sacrum (CTPelvic1K's over CTSpine1K's), 8 left hip, 9 right hip.
In a record with one annotation, every other voxel is 10 = ignore, not background (1 May).
1 spine-only patient not exported (reason not recorded). 802 patients, 1,153 records
fused
n = 342
342 records
separate
n = 351
702 records (2 each)
spine only
n = 89
89 records
pelvis only
n = 20
20 records
six-way subtype per patient, from the source annotations; no image was read
CTSpine1K lumbar count and CTPelvic1K filename tag, ambiguous where they disagree.
A substring parser had read semisacralization as sacralization (2 patients); fixed 30 April.
normal
769
lumbarization
14
sacralization
(count-style)
9
sacralization
(full)
6
semi-
sacralization
2
ambiguous
2
training view (nnU-Net, 3 May)
L5 and L6 merged into last_lumbar: nine classes plus ignore. 174 records held out; five folds over the other 979. The split is by record: 92 separate patients have one record on each side.
Castellvi I–IV, the 33 LSTV patients
(added after the release)
Two radiology residents' reads, complete by 4 September 2026, agree on 27 of 33 (κ = 0.74); consensus on the other 6.
Ib 2, IIa 4, IIb 6, IIIa 1, IIIb 16, IV 4