

# CTSpinoPelvic1K: spine, pelvis, ribs and femora in one coordinate frame, annotated for lumbosacral transitional anatomy

Gregory Schwing<sup>a)</sup>

*Department of Surgery, Detroit Medical Center and Wayne State University, Detroit, Michigan, USA*

Ashley Schehr, Annika Tekumulla, Margret Khoushi, Ryan Christian, Dane Hubers, Faris Mahjoub, Hassan Saad, Mia Sooch, Sathyagopal Siddapureddy, Michael McLellan, and Jerick Kim

*School of Medicine, Wayne State University, Detroit, Michigan, USA*

Miraziz Ismoilov and Nizar Alnabahneh

*Department of Radiology, Detroit Medical Center and Wayne State University, Detroit, Michigan, USA*

(Dated: 10 September 2026)

**Purpose:** A vertebra at the lumbosacral junction is named by counting. The gold standard counts caudally from C2 on whole-spine imaging<sup>1</sup>; however, a lumbar surgical case is planned on lumbar-only imaging<sup>2-4</sup> (T12 to S1) without C2. Abdominopelvic computed tomography (CT) holds that span plus the lowest ribs and pelvis, standing in for the preoperative view. Where a lumbosacral transitional vertebra (LSTV) alters the count, the local anatomy is ambiguous. Four rib-free vertebrae may be an L1 with a lumbar rib or an L5 assimilated to the sacrum, and six may be a sixth lumbar vertebra, a T12 with aplastic ribs, or a lumbarized S1<sup>5</sup>. CTSpinoPelvic1K asks whether local morphology resolves that ambiguity without the count. Two public label sets covered this imaging, CTSpine1K’s vertebrae and CTPelvic1K’s pelvis, on the same colonography patients, never joined; this release joins them on one series and annotates the bones neither had. It provides 802 CT records, per-level ribs and femora, lumbar levels anchored on the lowest rib-bearing vertebra and S1, and classes for L6, T13, a separate S1 and lumbar ribs, so anomalies are recorded as themselves.

**Acquisition and Validation Methods:** Records pair CTSpine1K<sup>6</sup> and CTPelvic1K<sup>7</sup> labels on each patient’s bone-richest acquisition under a VerSe-native scheme. Validation covered geometric invariants (802/802 pass), rib-vertebra incidence across 5,749 ribs (0.035% offset), and spinopelvic measures matching published values.

**Data Format and Usage Notes:** NIfTI image/label pairs with patient-grouped LSTV-stratified five-fold splits and a loader; archived at <https://doi.org/10.5281/zenodo.22139642>.

**Potential Applications:** Classifying a vertebra from local features; updating cadaveric morphometry; spinopelvic assessment; opportunistic screening; and, absent a public preoperative lumbar cohort, surgical planning research (377 records prone). *Limitations:* thoracic ground truth is field-of-view limited; postural angles are supine; no held-out test set; ribs are triaged-review pseudolabels; Castellvi grades are two-reader consensus on 33 records.

## I. INTRODUCTION

The spine commonly carries seven cervical, twelve thoracic and five lumbar vertebrae, and a vertebra is identified by where it falls in that sequence. Transitional variants sit at the two ends of the lumbar column, the thoracolumbar transitional vertebra (TLTV) above and the lumbosacral transitional vertebra (LSTV) below, and they disturb identification in two ways at once. They change what the border vertebra and its ribs *look like*: a thoracolumbar border vertebra may carry a full rib, a hypoplastic stump, or none at all, and a lumbosacral one may be partly or completely assimilated to the sacrum (*sacralization*), with an anteriorly wedged body, a reduced disc beneath it and hypoplastic facet joints, or conversely separated from it (*lumbarization*), with a squared

body, lumbar-type facets and a full-height disc<sup>5</sup>. They also change *how many* vertebrae each region contains, eleven or thirteen thoracic and four or six lumbar.

Either change alone would be tractable. Together they make it difficult to state definitively what any vertebra at these borders *is*. Four rib-free vertebrae above the sacrum may mean an L1 bearing a lumbar rib or an L5 assimilated to it; six may mean a true sixth lumbar vertebra, a T12 whose ribs are aplastic, or an S1 lumbarized away from the sacrum. Each set yields the same count (Fig. 2), and the morphology that would separate the readings is not decisive to a human reader in every case. Whether it differs enough for a learned classifier to separate them, or to assign each reading a probability, is the question this release is built to support. Absent

that, what remains is the count itself. The gold standard for numeration in transitional anatomy is whole-spine imaging, counting caudally from C2<sup>1</sup>, which fixes how many vertebrae and how many rib pairs the column holds, and no spine-limited acquisition can supply either count. LSTV is common, reported at 4–30% in the general population<sup>5</sup> and in 16.3% of a whole-spine CT series,<sup>8</sup> and wrong-level surgery is its most serious consequence.

Existing public collections cannot address this, and size is not the reason (Table I): spine datasets omit the pelvis, pelvic datasets do not number the vertebrae, TotalSegmentator has no class for a sixth lumbar vertebra or for a rib on a lumbar vertebra, and VerSe, the one collection with an L6 class, stops at the sacrum and carries no ribs. No public collection can therefore record a six-lumbar spine together with both of its borders, and a segmenter trained on them must, faced with an enumeration anomaly, shift the whole column by a level or absorb a vertebra into its neighbor. The size of that problem has now been measured on CT, where prior labeling methods assigned every vertebra correctly in 77% of subjects and an anomaly-aware extension of SPINEPS<sup>9</sup> raised that to 99%<sup>10</sup>. Its weights label L6 and T13 on CT, but its 1,536-scan CT cohort is in-house and unreleased, so the capability exists as weights, not as data.

The intraoperative side has prior art of its own. LevelCheck registers the intraoperative radiograph to the preoperative CT and projects the CT’s vertebral labels onto it<sup>11–13</sup>, but those labels are placed by hand and verified by the surgeon<sup>11</sup>, so the method assumes a correctly labeled CT and does not supply one. Automating that step where the anatomy is anomalous is what a dataset carrying the anomaly classes is for.

CTSpinoPelvic1K places the spine, pelvis, ribs and femora in one coordinate frame and gives a class to each structure an enumeration anomaly produces, a sixth lumbar vertebra and a rib borne by a lumbar vertebra. A scheme without those classes lacks a class vocabulary expressive enough to describe anatomic anomalies.

The project began with a surgeon’s question, whether a vertebra can be named from the anatomy in front of the reader, on the images a lumbar surgical case is actually planned on. By guideline those are lumbar-only imaging studies<sup>2,3,14</sup>, T12 to S1, the span the ACR spine parameters state for the lumbar spine<sup>4</sup>, and whole-spine coverage is reserved for trauma with an identified injury<sup>15,16</sup> and for deformity radiographs. An abdominopelvic CT runs from the diaphragm to the pelvic floor, so it holds that span together with the lowest ribs and the whole pelvis, and it lacks C2 exactly as the planning study does. Every record comes from one prospective trial protocol, ACRIN 6664<sup>17,18</sup>, which scanned 802 patients aged 50 and over, supine and prone, on multidetector CT at 15 centers, on five scanner vendors and nine models, with manufacturer, model and reconstruction kernel recorded per record. To our knowledge it is the largest spine-and-pelvis annotated CT cohort acquired under a single protocol, where the

comparators pool collections, are multi-site by design, or are routine clinical scans under many protocols (Table I). With the protocol held fixed, scanner effects on a model become measurable rather than confounded. The price is an abdominal field of view, in which only the lowest thoracic levels are present (Sec. V).

The gap was in plain sight. CTPelvic1K<sup>7</sup> and CT-Spine1K<sup>6</sup>, released months apart by an overlapping author group, each annotated the COLONOG collection under radiologist supervision, one the pelvis and the other the vertebrae, on the same patients, and no one joined them, although the join needs no new imaging and no new radiologist. It is not a file merge, because each annotation had to be traced back to the series it was drawn on (Sec. II.A). Done once, that crosswalk yields a spine-and-pelvis frame at 802 patients, and the bones neither set had, ribs, femora, S1 and hardware, could then be annotated against it rather than from scratch.

### I.A. Vertebra labeling when the scan cannot settle the count

*The limitation, stated plainly.* No scan in this corpus contains C2, so the conventional count cannot be performed, a property of abdominal imaging rather than of the annotation. A thirteenth thoracic vertebra and a lumbar rib are different phenotypes, distinguished as separate subtypes in cadaveric classifications of the thoracolumbar junction<sup>19,20</sup>: the first is an additional rib-bearing segment, usually above five lumbar vertebrae, so the column holds 25 presacral vertebrae; the second is a rudimentary rib on a lumbar-type L1 in a column of the usual 24. Between the two borders used here, the last rib-bearing vertebra and the sacrum, none of them is separable by counting: four rib-free vertebrae fit a lumbar rib, a cranially shifted T13 or an L5 assimilated to the sacrum, and six fit an aplastic twelfth rib or a sixth lumbar vertebra. A *stump rib*, hypoplastic and a cardinal indicator of a thoracolumbar transitional vertebra, is still a rib, so only its length records the anomaly. Rib anomalies travel with the lumbosacral border: on whole-spine CT hypoplastic twelfth ribs occur with sacralization and lumbar ribs with lumbarization.<sup>8</sup>

*What the release does in addition.* Every vertebra carries the identifier its radiologist-sourced annotation assigned, corrected where the source was wrong; those labels are the ground truth. Every released measurement is also expressed against the two landmarks present in essentially every abdominal scan, the sacrum below and the lowest rib-bearing vertebra above, so that the measurement is positional rather than nominal: a vertebra by how many rib-free vertebrae separate it from the sacrum, a lowest rib by its length as a fraction of the rib above it, a transverse process by its span and its distance from the ala. None of these changes with the name a reader gives the junction, so cases that disagree about the name remain comparable.

*But the vertebrae themselves are known to differ.*

Level-specific morphometry is long established: pedicle width and height change systematically from the thoracic to the lumbar spine,<sup>21</sup> vertebral body, endplate and canal dimensions differ by level,<sup>22–24</sup> and the transverse process and the iliolumbar ligament mark the last lumbar vertebra independently of any count.<sup>5,25</sup> If those differences hold at the boundary itself the phenotype is decidable locally, and on this corpus it does. Separating T12 from L1 on shape alone gives an area under the curve of 0.990 and 97.3% accuracy over 1485 vertebrae: a logistic regression on five released per-level measures and six ratios between them, each measure divided by the same patient’s median across their levels, so that size, which separates thoracic from lumbar trivially and says nothing at the junction, is removed; folds are grouped by case and the curve is taken on pooled out-of-fold scores (`test_morphometric_separability.py` in the released code). Schinz et al. reach the same conclusion from the other side: on 1,242 whole-thoracolumbar CTs a shape-based labeling of the junction matched nerve morphology in every case, against 92.6–97.2% for counting- and rib-based rules.<sup>26</sup>

## II. ACQUISITION AND VALIDATION METHODS

### II.A. Sources, and why a crosswalk was necessary

Both sources draw part of their imaging from The Cancer Imaging Archive (TCIA) CT colonography collection (COLONOG)<sup>17,18</sup>. CTSpine1K’s 1,005 volumes come from four collections and CTPelvic1K’s 1,184 from seven; COLONOG is the only collection common to both and the only one acquired under a single protocol, meaning one scan prescription and patient preparation for every case, so this release is its COLONOG subset, 782 patients with a CTSpine1K vertebral annotation and 20 with a CTPelvic1K pelvic annotation only, 802 in all. Neither source released the mapping from an annotation to the specific CT series it was drawn on.

That mapping matters because every patient in CT COLONOGRAPHY was scanned *twice*, prone and supine, and an annotation carries a patient identifier and nothing finer, so neither collection recorded which acquisition it had labeled. Turning a patient from supine to prone alters lumbar lordosis and the alignment of each segment against the next, so outlines drawn on one series are the wrong *shape* for the other and no rigid realignment recovers them. The crosswalk (Fig. 1) therefore resolves each annotation to a patient and then, within that patient’s volumes, to the series whose bone *agrees* with it, where candidates are thresholded at bone attenuation and scored by overlap with the mask. It mattered for 351 patients, nearly half the cohort, whose two collections had labeled different acquisitions of the same person; those records are released but held out of any analysis that assumes a single posture, two labeled postures of one patient being a resource rather than a defect.




FIG. 1 How the dataset was built.  $n$  counts scans, or corrections, at each step.

### II.B. Densification, and why it is asymmetric

A corpus in which half the records lack a pelvis is not a spinopelvic dataset, so the missing structures were pseudolabeled, asymmetrically (Fig. 1). Pelvis onto spine-only records is the safe direction, since the sacrum and innominate bones carry no enumeration to get wrong, and quality is reported as held-out performance on the *pelvic-only* records, withheld from training for that purpose. The reverse direction carries the whole problem this dataset addresses, because a pseudolabeled spine must commit to a count, and on a transitional vertebra it commits to the commoner reading; those 20 records were corrected by hand.

### II.C. Ribs

No public rib annotation covers this cohort, and the two available automatic sources<sup>30,31</sup> fail in complementary ways.

Möller’s binary rib network<sup>31</sup> reports high accuracy,

TABLE I Public CT collections at the lumbosacral junction. Black, class present; outlined, graded only to exclude it<sup>27</sup>.

| Collection                     | Scans | Count | L6 | Sacrum | S1 | Pelvis | Ribs | L. rib | Femora | Grade |
|--------------------------------|-------|-------|----|--------|----|--------|------|--------|--------|-------|
| CTSpine1K <sup>6</sup>         | 1,005 |       |    |        |    |        |      |        |        |       |
| CTPelvic1K <sup>7</sup>        | 1,184 |       |    |        |    |        |      |        |        |       |
| VerSe <sup>28</sup>            | 374   |       |    |        |    |        |      |        |        |       |
| RibSeg v2 <sup>29</sup>        | 660   |       |    |        |    |        |      |        |        |       |
| TotalSegmentator <sup>30</sup> | 1,204 |       |    |        |    |        |      |        |        |       |
| <b>CTSpinoPelvic1K</b>         | 802   |       |    |        |    |        |      |        |        |       |

and its authors show that stump ribs can be classified from a partial field of view; that result concerns the morphological features computed on a rib once segmented. On this corpus the segmentation itself failed on ribs that were *partially outside* the field of view. A rib crossing the boundary of an abdominal acquisition was liable to be missed altogether rather than segmented to the edge. In a colonography cohort the lower ribs are routinely clipped, and those are the ribs the count depends on.

TotalSegmentator<sup>30</sup> does segment clipped ribs and supplies the per-rib numbering that a binary network cannot. Its characteristic failure is at the other end of the bone, where the most medial portion of the rib (roughly the last tenth to sixth approaching the costovertebral joint) is frequently absent. A rib truncated at its medial end never reaches the vertebra it articulates with, so the rib-vertebra incidence check (the quality-control gate on the rib class labels, which matches each rib to the vertebra its number implies) has nothing to test.

The two were therefore unioned (Fig. 1), TotalSegmentator supplying the numbering and the medial-clipped cases, Möller the detection where its output is more complete.

The result is a pseudolabel whose review was triaged rather than exhaustive. A label-based rule flagged every record in which one rib bone carried more than one identifier, or in which a rib failed to reach the vertebra it articulates with, since a single bone carrying two numbers is a numbering error whatever the anatomy. The rule selected 152 records; 148 were reviewed and corrected, and four were not and are identified in the release. Reviewers were medical students working through OpenSpineConsortium<sup>32</sup>, trained against a written labeling protocol, with every correction attributed to its annotator. Records passing the rule were not individually inspected; the residual rib-vertebra offsets reported in Sec. II.F are the independent check on what the rule missed.

## II.D. Label scheme and counting anchors

The scheme is VerSe-native. Vertebrae keep their VerSe identifiers (C1–C7 = 1–7, T1–T12 = 8–19, L1–L6 = 20–25, sacrum 26, coccyx 27, T13 = 28), and every non-VerSe structure takes a fixed identifier above that range, S1 carved from the sacrum (29), hips (30–31), femora (32–33), thirteen ribs per side (34–46 left, 47–59 right), lumbar ribs (60–61) and surgical hardware (62–

68). The space is contiguous. Rib 13 is the true rib of a T13 and is empty in this release, so a thirteenth thoracic vertebra and a lumbar rib are recorded as the distinct phenotypes they are.

Two anchors give every record the same positional frame, and Fig. 2 shows both, the lowest rib-bearing vertebra above, **S1** carved from the sacrum below. S1 supplies the sacral endplate landmark that sacral slope and pelvic incidence are measured from, and with both brackets present L5 and L6 are distinguishable where a spine-limited view alone would not be.

*How S1 is cut.* The sacrum’s outer boundary is the source annotation and is never altered; S1 is the cranial part of that same bone, separated by a plane. The plane’s normal is the sacrum’s own cranio-caudal axis, made orthogonal to a left–right axis measured directly rather than assumed, so as to remove the patient’s rotation within the scanner, which is not negligible: a median of 4.5°, a maximum of 16.1°, and 508 of 801 records above 3°. The cut preserves the sub-division volume of the preceding release, changing the orientation of the S1/S2 boundary and not how much sacrum is called S1; per-record geometry ships with the archive.

VerSe<sup>28</sup> carries L6, whose identifier this release keeps, so its L6 is VerSe’s.

**A rib on a lumbar body receives its own class**, and this is the one class here with no published counterpart. A lumbar rib and a stump rib are the same object under two counts, so a scheme that numbers every rib 1–12 leaves the annotator either calling it rib 12, asserting the vertebra beneath it is thoracic, the very question at issue, or discarding it. TotalSegmentator carries ribs 1–12 per side and no such class, and VerSe’s T13 is a vertebra rather than a rib. This scheme records both: VerSe’s T13 (28) with its rib pair (46, 59) for a thoracic-type thirteenth vertebra, and 60/61 for a lumbar-type body bearing a rudimentary rib. No record here carries a T13; the sixteen lumbar-rib cases were read as lumbar-type bodies, and where a border vertebra could not be settled by its readers the label stands as read.

Figure 2 shows why the interval is the primitive and the label is derived. Among the records with four rib-free vertebrae, some reach that count from above, through a rib on L1, and some from below, through a lumbosacral segment fused to the sacrum, in Fig. 2(c) together with stump twelfth ribs; the count alone cannot tell them apart, and the release can because ribs, rib lengths and the S1 boundary are all labeled.






FIG. 2 The two anchors, from the released labels. Red, lowest rib-bearing vertebra and its rib; blue, S1; yellow, the rib-free vertebrae between. (b) and (c) both count four. In (b) a long rib sits on L1; in (c) the twelfth ribs are stumps and the segment below the fourth lumbar body is fused to the sacrum (Castellvi IIb) and carries the S1 identifier.

Sixteen records carry a lumbar rib, thirteen bilateral and three unilateral, all on the right, too few to speak to side preference.

Eighteen records carry an L6. Seventeen of them carry a consensus Castellvi grade; the eighteenth was not graded. The source transitional label (pelvic where CT-Pelvic1K flagged one, otherwise the lumbar count) reads lumbarization in fourteen, sacralization in two and semi-sacralization in one, and is unremarkable in the ungraded record. A spine can carry six lumbar-type vertebrae and still be read as a sacralization, depending on where the reader started the count.

Both counts, and the manifest fields that report them (`has_l6`, `has_lumbar_rib`, `n_lumbar_labels`, `lumbar_rib_side`), are computed from the released label volumes by counting identifiers 25 and 60–61.

## II.E. Computational tools

*Access.* Every step is performed by open-source code archived with the dataset, together with the reference loader, the label scheme and the quality-control scripts that generated the tables in this manuscript.

*Versions and parameters.* Segmentation of femora, the sacral sub-division and the per-level rib numbering used TotalSegmentator<sup>30</sup> (total task, release 2.x) on a single graphics processing unit. The binary rib network is Möller’s<sup>31</sup> released weights, applied through the nnU-Net v2 framework<sup>33</sup>. The pelvic completion for records whose pelvis was absent used a five-fold nnU-Net v2 ensemble applied out-of-fold, so that no record is completed by a model that has seen it. The five fold checkpoints, with their nnU-Net plans and dataset descriptor, are released alongside the data (<https://huggingface.co/OpenSpineConsortium/spinopelvic-seg-checkpoints>, trained with <https://github.com/Gregory-Schwing-MD-PhD/spinesurg-ct-nnnet> at commit 50b209f). Image handling throughout used NiBabel and SciPy; no image is resampled or reoriented when a label is written, so every label shares its image’s grid and affine exactly.

*Measurement.* Dimensions here come from code

released with the dataset. A reusable implementation that reads each between landmarks in the vertebra’s own frame, rather than along the scanner’s axes, is released separately as OpenSpineToolkit, a unit-tested open-source package (<https://github.com/OpenSpineConsortium/OpenSpineToolkit>). The two agree within a millimetre on canal depth; they differ most on posterior body height, where the frame removes a tilt term worth several millimetres at L5.

*Generative artificial intelligence.* A large language model (Claude, Anthropic) assisted with drafting and editing the manuscript text and with writing the analysis and figure code, all under author review. It was not used to generate, annotate or interpret imaging data; every number reported here is computed by the released code from the released volumes.

## II.F. Validation

Validation has three layers. Geometry is checked first, because a measurement taken on a corpus with a geometric fault still returns a plausible-looking number.

*Geometric invariants.* Every case is checked for label geometry agreeing with its CT in shape, affine and voxel spacing; for identifiers within the published scheme; for a non-empty label; and for sided structures falling on the correct sides, with the left–right axis read from the affine rather than assumed.

The check compares *each sided pair separately*. Testing the ribs alone passed all 802 records, four of which carried a `left_hip` label on the patient’s right side; pooling every sided structure into one test still missed one of those four records, because a large correctly-sided structure masks a smaller transposed one. A gate that passes everything is not evidence of a clean corpus until it has been shown capable of failing.

*Rib–vertebra incidence.* Each rib is matched to its nearest vertebra and compared against the vertebra its number implies. Of 11,548 ribs, 5,788 are not evaluable because the expected vertebra lies outside the field of view and 11 have no vertebra within the anchor distance; of the 5,749 evaluable, 5,747 match, two (0.035%) are

offset by one level, and no case is misnumbered. The denominator is stated because quoting the two offsets against all 11,548 ribs would halve the rate by counting ribs the check never examined. Both offsets are field-of-view truncations, one at +1 level and one at -1.

The same check carries an invariant that reads as a null result. Of the 5,749 evaluable ribs, *zero* are assigned to a lumbar vertebra, not because no lumbar ribs exist, since 16 records carry one, but because such a rib takes class 60 or 61 and never enters the numbered set the test examines. A numbered rib landing on a lumbar vertebra would be a counting error of exactly the kind this dataset exists to expose.

*Agreement with published values.* Derived spinopelvic measures are compared against Vialle *et al.*<sup>34</sup> (Table II). Pelvic incidence is the strongest of these checks because it is a morphological property of the pelvis rather than a posture, and so is directly comparable to a standing reference cohort; it agrees to within a degree. Lumbar lordosis sits below the standing reference, the direction recumbency predicts.

## II.G. Castellvi typing

Every record whose vertebral count departs from five carries a Castellvi grade<sup>35</sup> in the released manifest, 33 cases typed I–IV with the *a/b* unilateral–bilateral qualifier. The grade and the count are *different axes*. Grade IIIb occurs here at rib-free counts of four, five and six alike, and seven of the 33 graded cases carry a normal count of five.

Two radiology resident physicians graded every case independently and resolved their six disagreements by consensus; both reads and the consensus are in the manifest. The distinction the readers disagreed on, type II (articulation) against type III (bony fusion)<sup>5</sup>, is the one that matters clinically, and four of the six disagreements were III against II.

## II.H. Surgical hardware, and why a threshold is not enough

Metal is trivial to detect and hard to interpret, and the difference matters here because **an iatrogenic fusion is indistinguishable from a congenital one to a distance measurement**. A cage-bridged interspace reads as “no gap” exactly as a fused transitional vertebra does.

*II.H.0.a. Detection over-calls by a factor of five.* A 2500 HU threshold inside a shell around the labeled skeleton flags 52 of the 802 records. All 52 were reviewed manually, confirming 11 and rejecting 41 (79%) as contrast, calcification and reconstruction artefact. Attenuation does not separate the two groups (both saturate); site and volume do, every confirmed implant at 2,586 mm<sup>3</sup> or above and every rejection at 1,768 mm<sup>3</sup> or below.

*II.H.0.b. The subtype block had to be extended.* Identifiers 62–65 name spinal instrumentation (generic, cage, screw-and-rod, plate) and cannot describe what this cohort holds, so **66 arthroplasty**, **67 sacroiliac screw**

and **68 osteosynthesis** were added. Osteosynthesis holds parts of one bone together where arthroplasty replaces a joint, and a shape rule calls a femoral stem a rod. The confirmed cases are eight arthroplasties, one osteosynthesis, one sacroiliac fixation and one pair of interbody cages (Fig. 3).

*II.H.0.c. Metal outranks bone, and that is a subtraction.* In every confirmed case the implant was already inside a bone label, so naming it reclaims 1,538,852 voxels rather than adding them. In eight of these cases the femoral head that pelvic incidence and tilt are measured from is an implant.

## III. DATA FORMAT AND USAGE NOTES

Each record is a gzipped NIFTI image/label pair sharing an identical 4 × 4 affine, canonicalized to PIR, requiring no resampling. Masks are NIFTI rather than DICOM, which is what volumetric segmentation tooling expects.

Splits are patient-grouped and LSTV-stratified five-fold, frozen and shipped with the data as `splits_5fold.json`. The five validation folds partition all 802 records, so the release provides cross-validation and *not* a held-out test set; users reporting a single held-out number should carve one out and state which records it holds.

Stratification is on the transitional subtype rather than a binary LSTV flag. Across all 802 records the source transitional label reads lumbarization in 14, sacralization in 17 and semi-sacralization in two; separately, 18 records carry a sixth lumbar-type vertebra, nine carry only four lumbar labels, and 33 carry a consensus Castellvi grade. A record called sacralization by its four lumbar labels enters the same stratum as one called sacralization by the source label, so the two rare strata hold 15 records each and every validation fold carries three of each. That is the most stratification can guarantee, and it is why a fold-level metric on the rare classes carries an error bar far wider than its own decimal places.

The archive of record is Zenodo (<https://doi.org/10.5281/zenodo.22139642>). Code and documentation are on GitHub at the tagged release commit; any model-hub copy is a convenience mirror, not the archive.

## IV. DESCRIPTIVE ANALYSIS

The cohort is a colorectal cancer screening population of 802 records. 738 are recorded female (393) or male (345); 11 carry DICOM’s *other* value and 53 no sex field. Age is present for 709 (median 59, range 50–89). All 802 are counted in totals; the eleven outside the female/male dichotomy are excluded from sex-stratified measures.

*Count-free measures* (Fig. 4). Five rib-free vertebrae above the sacrum is typical; four and six are where transitional anatomy sits, and which is present does not by itself determine what to call it. The lowest-rib length ratio is bimodal, and its lower mode, below 0.33 of the rib above, marks a stump twelfth rib in 98 of the 788 records

TABLE II Derived measures against a standing reference ( $n = 260$ )<sup>34</sup>; this cohort is supine.

| Measure          | This dataset (supine) | Reference (standing) |
|------------------|-----------------------|----------------------|
| Pelvic incidence | 54.7°                 | 55 ± 10.6            |
| Pelvic tilt      | 17.3°                 | 13 ± 6               |
| Sacral slope     | 36.9°                 | 41.0 ± 8.4           |
| Lumbar lordosis  | 52.7°                 | 60 ± 10              |
| PI – LL mismatch | 2.6°                  | centered near 0      |


FIG. 3 Instrumentation in the release, one exemplar each, drawn through bone. (a) Hip arthroplasty, 66,  $n = 8$ . (b) Femoral-neck screws, 68,  $n = 1$ . (c) Sacroiliac screws, 67,  $n = 1$ . (d) Interbody cages, 63,  $n = 1$ . Bar, 5 cm.

TABLE III Data types and metadata, and the records populating each, from the released manifest.

| Field                                                                         | Records | Available |
|-------------------------------------------------------------------------------|---------|-----------|
| Identifiers and paths, configuration, match type, image/label alignment check | 802     | 100.0%    |
| Patient position, tube potential, section thickness, kernel                   | 802     | 100.0%    |
| Transitional label and class, annotation origin, instrumentation flags        | 802     | 100.0%    |
| Spine series identifier; bone-fraction check                                  | 782     | 97.5%     |
| Scanner manufacturer and model                                                | 768     | 95.8%     |
| Sex                                                                           | 749     | 93.4%     |
| Age                                                                           | 709     | 88.4%     |
| Pelvic series identifier                                                      | 713     | 88.9%     |
| Castellvi grade (consensus of two readers)                                    | 33      | 4.1%      |

with a measurable pair (12.4%, against 12.6% on whole-spine CT<sup>8</sup>); 16 records carry a lumbar rib (2.0%, against 1.5%). The co-occurrence reported there is testable here only where the junction was read; stump ribs accompany four of the 14 source-labeled sacralizations and 94 of the other 758 records (odds ratio 2.8,  $p = 0.09$ , Fisher’s exact test), and no lumbarization carries a lumbar rib.

*Spinopelvic measures* (Fig. 5). Pelvic incidence sits on the standing reference, as a morphological property rather than a posture should. Sacral slope falls below it and pelvic tilt above, the direction recumbency predicts.

*Opportunistic measures.* Every scan carries a vertebral bone density measurement at no additional dose.<sup>36</sup> L1 trabecular attenuation is reported at the published standard site.

## V. POTENTIAL APPLICATIONS AND LIMITATIONS

The principal use is the *spinopelvic interface*. Spine and pelvis carry radiologist-sourced annotation on one coordinate frame in 802 records, making sacral morphology measurable against the lumbar column rather than in isolation, among them the sacral endplate and its slope, the alar corridors constraining S2-alar-iliac and iliac screw trajectories, and the L5–S1 geometry a lum-

bosacral fusion must cross. Pelvic incidence is derived from the segmented sacrum and femoral heads, so it is reproducible from the labels themselves.

The transitional layer serves that use rather than competing with it, since no trajectory can be planned across a junction whose segments cannot be named.

*A proxy for the preoperative view.* No public preoperative lumbar CT cohort exists that we know of, so this release is the closest available stand-in for the view a lumbar case is planned on: it holds the T12-to-S1 span with the lowest ribs and the pelvis, 377 of its 802 records were acquired prone, the position of posterior lumbar surgery, and it carries classes for the anomalies that make level identification fail. It is a screening population at colonography dose, not a surgical series, so it supports method development for level identification and instrumentation geometry rather than validation of a planning decision.

### V.A. Three uses the released measurements support

*Naming a level from its own geometry.* Wrong-level spine surgery runs at roughly one in 3,110 spinal procedures, and among the predominant contributing factors is the transitional anatomy this cohort was assembled around.<sup>37,38</sup> Per-level body height, canal width, endplate width, transverse-process span and wedge ratio are re-




FIG. 4 Count-free measures. (a) Vertebrae between the lowest rib and the sacrum. (b) Lowest rib length over the rib above. (c) Lowest lumbar transverse span against its gap to the ala; transitional labels marked.




FIG. 5 Spinopelvic measures against standing reference bands (mean  $\pm$  SD)<sup>34</sup>.

leased for T11–L5 across all 802 records, and because the labels were assigned against the twelfth rib rather than by enumeration, a model trained on them is not learning the convention that fails.

*Reference morphometry with its spread.* The values a surgeon consults for vertebral body height, canal dimensions and pedicle width come from cadaveric series of twelve to thirty specimens<sup>22–24</sup> and are redrawn in the textbooks as one curve per level.<sup>39</sup> A standard error, where one is reported, describes the mean and not the spread of individuals, so neither tells the reader whether the patient in front of them is unusual. The same measures are given here from 802 records with the distribution attached (Fig. 6), reproducing two properties of the classical figures, body height crossing over at T11–T12 and reversing by L4–L5 and pedicle width widening into the lower lumbar spine, and adding the width of each distribution, the part needed to judge an individual.

*Patient-specific models.* Finite-element studies increasingly apply patient-specific modeling to alignment planning and implant design.<sup>40</sup> Such a model needs spine, pelvis and femoral heads in one frame, which is this release’s construction; in 351 patients two postures give a within-patient change in alignment against which a predicted postural response can be checked.

*Limitations: coverage and cohort.* Thoracic ground truth is field-of-view limited (Fig. 7) and does not reach T1. Postural angles are supine, though pelvic incidence is not postural and needs no such caveat. The cohort is one acquisition protocol of a colorectal screening population aged 50 and over, so its distributions should not be

read as a surgical one, and generalization to pathologic anatomy is untested.

*Label strength varies by structure, and the release does not average over that.* Vertebral labels derive from radiologist-supervised source annotations, corrected where those sources were wrong; pelvic labels on records that lacked one are pseudolabeled. The rib layer is a pseudolabel whose review was triaged, so its guarantee is the absence of the failure modes that rule detects. Transitional labels come from two independent sources and are not uniformly adjudicated, which is why the measures reported here are count-free; the Castellvi grades are a consensus of two radiology resident physicians. S1 is carved by an automatic estimate of the S1–S2 boundary; in 100 records that estimate is implausible, making S1 more than half the sacrum’s height or under fifteen percent of it, and those records are flagged in the quality-control table for exclusion from any measurement that depends on the sacral endplate.

*Absent structures, and no classifier at the lower border.* Sacrum, hips and femora are present in all 802 records; one record carries no S1 and one no T12, each outside the field of view. Nine records carry no L5 identifier because their source annotation counted four lumbar vertebrae, all nine are Castellvi IIb, and the fused transitional segment carries the S1 identifier as the caudal anchor; the manifest names them. No shape-based classifier is attempted at the lumbosacral junction, where 15 records in each rare stratum leave three per fold, too few to estimate from; the 0.990 area under the curve reported above is the thoracolumbar boundary, not this one.






FIG. 6 Morphometry by level, as median, interquartile range and 5th-95th percentile, with  $n$  at right. (a) Body height. (b) Endplate width. (c) Canal. (d) Pedicle width. (e) Disc height. (f) Trabecular attenuation. Black spans give the range of published means for the same measure, five to eighteen series per panel, one tick each; they disagree among themselves by 21 to 54% of their own median.




FIG. 7 Records carrying each vertebral level, cranial to caudal. Counts per identifier in Table S1.

## VI. DISCUSSION

*Comparison with related material.* VerSe<sup>27,28</sup> comes closest in intent but has no sacral mask for the structure a Castellvi grade describes; this release segments it, as L6 or as a carved S1. Extending VerSe with a sacrum would not have supplied the pelvis, femora or prone acquisitions, and what this release adds over TotalSegmentator<sup>30</sup> is the classes an anomaly needs, not more anatomy per scan.

*Future directions.* Applying the released weights to VerSe would add the S1 anchor and the anomaly classes to a bone-kernel collection. A Castellvi read of the whole cohort would test the co-occurrence of rib and lumbosacral anomalies at full power and supply the labels a lumbosacral classifier needs.

## VII. CONCLUSION

CTSpinoPelvic1K places spine, pelvis, per-level ribs and femora on one coordinate frame in 802 records, and gives the anatomy that makes lumbar numbering ambiguous a class of its own. Because both counting anchors are explicit, a level can be identified from a field of view that cannot support the conventional count downward

from C2.

## DATA AVAILABILITY

The release described here is v10, permanently archived at <https://doi.org/10.5281/zenodo.22642578>; the concept DOI <https://doi.org/10.5281/zenodo.22139642> resolves to the current version. The archive carries the loader, the label scheme, the quality-control tables and the analysis code under an open-source licence. Code: <https://github.com/OpenSpineConsortium/CTSpinoPelvic1K>; convenience mirror: <https://huggingface.co/datasets/OpenSpineConsortium/CTSpinoPelvic1K>; measurement toolkit: <https://github.com/OpenSpineConsortium/OpenSpineToolkit>.

## ACKNOWLEDGEMENTS

Acknowledgements are given on the title page, per the journal's double-anonymized review policy. This work received no funding.

## CONFLICT OF INTEREST

The authors have no relevant conflicts of interest to disclose.

## ETHICS

This work uses publicly available, de-identified imaging and annotations derived from it. The authors' institutional review board determined that it is not human

participant research under 45 CFR 46 and requires no oversight.

<sup>a</sup>) Electronic mail: [gregory.schwing@med.wayne.edu](mailto:gregory.schwing@med.wayne.edu)

- <sup>1</sup>J. Lian, N. Levine, and W. Cho, A review of lumbosacral transitional vertebrae and associated vertebral numeration, *Eur. Spine J.* **27**, 995–1004 (2018), doi:10.1007/s00586-018-5554-8.
- <sup>2</sup>T. A. Hutchins et al., ACR Appropriateness Criteria Low Back Pain: 2021 Update, *J. Am. Coll. Radiol.* **18**, S361–S379 (2021), doi:10.1016/j.jacr.2021.08.002.
- <sup>3</sup>American College of Radiology, ACR–ASNR–ASSR–SPR practice parameter for the performance of computed tomography (CT) of the spine, revised 2022 (Resolution 23), American College of Radiology, Reston, VA. <https://www.asnr.org/wp-content/uploads/2019/06/CT-Spine-1.pdf>, 2022.
- <sup>4</sup>American College of Radiology, ACR–ASNR–SABI–SSR practice parameter for the performance of magnetic resonance imaging (MRI) of the adult spine, revised 2023 (Resolution 6), American College of Radiology, Reston, VA. <https://www.asnr.org/wp-content/uploads/2019/06/MR-Adult-Spine.pdf>, 2023.
- <sup>5</sup>G. Konin and D. Walz, Lumbosacral transitional vertebrae: classification, imaging findings, and clinical relevance, *AJNR Am. J. Neuroradiol.* **31**, 1778–1786 (2010), doi:10.3174/ajnr.A2036.
- <sup>6</sup>Y. Deng et al., CTSpine1K: a large-scale dataset for spinal vertebrae segmentation in computed tomography, *Mach. Learn. Biomed. Imaging* **3**, 824–832 (2025), doi:10.59275/j.melba.2025-gf84.
- <sup>7</sup>P. Liu et al., Deep learning to segment pelvic bones: large-scale CT datasets and baseline models, *Int. J. Comput. Assist. Radiol. Surg.* **16**, 749–756 (2021), doi:10.1007/s11548-021-02363-8.
- <sup>8</sup>S. Nagata et al., Lumbosacral transitional vertebra on whole-spine CT: prevalence and association with rib abnormalities, *Skeletal Radiol.* **54**, 2169–2177 (2025), doi:10.1007/s00256-025-04952-z.
- <sup>9</sup>H. Möller et al., SPINEPS—automatic whole spine segmentation of T2-weighted MR images using a two-phase approach to multi-class semantic and instance segmentation, *Eur. Radiol.* **35**, 1178–1189 (2025), doi:10.1007/s00330-024-11155-y.
- <sup>10</sup>H. Möller et al., VERIDAH: Solving Enumeration Anomaly Aware Vertebra Labeling across Imaging Sequences, arXiv, 2026, arXiv:2601.14066. doi:10.48550/ARXIV.2601.14066.
- <sup>11</sup>Y. Otake et al., Automatic localization of vertebral levels in x-ray fluoroscopy using 3D-2D registration: a tool to reduce wrong-site surgery, *Phys. Med. Biol.* **57**, 5485–5508 (2012), doi:10.1088/0031-9155/57/17/5485.
- <sup>12</sup>S.-f. L. Lo et al., Automatic Localization of Target Vertebrae in Spine Surgery: Clinical Evaluation of the LevelCheck Registration Algorithm, *Spine (Phila. Pa. 1976)* **40**, E476–E483 (2015), doi:10.1097/BRS.0000000000000814.
- <sup>13</sup>T. De Silva et al., Utility of the LevelCheck Algorithm for Decision Support in Vertebral Localization, *Spine (Phila. Pa. 1976)* **41**, E1249–E1256 (2016), doi:10.1097/BRS.0000000000001589.
- <sup>14</sup>D. S. Kreiner et al., An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spinal stenosis (update), *Spine J.* **13**, 734–743 (2013), doi:10.1016/j.spinee.2012.11.059.
- <sup>15</sup>S. Sixta et al., Screening for thoracolumbar spinal injuries in blunt trauma: an Eastern Association for the Surgery of Trauma practice management guideline, *J. Trauma Acute Care Surg.* **73**, S326–S332 (2012), doi:10.1097/TA.0b013e31827559b8.
- <sup>16</sup>American College of Surgeons Committee on Trauma, ACS TQIP best practices guidelines in imaging, American College of Surgeons, Chicago. [https://www.facs.org/media/oxdjw5zj/imaging\\_guidelines.pdf](https://www.facs.org/media/oxdjw5zj/imaging_guidelines.pdf), 2018.
- <sup>17</sup>K. Smith et al., Data from CT COLONOGRAPHY (ACRIN 6664), The Cancer Imaging Archive, version 2 [dataset], 2015, doi:10.7937/K9/TCIA.2015.NWTESAY1.
- <sup>18</sup>K. Clark et al., The Cancer Imaging Archive (TCIA): maintaining and operating a public information repository, *J. Digit. Imaging* **26**, 1045–1057 (2013), doi:10.1007/s10278-013-9622-7.
- <sup>19</sup>A. M. Du Plessis, L. M. Greyling, and B. J. Page, Differentiation and classification of thoracolumbar transitional vertebrae, *J. Anat.* **232**, 850–856 (2018), doi:10.1111/joa.12781.
- <sup>20</sup>A. M. Poolman, Q. Wessels, A. V. Schoor, and N. Keough, Thoracolumbar transitional vertebrae: Quantitative differentiation and associated numeric variation in the vertebral column using skeletal remains, *J. Anat.* **243**, 311–318 (2023), doi:10.1111/joa.13865.
- <sup>21</sup>M. R. Zindrick et al., Analysis of the morphometric characteristics of the thoracic and lumbar pedicles, *Spine (Phila. Pa. 1976)* **12**, 160–166 (1987), doi:10.1097/00007632-198703000-00012.
- <sup>22</sup>M. M. Panjabi et al., Thoracic human vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **16**, 888–901 (1991), doi:10.1097/00007632-199108000-00006.
- <sup>23</sup>M. M. Panjabi et al., Human lumbar vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **17**, 299–306 (1992), doi:10.1097/00007632-199203000-00010.
- <sup>24</sup>J. L. Berry, J. M. Moran, W. S. Berg, and A. D. Steffee, A morphometric study of human lumbar and selected thoracic vertebrae, *Spine (Phila. Pa. 1976)* **12**, 362–367 (1987), doi:10.1097/00007632-198705000-00010.
- <sup>25</sup>R. J. Hughes and A. Saifuddin, Numbering of lumbosacral transitional vertebrae on MRI: role of the iliolumbar ligaments, *AJR Am. J. Roentgenol.* **187**, W59–W65 (2006), doi:10.2214/AJR.05.0415.
- <sup>26</sup>D. Schinz et al., A vertebral shape-based classification for labeling the thoracolumbar junction on CT imaging, *Eur. Spine J.* **35**, 2195–2203 (2026), doi:10.1007/s00586-025-09364-0.
- <sup>27</sup>H. Liebl et al., A computed tomography vertebral segmentation dataset with anatomical variations and multi-vendor scanner data, *Sci. Data* **8**, 284 (2021), doi:10.1038/s41597-021-01060-0.
- <sup>28</sup>A. Sekuboyina et al., VerSe: a vertebrae labelling and segmentation benchmark for multi-detector CT images, *Med. Image Anal.* **73**, 102166 (2021), doi:10.1016/j.media.2021.102166.
- <sup>29</sup>L. Jin et al., RibSeg v2: a large-scale benchmark for rib labeling and anatomical centerline extraction, *IEEE Trans. Med. Imaging* **43**, 570–581 (2024), doi:10.1109/TMI.2023.3313627.
- <sup>30</sup>J. Wasserthal et al., TotalSegmentator: robust segmentation of 104 anatomic structures in CT images, *Radiol. Artif. Intell.* **5**, e230024 (2023), doi:10.1148/ryai.230024.
- <sup>31</sup>H. Möller et al., Automated thoracolumbar stump rib detection and analysis in a large CT cohort, arXiv, 2025, arXiv:2505.05004. doi:10.48550/arXiv.2505.05004.
- <sup>32</sup>A. Schehr, J. Kim, and G. Schwing, OpenSpineConsortium: an open-source framework for medical student engagement in computational spine imaging research, *Cureus* (2026), doi:10.7759/cureus.112661.
- <sup>33</sup>F. Isensee, P. F. Jaeger, S. A. A. Kohl, J. Petersen, and K. H. Maier-Hein, nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation, *Nat. Methods* **18**, 203–211 (2021), doi:10.1038/s41592-020-01008-z.
- <sup>34</sup>R. Vialle, N. Levassor, L. Rillardon, A. Templier, W. Skalli, and P. Guigui, Radiographic analysis of the sagittal alignment and balance of the spine in asymptomatic subjects, *J. Bone Joint Surg. Am.* **87**, 260–267 (2005), doi:10.2106/JBJS.D.02043.
- <sup>35</sup>A. E. Castellvi, L. A. Goldstein, and D. P. Chan, Lumbosacral transitional vertebrae and their relationship with lumbar extradural defects, *Spine (Phila. Pa. 1976)* **9**, 493–495 (1984), doi:10.1097/00007632-198407000-00014.
- <sup>36</sup>P. J. Pickhardt, B. D. Pooler, T. Lauder, A. M. del Rio, R. J. Bruce, and N. Binkley, Opportunistic screening for osteoporosis using abdominal computed tomography scans obtained for other indications, *Ann. Intern. Med.* **158**, 588–595 (2013), doi:10.7326/0003-4819-158-8-201304160-00003.
- <sup>37</sup>M. G. Mody, A. Nourbakhsh, D. L. Stahl, M. Gibbs, M. Al-fawareh, and K. J. Garges, The Prevalence of Wrong Level Surgery Among Spine Surgeons, *Spine (Phila. Pa. 1976)* **33**, 194–198 (2008), doi:10.1097/BRS.0b013e31816043d1.
- <sup>38</sup>N. Epstein, A perspective on wrong level, wrong side, and wrong site spine surgery, *Surg. Neurol. Int.* **12**, 286 (2021), doi:10.25259/SNI.402.2021.
- <sup>39</sup>E. C. Benz, *Biomechanics of Spine Stabilization*, Thieme, New York, 3rd edition, 2015, ISBN 978-1-60406-924-2.
- <sup>40</sup>E. Beaulieu et al., A review of finite element analysis in spine surgery decision-making, *J. Clin. Med.* **15**, 2584 (2026), doi:10.3390/jcm15072584.