

**CTSpinoPelvic1K: spine, pelvis, ribs and femora in one coordinate frame, annotated
for lumbosacral transitional anatomy**

[Author names removed for double-anonymized review]

[Affiliations removed for double-anonymized review]

(Dated: 6 September 2026)

Purpose: Lumbar levels cannot be numbered reliably when a lumbosacral transitional vertebra (LSTV) is present. Counting down from C2, the reference standard for numeration¹, is impossible in an abdominal field of view, which leaves a specific ambiguity: four rib-free lumbar vertebrae may mean an L1 bearing a lumbar rib or an L5 assimilated to the sacrum, and six may mean a true sixth lumbar vertebra or a T12 whose ribs are aplastic². CTSpinoPelvic1K is built to test whether local morphology can resolve them without counting from the C2 anchor. It provides 802 CT records anchoring each lumbar level on the lowest rib-bearing vertebra above and the first sacral segment below, with radiologist-sourced spine *and* pelvic annotation on one frame, per-level ribs and femora. Classes for a sixth lumbar vertebra, a thirteenth thoracic vertebra, a separate first sacral segment and lumbar ribs cover the anomalies.

Acquisition and Validation Methods: Records pair TCIA CT COLONOGRAPHY imaging with CTSpine1K³ vertebral and CTPelvic1K⁴ pelvic labels on the acquisition with highest bone coverage, under a VerSe-native scheme. Validation: geometric invariants (802/802 pass); rib-vertebra incidence across 5,749 evaluable ribs (2 offsets, 0.035%); and agreement of spinopelvic measures with published values.

Data Format and Usage Notes: Gzipped NIfTI image/label pairs sharing one affine, canonicalized to PIR, with frozen patient-grouped LSTV-stratified five-fold splits and a reference loader; archived under a persistent identifier, <https://doi.org/10.5281/zenodo.22139642>.

Potential Applications: Testing whether a network can classify a vertebra from local features alone; updating textbook morphometry drawn from small cadaveric series; spinopelvic assessment at the lumbosacral interface; variant studies; opportunistic screening. *Limitations:* thoracic ground truth is field-of-view limited; postural angles supine; no held-out test set; the rib layer is a triaged-review pseudolabel; Castellvi grades are a two-reader consensus.

6 I. INTRODUCTION

7 The spine commonly carries seven cervical, twelve thoracic and five lumbar vertebrae, and
8 a vertebra is identified by where it falls in that sequence. Transitional variants sit at the two
9 ends of the lumbar column — the thoracolumbar transitional vertebra (TLTV) above, the
10 lumbosacral transitional vertebra (LSTV) below — and they disturb identification in two
11 ways at once. They change what the border vertebra and its ribs *look like*: a thoracolumbar
12 border vertebra may carry a full rib, a hypoplastic stump, or none at all, and a lumbosacral
13 one may be partly or completely assimilated to the sacrum, with an anteriorly wedged body,
14 a reduced disc beneath it and hypoplastic facet joints, or conversely separated from it with
15 a squared body, lumbar-type facets and a full-height disc². They also change *how many*
16 vertebrae each region contains — eleven or thirteen thoracic, four or six lumbar.

17 Either change alone would be tractable. Together they make it difficult to state defini-
18 tively what any vertebra at these borders *is*. Four rib-free vertebrae above the sacrum may
19 mean an L1 bearing a lumbar rib or an L5 assimilated to it; six may mean a true sixth
20 lumbar vertebra or a T12 whose ribs are aplastic. Each pair yields the same count, and the
21 morphology that would separate the two readings is not decisive to a human reader in every
22 case. Whether it nonetheless differs enough for a learned classifier to separate them, or at
23 least to assign each reading a probability, is the question this release is built to support.
24 Absent that, what remains is the sequence — conventionally established by counting down-
25 ward from C2, the reference standard for numeration¹, and unavailable in any spine-limited
26 acquisition. LSTV is common, reported between 4% and 30% depending on definition, and
27 wrong-level surgery is its most serious consequence.

28 Existing public collections cannot address this, and size is not the reason (Table I). Spine
29 datasets omit the pelvis; pelvic datasets do not number the vertebrae; and TotalSegmentator,
30 the whole-body scheme, has no class for a sixth lumbar vertebra or for a rib on a lumbar
31 vertebra. A six-lumbar spine therefore cannot be recorded in any of them, however good
32 the segmentation. VerSe^{5,6} makes the point most clearly (Table I): it deliberately enriched
33 for anatomical variants and graded them by Castellvi, yet left the graded vertebrae fused to
34 the sacrum unsegmented.

35 CTSpinoPelvic1K places the spine, pelvis, ribs and femora in one coordinate frame and
36 gives a class to each structure an enumeration anomaly produces: a sixth lumbar vertebra,

37 and a rib borne by a lumbar vertebra. A scheme without those classes must record such
38 anatomy as something it is not.

39 *Where the material is.* Everything described here is v8 of a single archived dataset,
40 deposited at <https://doi.org/10.5281/zenodo.22293878>. Citations should use <https://doi.org/10.5281/zenodo.22139642>, the permanent identifier for the dataset across all
41 of its versions, which resolves to whichever version is current; the version identifier above
42 pins the exact release measured in this manuscript. Documentation, the label scheme, the
43 reference loader and the per-record quality-control tables are distributed with the archive
44 and are described in Sec. III.

46 I.A. Vertebra labeling when the scan cannot settle the count

47 *The limitation, stated plainly.* No scan in this corpus contains C2 or the whole cervical
48 spine, so the conventional count cannot be performed. That is a property of abdominal
49 imaging rather than of the annotation, and it means a variant at the thoracolumbar junction
50 cannot be *resolved by counting* here at all: a thirteenth thoracic vertebra, a lumbar rib, and
51 a *stump rib* — a hypoplastic twelfth rib, one of the cardinal indicators of a thoracolumbar
52 transitional vertebra — produce overlapping appearances.

53 *What the release does instead.* No vertebral label is asserted where it cannot be supported.
54 Every quantity is expressed against the two landmarks present in essentially every abdominal
55 scan — the sacrum below and the lowest rib-bearing vertebra above — so the description is
56 positional rather than nominal: a vertebra by how many rib-free vertebrae separate it from
57 the sacrum, a lowest rib by its length as a fraction of the rib above it, a transverse process
58 by its span and its distance from the ala. None of these changes according to whether a
59 reader would have said L5 or S1, so cases that disagree about the name remain comparable.
60 Where the junction is in view the count is determined and the conventional name follows,
61 and the two descriptions agree. The positional description is, by construction, blind to the
62 ambiguity: an L1 bearing a lumbar rib and an L5 assimilated to the sacrum both leave
63 four rib-free vertebrae above the sacrum, and a true sixth lumbar vertebra and a T12 with
64 aplastic ribs both leave six. The count alone cannot separate the readings.

65 *But the vertebrae themselves are known to differ.* Counting is not the only route to an
66 identity. Level-specific morphometry is long established: pedicle width and height change

TABLE I Public CT collections at the lumbosacral junction. Black, class present; outlined, graded only to exclude it⁶.

Collection	Scans	Count	L6	Sacrum	S1	Pelvis	Ribs	L. rib	Femora	Grade
CTSpine1K ³	1,005									
CTPelvic1K ⁴	1,184									
VerSe ⁵	374									
RibSeg v2 ¹²	660									
TotalSegmentator ¹⁶	1,204									
CTSpinoPelvic1K	802									

systematically from the thoracic to the lumbar spine,⁷ vertebral body, endplate and canal dimensions differ by level,^{8–10} and the transverse process and the iliolumbar ligament mark the last lumbar vertebra independently of any count.^{2,11} If those differences hold at the boundary itself, the phenotype is decidable from the vertebra in front of the reader without any global count. On this corpus it does hold: separating T12 from L1 on shape alone, with each patient’s own size divided out and cross-validation grouped by case, gives an area under the curve of 0.990 and 97.3% accuracy over 1485 vertebrae. Size is deliberately removed, because a classifier given raw millimeters separates thoracic from lumbar immediately and has learned nothing that helps at the junction, where the two are nearly the same size.

That is reported as a property of the released measurements rather than as a method: the information needed to settle these variants is present locally.

II. ACQUISITION AND VALIDATION METHODS

II.A. Sources, and why a crosswalk was necessary

Both CTSpine1K³ and CTPelvic1K⁴ draw part of their imaging from the TCIA CT colonography collection^{13,14}, and the vertebral and pelvic annotations in both were produced under board-certified radiologist supervision. Neither released the mapping from an annotation to the specific CT series it was drawn on.

That mapping matters because every patient in CT COLONOGRAPHY was scanned *twice*, prone and supine, and an annotation carries a patient identifier and nothing finer.




FIG. 1 How the dataset was built. n counts scans, or corrections, at each step.

86 Turning a patient from supine to prone alters lumbar lordosis and the alignment of each
 87 segment against the next, so outlines drawn on one series are the wrong *shape* for the other
 88 and no rigid realignment recovers them. Such a mask looks competent in isolation and
 89 reveals itself only overlaid on the image. The crosswalk (Fig. 1) therefore resolves each
 90 annotation to a patient and then, within that patient's volumes, to the series whose bone
 91 *agrees* with it: candidates are thresholded at bone attenuation and scored by overlap with
 92 the mask.

93 II.B. Fused and split records

94 CTSpine1K and CTPelvic1K each annotated a patient independently, so their labels are
95 not guaranteed to come from the same CT series (Fig. 1). This turned out to matter for
96 351 patients, nearly half the cohort: the two collections had labeled different acquisitions of
97 the same person, one prone and one supine, with neither recording which. Because posture
98 changes spinal alignment, a spine label from one acquisition cannot simply be paired with a
99 pelvic label from the other. These 351 records are released but held out of any analysis that
100 assumes a single posture. Two labeled acquisitions of the same patient in different postures
101 are, instead, a resource for studying postural alignment, not a defect.

102 II.C. Densification, and why it is asymmetric

103 A corpus in which half the records lack a pelvis is not a spinopelvic dataset, so the missing
104 structures were pseudolabeled, asymmetrically (Fig. 1). Pelvis onto spine-only records is the
105 safe direction: the sacrum and innominate bones carry no enumeration to get wrong, and
106 quality is reported as held-out performance on the *pelvic_only* records, withheld from training
107 for that purpose. The reverse direction carries the whole problem this dataset addresses,
108 since a pseudolabeled spine must commit to a count and on a transitional vertebra commits
109 to the commoner reading; those 20 records were corrected by hand, tractable at that scale.

110 II.D. Ribs

111 No public rib annotation covers this cohort, and the two available automatic sources fail
112 in complementary ways.

113 Möller’s binary rib network¹⁵ reports high accuracy, and its authors show that stump ribs
114 can be classified from a partial field of view; that result concerns the morphological features
115 computed on a rib once segmented. On this corpus the segmentation itself failed on ribs
116 that were *partially outside* the field of view: a rib crossing the boundary of an abdominal
117 acquisition was liable to be missed altogether rather than segmented to the edge. This is
118 not a limitation reported by its authors: their evaluation excluded subjects in which the last
119 rib was not visible, and their failure analysis attributes errors to costal-process fusion, so it
120 is recorded here as an observation from this cohort. Since this is a colonography cohort, the

121 lower ribs are routinely clipped, and those are precisely the ribs the count depends on.

122 TotalSegmentator¹⁶ does segment clipped ribs and supplies the per-rib numbering that a
123 binary network cannot. Its characteristic failure is at the other end of the bone: the most
124 medial portion of the rib (roughly the last tenth to sixth approaching the costovertebral
125 joint) is frequently absent. A rib truncated at its medial end never reaches the vertebra
126 it articulates with, so the rib–vertebra incidence check (the quality-control gate on the rib
127 class labels, which matches each rib to the vertebra its number implies) has nothing to test.

128 The two were therefore unioned (Fig. 1): TotalSegmentator supplies numbering and the
129 medial-clipped cases, Möller supplies detection where its output is more complete, and the
130 union carries TotalSegmentator’s per-level identities.

131 The result is a pseudolabel whose review was triaged rather than exhaustive, which
132 is worth stating plainly because a layer described only as “human-corrected” invites the
133 reader to assume every bone was looked at. A label-based rule flagged every record in
134 which one rib bone carried more than one identifier, or in which a rib failed to reach the
135 vertebra it articulates with, since a single bone carrying two numbers is a numbering error by
136 construction, whatever the anatomy. The rule selected 152 records; 148 were reviewed and
137 corrected, and 4 were not and are identified in the release. Reviewers were medical students
138 working through a student research consortium¹⁷, trained against a written labeling protocol,
139 with every correction recorded against the annotator who made it. Records passing the rule
140 were not individually inspected, so the layer only guarantees freedom from the failure modes
141 the rule detects, no more. The residual rib–vertebra offsets reported in Sec. II.G are the
142 independent check on what the rule missed.

143 II.E. Label scheme and counting anchors

144 The scheme is VerSe-native: vertebrae keep their VerSe identifiers (C1–C7 = 1–7, T1–T12
145 = 8–19, L1–L6 = 20–25, sacrum 26), and every non-VerSe structure takes a fixed identifier
146 above that range — S1 carved from the sacrum (29), hips (30–31), femora (32–33), ribs
147 (34–45 left, 46–57 right), lumbar ribs (74–75) and surgical hardware (76–82). Identifiers
148 58–73 are unassigned: a soft-tissue block once reserved there was never populated and is
149 retired.

150 Two anchors make the count decidable, and Fig. 2 shows both: the lowest rib-bearing

151 vertebra above, **S1** carved from the sacrum below. Each earns its place by what it supports
152 rather than by being an endpoint — S1 supplies the sacral endplate landmark that sacral
153 slope and pelvic incidence are measured from, and with both brackets present L5 and L6
154 are distinguishable where a spine-limited view alone would not be.

155 *How S1 is cut.* The sacrum’s outer boundary is the source annotation and is never altered;
156 S1 is the cranial part of that same bone, separated by a plane. The plane’s normal is the
157 sacrum’s own cranio-caudal axis, made orthogonal to a left–right axis measured directly
158 rather than assumed, so as to remove the patient’s rotation within the scanner. That
159 rotation is not negligible: median 4.5°, maximum 16.1°, and 508 of 801 records above 3°.
160 The level of the cut preserves the sub-division volume of the preceding release, so the plane
161 changes the shape and orientation of the S1/S2 boundary and not how much of the sacrum
162 is called S1. Per-record geometry ships with the archive.

163 Neither anchor is novel on its own: TotalSegmentator¹⁶ carries a separate S1 class and
164 per-level ribs, and VerSe⁵ carries L6. Because this release uses the VerSe identifiers, its L6
165 *is* VerSe’s, interoperable rather than local to this work. L6 is retained not for novelty but
166 because a scheme without it cannot record a six-lumbar spine at all, and must renumber the
167 column or drop a level to fit.

168 **A rib on a lumbar body receives its own class**, and this is the one class here with no
169 published counterpart. A lumbar rib and a stump rib are the same object under two counts,
170 so assigning it a number would commit the label to one reading of an enumeration anomaly.
171 A scheme that numbers every rib 1–12 has nowhere to put a thirteenth: the annotator must
172 either call it rib 12 — asserting the vertebra beneath it is thoracic, the very question at
173 issue — or discard it. TotalSegmentator carries ribs 1–12 per side and no such class, and
174 VerSe’s T13 is a vertebra rather than a rib. 16 released records carry a lumbar rib and 18
175 carry an L6.

176 Figure 2 shows why the interval is the primitive and the label is derived. Among the
177 records with four rib-free vertebrae, some reach that count through a lumbar rib and some
178 without one; the two are indistinguishable by the count alone and are distinguished here
179 only because rib presence is itself labeled.

180 Sixteen records carry a lumbar rib in the released volumes: thirteen bilateral and three
181 unilateral, all three on the right. That laterality is reported because it is what the release
182 contains and not as a finding — sixteen records cannot support a statement about side




FIG. 2 The two anchors, from the released labels. Red, lowest rib-bearing vertebra and its rib; blue, S1; yellow, the rib-free vertebrae between. (b) and (c) both count four.

183 preference, and none is made.

184 Eighteen records carry an L6. Fourteen are labeled lumbarization, two sacralization,
 185 one semi-sacralization and one normal, which is worth stating because it shows the two
 186 annotation axes disagreeing in the open: a record can carry six lumbar-type vertebrae and
 187 still be read as a sacralization, or as unremarkable, depending on where the reader started
 188 the count.

189 Both counts are derived from the released label volumes, and so are the manifest fields
 190 that report them. That is a correction rather than a design note: through v5 the `has_l6`
 191 flag was true for exactly one record, which contains no L6, and false for all eighteen that
 192 do, while `has_lumbar_rib` was false everywhere and `n_lumbar_labels` read zero in 799 of
 193 802 records. A reader selecting the six-lumbar cases on those fields would have received one
 194 record that is not one, and a reader selecting lumbar ribs an empty set, with nothing to
 195 indicate either. The fields are rebuilt for this release by counting identifiers 25 and 74–75 in
 196 each volume, and `lumbar_rib_side` is added alongside them.

197 II.F. Computational tools

198 *Access.* Every processing step described above is performed by code released under an
 199 open-source licence and archived with the dataset; no step depends on software that is
 200 not publicly obtainable. The release archive carries the reference loader, the label-scheme
 201 definition and the quality-control scripts that generated the tables in this manuscript.

202 *Versions and parameters.* Segmentation of femora, the sacral sub-division and the per-

level rib numbering used TotalSegmentator¹⁶ (`total` task, release 2.x) on a single graphics processing unit. The binary rib network is Möller’s¹⁵ released weights, applied through the nnU-Net v2 framework¹⁸. The pelvic completion for records whose pelvis was absent used a five-fold nnU-Net v2 ensemble, applied out-of-fold: each record is completed by the fold that never trained on it, so no record is completed by a model that has seen it. The five fold checkpoints, with their nnU-Net plans and dataset descriptor, are released alongside the data (<https://huggingface.co/anonymous-mlhc/spinopelvic-seg-checkpoints>). Image handling throughout used NiBabel and SciPy; no image is resampled or reoriented when a label is written, so every label shares its image’s grid and affine exactly.

Generative artificial intelligence. Assistance from large language models was used for manuscript preparation — drafting and editing of text, and generation of analysis and figure code — under author review. It was not used to generate, annotate or interpret any imaging data, and no scientific claim in this manuscript originates from it. All numbers reported here are computed by the released code from the released volumes.

II.G. Validation

Validation is layered, and the ordering matters: measurements computed on a corpus that has not established internal consistency do not look broken, they look finished.

Geometric invariants. Every case is checked for label geometry agreeing with its CT in shape, affine and voxel spacing; for identifiers within the published scheme; for a non-empty label; and for sided structures falling on the correct sides, with the left–right axis read from the affine rather than assumed. The check exists because a renumbering pass once wrote a reoriented array under a canonical affine and transposed a label away from its image, a failure invisible downstream.

The check compares *each sided pair separately*, and both simpler alternatives miss cases it catches. Testing the ribs alone passed all 802 records while four of them carried `left hip` on the patient’s right. Pooling all sided structures into one test recovered three of those four and missed the fourth, because hips and femora outweigh the ribs by voxel count, so a large correctly-sided structure masks a smaller transposed one. A gate that passes everything is not evidence of a clean corpus until it has been shown capable of failing.

Rib–vertebra incidence. Each rib is matched to its nearest vertebra and compared against

the vertebra its number implies. Of 11,548 ribs, 5,788 are not evaluable because the expected vertebra lies outside the field of view and 11 have no vertebra within the anchor distance; of the 5,749 evaluable, 5,747 match, two (0.035%) are offset by one level, and no case is misnumbered. The denominator is stated because quoting the two residual offsets against all 11,548 ribs would halve the rate by counting ribs the check never examined — and because more than half the ribs in a screening abdominal CT being unevaluable is itself a fact about the corpus worth reporting. Both offsets are field-of-view truncations on individually characterized cases, one at +1 level and one at −1, and are reported rather than suppressed: a threshold tuned until the count reaches zero yields a cleaner number and a less informative one.

The same check carries an invariant that reads as a null result. Of the 5,749 evaluable ribs, *zero* are assigned to a lumbar vertebra — not because no lumbar ribs exist, since 16 records carry one, but because such a rib takes class 74 or 75 and never enters the numbered set the test examines. A numbered rib landing on a lumbar vertebra would be a counting error of exactly the kind this dataset exists to expose.

Agreement with published values. Derived spinopelvic measures are compared against Vialle *et al.*¹⁹ (Table II). Pelvic incidence is the strongest of these checks because it is a morphological property of the pelvis rather than a posture, and so is directly comparable to a standing reference cohort.

II.H. Castellvi typing

Every record whose vertebral count departs from five carries a Castellvi grade²⁰ in the released manifest: 33 cases, typed I–IV with the *a/b* unilateral–bilateral qualifier. This is released as an annotation layer because the grade and the count are *different axes* and the dataset is built to show that they are: grade IIb occurs here at rib-free counts of four, five and six alike, and seven of the 33 graded cases carry a perfectly normal count of five. A corpus that recorded only the count would describe those seven as unremarkable.

Two radiology resident physicians graded every case independently and resolved their six disagreements by consensus; both reads and the consensus are in the manifest. The distinction the classification turns on — bony fusion against articulation without it² — is the one carrying clinical weight, and it is where the disagreements lay: four of the six were

TABLE II Derived measures against a standing reference ($n = 260$)¹⁹; this cohort is supine.

Measure	This dataset (supine)	Reference (standing)
Pelvic incidence	54.7°	54.7 ± 10.6
Pelvic tilt	17.3°	13 ± 6
Sacral slope	36.9°	41.0 ± 8.4
Lumbar lordosis (supine)	52.7°	43 ± 11.2 (standing)
PI – LL mismatch	2.6°	centered near 0

III against II.

II.I. Surgical hardware, and why a threshold is not enough

Metal is trivial to detect and hard to interpret, and the difference matters here because **an iatrogenic fusion is indistinguishable from a congenital one to a distance measurement**: a cage-bridged interspace reads as “no gap” exactly as a fused transitional vertebra does. An unrecorded instrumented case is therefore a false positive on the central measure.

II.I.0.a. Detection over-calls by a factor of five. A 2500 HU threshold inside a shell around the labeled skeleton flags 52 of the 802 records. One of the two radiology resident physicians read all 52 and confirmed 11, rejecting 41 (79%) as contrast, calcification and reconstruction artefact. Attenuation does not separate the two groups (both saturate); site and volume do, every confirmed implant at 2,586 mm³ or above and every rejection at 1,768 mm³ or below. Instrumentation prevalence is 1.4%, not the 6.5% the threshold alone reports.

II.I.0.b. The subtype block had to be extended. Identifiers 76–79 name spinal instrumentation (generic, cage, screw-and-rod, plate) and cannot describe what this cohort holds, so **80 arthroplasty**, **81 sacroiliac screw** and **82 osteosynthesis** were added: osteosynthesis holds parts of one bone together where arthroplasty replaces a joint, and a shape rule calls a femoral stem a rod. The confirmed cases are 8 arthroplasties, one osteosynthesis, one sacroiliac fixation and one pair of interbody cages (Fig. 3).

II.I.0.c. Metal outranks bone, and that is a subtraction. In every confirmed case the implant was already inside a bone label, so naming it reclaims 1,538,852 voxels rather than




FIG. 3 Instrumentation in the release, one exemplar each, drawn through bone. (a) Hip arthroplasty, 80, $n = 8$. (b) Femoral-neck screws, 82, $n = 1$. (c) Sacroiliac screws, 81, $n = 1$. (d) Interbody cages, 77, $n = 1$. Bar, 5 cm.

adding them. Pelvic incidence and tilt are measured from the femoral head center, and in 8 of these cases that head is an implant: any spinopelvic parameter previously computed from them was measured on metal.

III. DATA FORMAT AND USAGE NOTES

Each record is a gzipped NIfTI image/label pair sharing an identical 4×4 affine, canonicalized to PIR, requiring no resampling. Masks are NIfTI rather than DICOM: the policy prefers DICOM where applicable, and for volumetric segmentation masks NIfTI is what the downstream tooling expects.

Splits are patient-grouped and LSTV-stratified five-fold, frozen and shipped with the data as `splits_5fold.json`. The five validation folds partition all 802 records, so the release provides cross-validation and *not* a held-out test set. Users reporting a single held-out number should carve one out and state which records it holds, because the folds shipped here will not do that job.

Stratification is on the transitional subtype rather than on a binary LSTV flag, and it enforces a minimum of each rare subtype in every validation fold: each fold’s validation set carries three sacralization and three lumbarization records, so no fold is evaluated on a cohort missing the class the dataset exists for. With 15 lumbarization and 15 sacralization records across 802, that is the practical limit of what stratification can guarantee, and it is why a fold-level metric on the rare classes carries an error bar far wider than its own decimal places.

TABLE III Data types and metadata, and the records populating each, from the released manifest.

Field	Records	Available
Identifier, image path, label path, configuration, match type	802	100.0%
Patient position, tube potential, section thickness, kernel	802	100.0%
Transitional label and class	802	100.0%
Spine and pelvic annotation origin	802	100.0%
Instrumentation and partial-annotation flags	802	100.0%
Image/label alignment check	802	100.0%
Spine series identifier; bone-fraction check	782	97.5%
Scanner manufacturer and model	768	95.8%
Sex	749	93.4%
Age	709	88.4%
Pelvic series identifier	713	88.9%
Castellvi grade (consensus of two readers)	33	4.1%

Table III lists the key data types and metadata against the fraction of records for which each is populated.

The archive of record is Zenodo (<https://doi.org/10.5281/zenodo.22139642>). Code and documentation are on GitHub at the tagged release commit; any model-hub copy is a convenience mirror, not the archive.

IV. DESCRIPTIVE ANALYSIS

The cohort is a colorectal cancer screening population of 802 records. 738 are recorded female (393) or male (345); 11 carry DICOM’s *other* value, a recorded value and not a missing one, and 53 carry no sex field. Age is present for 709 (median 59, range 50–89). All 802 are counted in totals, and records outside the female/male dichotomy are excluded from sex-stratified measures (a limitation of those measures rather than of the cohort), stated because folding eleven people into “unrecorded” would misdescribe the source. The lower age bound is the screening guideline rather than a selection applied here, and it is why the measures below sit closer to a reference range than a surgical series would.




FIG. 4 Count-free measures. (a) Vertebrae between the lowest rib and the sacrum. (b) Lowest rib length over the rib above. (c) Lowest lumbar transverse span against its gap to the ala; transitional labels marked.




FIG. 5 Spinopelvic measures against standing reference bands (mean $\pm$ SD)¹⁹.

319 *Count-free measures* (Fig. 4). Five rib-free vertebrae above the sacrum is typical; four
 320 and six are where transitional anatomy sits, and which of the two is present does not by
 321 itself determine what to call it. The lowest-rib length ratio is bimodal.

322 *Level gradients and spinopelvic measures* (Fig. 5). Vertebral dimensions increase caudally
 323 in step with the load each level carries. Pelvic incidence does not vary with age, while pelvic
 324 tilt increases — the descriptive signature of a pelvis retroverting as a spine ages.

325 *Opportunistic measures*. Every scan carries a vertebral bone density measurement at no
 326 additional dose.²¹ L1 trabecular attenuation is reported at the published standard site.

327 V. POTENTIAL APPLICATIONS AND LIMITATIONS

328 The principal use is the *spinopelvic interface*. Spine and pelvis carry radiologist-sourced
329 annotation on one coordinate frame in 802 records, making sacral morphology measurable
330 against the lumbar column rather than in isolation — the sacral endplate and its slope, the
331 alar corridors constraining S2-alar-iliac and iliac screw trajectories, and the L5–S1 geometry
332 a lumbosacral fusion must cross. Pelvic incidence is derived from the segmented sacrum and
333 femoral heads rather than from landmarks placed on a radiograph, so it is reproducible from
334 the labels themselves.

335 The transitional layer serves that use rather than competing with it: no trajectory can be
336 planned across a junction whose segments cannot be named. Level-numbering benchmarks,
337 variant studies and opportunistic screening follow from the same annotation.

338 V.A. Three uses the released measurements support

339 *Naming a level from its own geometry.* Wrong-level spine surgery runs at roughly one
340 in 3,110 spinal procedures, and the cause most often cited is the transitional anatomy this
341 cohort was assembled around.²² The failure is structural rather than careless: the count
342 is ambiguous exactly where the variant sits. Per-level body height, canal width, endplate
343 width, transverse-process span and wedge ratio are released for T11–L5 across all 802 records,
344 and because the labels were assigned against the twelfth rib rather than by enumeration, a
345 model trained on them is not learning the convention that fails.

346 *Reference morphometry with its spread.* The values a surgeon consults for vertebral
347 body height, canal dimensions and pedicle width come from cadaveric series of a few dozen
348 specimens, plotted as single values with no indication of spread.¹⁰ They cannot tell a reader
349 whether the patient in front of them is unusual, because they never show what usual looks
350 like. The same measures are given here from 802 records with the distribution attached
351 (Fig. 6), reproducing two properties of the classical figures — body height crosses over,
352 dorsal exceeding ventral at T11–T12 and reversing by L4–L5, and pedicle width widens
353 steeply into the lower lumbar spine — and adding the width of each distribution, which is
354 the part needed to judge an individual. A larger n sharpens a distribution without changing
355 the population it came from, and this one is a screening cohort aged 50 and over.




FIG. 6 Morphometry by level: median, interquartile range, 5th–95th percentile, n at right. (a) Body height. (b) Endplate width. (c) Canal. (d) Pedicle width. (e) Disc height. (f) Trabecular attenuation.

Patient-specific models. Planning is moving toward patient-specific biomechanical models rather than population norms.²³ Such a model needs spine, pelvis and femoral heads in one frame, which is this release’s construction. In 351 patients the annotation sits on two acquisitions in different positions, giving a within-patient change in alignment against which a predicted postural response can be checked.

Its limitations are stated at the level of detail a reader would need to catch them independently.

Coverage. Thoracic ground truth is field-of-view limited (Fig. 7) and does not extend to T1. Postural angles are supine; pelvic incidence is not postural and needs no such caveat. The cohort is a colorectal screening population aged 50 and over, so its distributions should not be read as representative of a surgical one.

Declared but empty, and confirmed so. The partial-annotation sentinel (255) is declared in the scheme and occurs in no record — established by counting identifiers across all 802 released volumes rather than asserted. It is retained because the protocol is part of the




FIG. 7 Records carrying each vertebral level, cranial to caudal. Counts per identifier in Table S1.

370 scheme and a later release may ship a partially traced record. The hardware identifiers were
 371 in this position until v6 and are no longer: 76–82 are populated in the eleven instrumented
 372 records described in Sec. II.I.

373 *Strength of the labels varies by structure, and the release does not average over that.*
 374 Vertebral labels derive from radiologist-supervised source annotations, corrected where those
 375 sources were wrong. Pelvic labels on records that lacked one are pseudolabeled. The rib
 376 layer is a pseudolabel whose human review was triaged by an automated rule rather than
 377 exhaustive, so its guarantee is the absence of the failure modes that rule detects. Transitional
 378 labels derive from two independent sources and are not uniformly adjudicated — which is
 379 precisely why the measures reported here are count-free — and the Castellvi grades are
 380 a consensus of two radiology resident physicians. The sacral sub-division inherits its level
 381 from an automatic method, and in 100 records that level places S1 above half the sacrum
 382 or below fifteen percent of it, which no first sacral segment is; those records are flagged in
 383 the released quality-control table and should be filtered before any measurement depending
 384 on the sacral endplate.

385 *Structures are not universally present.* The same census shows the release is not uniformly
 386 complete. Sacrum, hips and femora are present in all 802 records. One record carries no

387 S1, one lacks T12, and nine lack an L5 identifier — in every case because the structure
388 lies outside the field of view or, for L5, because the lowest lumbar segment is labeled L6 or
389 incorporated into the sacrum. Users filtering on the presence of the caudal anchor should
390 filter explicitly rather than assume it.

391 VI. DISCUSSION

392 *Comparison with related material.* Against the collections in Table I, the contribution
393 is the crosswalk placing spine and pelvis on the same series, together with classes for the
394 anatomy an enumeration anomaly actually produces. VerSe⁵ comes closest in intent: it
395 also enriched its cohort for transitional anatomy and graded each case by Castellvi. The
396 difference is what a graded case leaves behind. VerSe names the anomaly but does not
397 segment the sacrum, so the fused or articulating structure the grade describes has no mask
398 to go with it. This release segments that structure directly, as L6 or as a separately carved
399 S1, so a user has the anatomy itself to measure rather than only a grade that describes it.
400 TotalSegmentator¹⁶ invites a different comparison, since it segments far more of the body
401 than any single source used here. But its label scheme simply has no class for a sixth lumbar
402 vertebra or for a rib on a lumbar body. A six-lumbar spine cannot be recorded in it: the
403 column must be renumbered or a level dropped to fit. What this release adds, then, is not
404 more anatomy per scan (TotalSegmentator already covers more), but the specific classes an
405 enumeration anomaly needs in order to be recorded accurately.

406 *Limitations.* The cohort comes from one acquisition protocol of a colorectal-screening
407 population aged 50 and over, so how well it generalizes to a pathologic population is untested.
408 The rib layer’s human review was triaged by an automated rule rather than performed on
409 every bone, so it guarantees only the absence of the failure modes that rule detects.

410 *Future directions.* The version history shows the scheme being extended more than once:
411 femora and the S1 anchor, then per-level ribs, then lumbar ribs and surgical hardware, were
412 each folded into the existing release rather than requiring a new dataset. The same path is
413 open going forward: imaging from beyond this cohort’s single protocol, and the C2 anchor
414 this acquisition cannot supply, are the natural next additions.

415 **VII. CONCLUSION**

416 CTSpinoPelvic1K places spine, pelvis, per-level ribs and femora on one coordinate frame
417 in 802 records, and gives the anatomy that makes lumbar numbering ambiguous a class of
418 its own rather than recording it as something it is not. Because both counting anchors are
419 explicit, a level can be identified from a field of view that cannot support the conventional
420 count downward from C2, which is every record here.

421 **DATA AVAILABILITY**

422 The release described here is v8, permanently archived at [https://doi.org/10.5281/](https://doi.org/10.5281/zenodo.22293878)
423 [zenodo.22293878](https://doi.org/10.5281/zenodo.22293878); the concept DOI <https://doi.org/10.5281/zenodo.22139642> resolves
424 to the current version. The archive carries the loader, the label scheme, the quality-control
425 tables and the analysis code under an open-source licence. The repository URL identifies
426 the authors; it is on the title page and will be restored here for the camera-ready version.

427 **ACKNOWLEDGEMENTS**

428 Acknowledgements are given on the title page, per the journal’s double-anonymized re-
429 view policy. This work received no funding.

430 **CONFLICT OF INTEREST**

431 The authors have no relevant conflicts of interest to disclose.

432 **ETHICS**

433 This work uses publicly available, de-identified imaging and annotations derived from it.
434 The authors’ institutional review board determined that it is not human participant research
435 under 45 CFR 46 and requires no oversight.

REFERENCES

- ¹J. Lian, N. Levine, and W. Cho, A review of lumbosacral transitional vertebrae and associated vertebral numeration, *Eur. Spine J.* **27**, 995–1004 (2018), doi:10.1007/s00586-018-5554-8.
- ²G. P. Konin and D. M. Walz, Lumbosacral transitional vertebrae: classification, imaging findings, and clinical relevance, *AJNR Am. J. Neuroradiol.* **31**, 1778–1786 (2010), doi:10.3174/ajnr.A2036.
- ³Y. Deng et al., CTSpine1K: a large-scale dataset for spinal vertebrae segmentation in computed tomography, *Mach. Learn. Biomed. Imaging* **3**, 824–832 (2025), doi:10.59275/j.melba.2025-gf84.
- ⁴P. Liu et al., Deep learning to segment pelvic bones: large-scale CT datasets and baseline models, *Int. J. Comput. Assist. Radiol. Surg.* **16**, 749–756 (2021), doi:10.1007/s11548-021-02363-8.
- ⁵A. Sekuboyina et al., VerSe: a vertebrae labelling and segmentation benchmark for multi-detector CT images, *Med. Image Anal.* **73**, 102166 (2021), doi:10.1016/j.media.2021.102166.
- ⁶H. Liebl et al., A computed tomography vertebral segmentation dataset with anatomical variations and multi-vendor scanner data, *Sci. Data* **8**, 284 (2021), doi:10.1038/s41597-021-01060-0.
- ⁷M. R. Zindrick et al., Analysis of the morphometric characteristics of the thoracic and lumbar pedicles, *Spine (Phila. Pa. 1976)* **12**, 160–166 (1987), doi:10.1097/00007632-198703000-00012.
- ⁸M. M. Panjabi et al., Thoracic human vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **16**, 888–901 (1991), doi:10.1097/00007632-199108000-00006.
- ⁹M. M. Panjabi et al., Human lumbar vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **17**, 299–306 (1992), doi:10.1097/00007632-199203000-00010.
- ¹⁰E. C. Benzel, *Biomechanics of Spine Stabilization*, Thieme, New York, 3rd edition, 2015.
- ¹¹R. J. Hughes and A. Saifuddin, Numbering of lumbosacral transitional vertebrae on MRI: role of the iliolumbar ligaments, *AJR Am. J. Roentgenol.* **187**, W59–W65 (2006), doi:10.2214/AJR.05.0415.

¹²L. Jin et al., RibSeg v2: a large-scale benchmark for rib labeling and anatomical centerline extraction, *IEEE Trans. Med. Imaging* **43**, 570–581 (2024), doi:10.1109/TMI.2023.3313627.

¹³K. Smith et al., Data from CT COLONOGRAPHY, The Cancer Imaging Archive, 2015, doi:10.7937/K9/TCIA.2015.NWTESAY1.

¹⁴K. Clark et al., The Cancer Imaging Archive (TCIA): maintaining and operating a public information repository, *J. Digit. Imaging* **26**, 1045–1057 (2013), doi:10.1007/s10278-013-9622-7.

¹⁵H. Möller et al., Automated thoracolumbar stump rib detection and analysis in a large CT cohort, 2025, arXiv:2505.05004. doi:10.48550/arXiv.2505.05004.

¹⁶J. Wasserthal et al., TotalSegmentator: robust segmentation of 104 anatomic structures in CT images, *Radiol. Artif. Intell.* **5**, e230024 (2023), doi:10.1148/ryai.230024.

¹⁷[Reference to the authors’ own work, removed for double-anonymized review.]

¹⁸F. Isensee, P. F. Jaeger, S. A. A. Kohl, J. Petersen, and K. H. Maier-Hein, nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation, *Nat. Methods* **18**, 203–211 (2021), doi:10.1038/s41592-020-01008-z.

¹⁹R. Vialle, N. Levassor, L. Rillardon, A. Templier, W. Skalli, and P. Guigui, Radiographic analysis of the sagittal alignment and balance of the spine in asymptomatic subjects, *J. Bone Joint Surg. Am.* **87**, 260–267 (2005), doi:10.2106/JBJS.D.02043.

²⁰A. E. Castellvi, L. A. Goldstein, and D. P. K. Chan, Lumbosacral transitional vertebrae and their relationship with lumbar extradural defects, *Spine (Phila. Pa. 1976)* **9**, 493–495 (1984), doi:10.1097/00007632-198407000-00014.

²¹P. J. Pickhardt, B. D. Pooler, T. Lauder, A. M. del Rio, R. J. Bruce, and N. Binkley, Opportunistic screening for osteoporosis using abdominal computed tomography scans obtained for other indications, *Ann. Intern. Med.* **158**, 588–595 (2013), doi:10.7326/0003-4819-158-8-201304160-00003.

²²N. E. Epstein, A perspective on wrong level, wrong side, and wrong site spine surgery, *Surg. Neurol. Int.* **12**, 286 (2021), doi:10.25259/SNI.402.2021.

²³E. Beaulieu et al., A review of finite element analysis in spine surgery decision-making, *J. Clin. Med.* **15**, 2584 (2026), doi:10.3390/jcm15072584.