

**CTSpinoPelvic1K: spine, pelvis, ribs and femora in one coordinate frame, annotated
for lumbosacral transitional anatomy**

[Author names removed for double-anonymized review]

[Affiliations removed for double-anonymized review]

(Dated: 6 September 2026)

Purpose: For transitional anatomy the gold standard for numeration remains whole-spine imaging and counting caudally from C2¹. Abdominal CT does not contain C2, and where a lumbosacral transitional vertebra (LSTV) alters the count the local anatomy is ambiguous: four rib-free lumbar vertebrae may mean an L1 bearing a lumbar rib or an L5 assimilated to the sacrum, and six may mean a sixth lumbar vertebra, a T12 whose ribs are aplastic, or an S1 lumbarized from the sacrum². CT-SpinoPelvic1K is built so that this can be asked of the data: whether local morphology resolves them without counting from C2. It provides 802 CT records anchoring each lumbar level on the lowest rib-bearing vertebra above and the first sacral segment below, with radiologist-sourced spine *and* pelvic annotation on one frame, per-level ribs and femora. Classes for a sixth lumbar vertebra, a thirteenth thoracic vertebra, a separate first sacral segment and lumbar ribs cover the anomalies.

Acquisition and Validation Methods: Records pair TCIA CT COLONOGRAPHY imaging with CTSpine1K³ vertebral and CTPelvic1K⁴ pelvic labels on the acquisition with highest bone coverage, under a VerSe-native scheme. Validation: geometric invariants (802/802 pass); rib-vertebra incidence across 5,749 evaluable ribs (2 offsets, 0.035%); and agreement of spinopelvic measures with published values.

Data Format and Usage Notes: Gzipped NIfTI image/label pairs sharing one affine, canonicalized to PIR, with frozen patient-grouped LSTV-stratified five-fold splits and a reference loader; archived under a persistent identifier, <https://doi.org/10.5281/zenodo.22139642>.

Potential Applications: Testing whether a network can classify a vertebra from local features alone; updating textbook morphometry drawn from small cadaveric series; spinopelvic assessment; variant studies; opportunistic screening. *Limitations:* thoracic ground truth is field-of-view limited; postural angles supine; no held-out test set; the rib layer is a triaged-review pseudolabel; Castellvi grades are a two-reader consensus.

6 I. INTRODUCTION

7 The spine commonly carries seven cervical, twelve thoracic and five lumbar vertebrae, and
8 a vertebra is identified by where it falls in that sequence. Transitional variants sit at the
9 two ends of the lumbar column, the thoracolumbar transitional vertebra (TLTV) above and
10 the lumbosacral transitional vertebra (LSTV) below, and they disturb identification in two
11 ways at once. They change what the border vertebra and its ribs *look like*: a thoracolumbar
12 border vertebra may carry a full rib, a hypoplastic stump, or none at all, and a lumbosacral
13 one may be partly or completely assimilated to the sacrum (*sacralization*), with an anteriorly
14 wedged body, a reduced disc beneath it and hypoplastic facet joints, or conversely separated
15 from it (*lumbarization*), with a squared body, lumbar-type facets and a full-height disc².
16 They also change *how many* vertebrae each region contains: eleven or thirteen thoracic,
17 four or six lumbar.

18 Either change alone would be tractable. Together they make it difficult to state defini-
19 tively what any vertebra at these borders *is*. Four rib-free vertebrae above the sacrum may
20 mean an L1 bearing a lumbar rib or an L5 assimilated to it; six may mean a true sixth
21 lumbar vertebra, a T12 whose ribs are aplastic, or an S1 lumbarized away from the sacrum.
22 Each set yields the same count, and the morphology that would separate the readings is not
23 decisive to a human reader in every case. Whether it differs enough for a learned classifier
24 to separate them, or at least to assign each reading a probability, is the question this release
25 is built to support. Absent that, what remains is the count itself: the gold standard for
26 numeration in transitional anatomy is whole-spine imaging, counting caudally from C2¹,
27 which fixes how many vertebrae and how many rib pairs the column holds, and no spine-
28 limited acquisition can supply either count. LSTV is common, reported between 4% and
29 30% depending on definition, and wrong-level surgery is its most serious consequence.

30 Existing public collections cannot address this, and size is not the reason (Table I). Spine
31 datasets omit the pelvis; pelvic datasets do not number the vertebrae; and TotalSegmentator,
32 the whole-body scheme, has no class for a sixth lumbar vertebra or for a rib on a lumbar
33 vertebra. A six-lumbar spine therefore cannot be recorded in any of them, however good the
34 segmentation, and a segmenter trained on them inherits the gap: faced with an enumeration
35 anomaly it must shift the whole column by a level or absorb a vertebra into its neighbor.
36 The size of that problem has now been measured on CT, where prior labeling methods

37 assigned every vertebra correctly in 77% of subjects and an anomaly-aware method raised
38 that to 99%⁵. VerSe^{6,7} makes the point most clearly (Table I): it deliberately enriched for
39 anatomical variants and graded them by Castellvi, yet left the graded vertebrae fused to the
40 sacrum unsegmented.

41 The intraoperative side of the problem has prior art of its own. LevelCheck registers
42 the intraoperative radiograph to the preoperative CT and projects the CT’s vertebral labels
43 onto it, localizing the target level in a limited field of view to within a few percent of cases
44 in clinical use^{8–10}. The labels it projects are placed by hand on the CT, by a skilled operator
45 and verified by the surgeon⁸: the method assumes a correctly labeled CT and does not
46 supply one. Automating that step, and getting it right precisely where the anatomy is
47 anomalous, is what a dataset carrying the classes those anomalies produce is for.

48 CTSpinoPelvic1K places the spine, pelvis, ribs and femora in one coordinate frame and
49 gives a class to each structure an enumeration anomaly produces: a sixth lumbar verte-
50 bra, and a rib borne by a lumbar vertebra. A scheme without those classes lacks a class
51 vocabulary expressive enough to describe anatomic anomalies.

52 The project began as a surgeon’s question: whether a vertebra can be named from the
53 anatomy in front of the reader, on the abdominal and spine-limited images a surgeon actually
54 has, in the patients where counting fails. The cohort suits that question. Every record comes
55 from one prospective trial protocol, ACRIN 6664^{11,12}: 802 patients aged 50 and over, scanned
56 supine and prone on multidetector CT at 15 centers, on five scanner vendors and nine models,
57 with manufacturer, model and reconstruction kernel recorded per record. To our knowledge
58 it is the largest spine-and-pelvis annotated CT cohort acquired under a single protocol; the
59 comparators pool several collections (CTSpine1K four, CTPelvic1K seven), are multi-site
60 by design (VerSe), or are routine clinical scans under many protocols (TotalSegmentator).
61 With the protocol held fixed, scanner effects on a model become measurable rather than
62 confounded. The price is the field of view: abdominal, so the thoracic spine enters from
63 below and C2 is never present (Sec. V).

64 *Where the material is.* Everything described here is v8 of a single archived dataset,
65 deposited at <https://doi.org/10.5281/zenodo.22293878>. Citations should use [https:](https://doi.org/10.5281/zenodo.22139642)
66 [//doi.org/10.5281/zenodo.22139642](https://doi.org/10.5281/zenodo.22139642), the permanent identifier for the dataset across all
67 of its versions, which resolves to whichever version is current; the version identifier above
68 pins the exact release measured in this manuscript (Sec. III).

69 I.A. Vertebra labeling when the scan cannot settle the count

70 *The limitation, stated plainly.* No scan in this corpus contains C2, so the conventional
71 count cannot be performed: a property of abdominal imaging, not of the annotation. A
72 thirteenth thoracic vertebra, a lumbar rib and a *stump rib* (a hypoplastic twelfth rib, a
73 cardinal indicator of a thoracolumbar transitional vertebra) leave the same number of rib-
74 free vertebrae between the two borders of the lumbar spine, the last rib-bearing vertebra
75 and the sacrum.

76 *What the release does in addition.* Every vertebra carries the identifier its radiologist-
77 sourced annotation assigned, corrected where the source was wrong; those labels are the
78 ground truth. Alongside them, every released measurement is also expressed against the
79 two landmarks present in essentially every abdominal scan, the sacrum below and the lowest
80 rib-bearing vertebra above, so that the measurement is positional rather than nominal: a
81 vertebra by how many rib-free vertebrae separate it from the sacrum, a lowest rib by its
82 length as a fraction of the rib above it, a transverse process by its span and its distance from
83 the ala. None of these changes according to whether a reader would have said L5 or S1, so
84 cases that disagree about the name remain comparable, and where the junction is in view
85 the count is determined and the two descriptions agree.

86 *But the vertebrae themselves are known to differ.* Counting is not the only route to an
87 identity. Level-specific morphometry is long established: pedicle width and height change
88 systematically from the thoracic to the lumbar spine,¹³ vertebral body, endplate and canal
89 dimensions differ by level,^{14–16} and the transverse process and the iliolumbar ligament mark
90 the last lumbar vertebra independently of any count.^{2,17} If those differences hold at the
91 boundary itself, the phenotype is decidable from the vertebra in front of the reader without
92 any global count. On this corpus it does hold: separating T12 from L1 on shape alone, with
93 each patient’s own size divided out and cross-validation grouped by case, gives an area under
94 the curve of 0.990 and 97.3% accuracy over 1485 vertebrae. Size is removed because raw
95 millimeters separate thoracic from lumbar trivially and teach nothing about the junction,
96 where the two are nearly the same size.

97 That is reported as a property of the released measurements rather than as a method:
98 the information needed to settle these variants is present locally.

TABLE I Public CT collections at the lumbosacral junction. Black, class present; outlined, graded only to exclude it⁷.

Collection	Scans	Count	L6	Sacrum	S1	Pelvis	Ribs	L. rib	Femora	Grade
CTSpine1K ³	1,005									
CTPelvic1K ⁴	1,184									
VerSe ⁶	374									
RibSeg v2 ¹⁸	660									
TotalSegmentator ²⁰	1,204									
CTSpinoPelvic1K	802									

II. ACQUISITION AND VALIDATION METHODS

II.A. Sources, and why a crosswalk was necessary

Both CTSpine1K³ and CTPelvic1K⁴ draw part of their imaging from the TCIA CT colonography collection^{11,12}, and the vertebral and pelvic annotations in both were produced under board-certified radiologist supervision. Neither released the mapping from an annotation to the specific CT series it was drawn on.

That mapping matters because every patient in CT COLONOGRAPHY was scanned *twice*, prone and supine, and an annotation carries a patient identifier and nothing finer. Turning a patient from supine to prone alters lumbar lordosis and the alignment of each segment against the next, so outlines drawn on one series are the wrong *shape* for the other and no rigid realignment recovers them. Such a mask looks competent in isolation and reveals itself only overlaid on the image. The crosswalk (Fig. 1) therefore resolves each annotation to a patient and then, within that patient’s volumes, to the series whose bone *agrees* with it: candidates are thresholded at bone attenuation and scored by overlap with the mask.

II.B. Fused and split records

CTSpine1K and CTPelvic1K each annotated a patient independently, so their labels are not guaranteed to come from the same CT series (Fig. 1). This turned out to matter for




FIG. 1 How the dataset was built. n counts scans, or corrections, at each step.

117 351 patients, nearly half the cohort: the two collections had labeled different acquisitions of
 118 the same person, one prone and one supine, with neither recording which. Because posture
 119 changes spinal alignment, a spine label from one acquisition cannot simply be paired with a
 120 pelvic label from the other. These 351 records are released but held out of any analysis that
 121 assumes a single posture. Two labeled acquisitions of the same patient in different postures
 122 are, instead, a resource for studying postural alignment, not a defect.

II.C. Densification, and why it is asymmetric

A corpus in which half the records lack a pelvis is not a spinopelvic dataset, so the missing structures were pseudolabeled, asymmetrically (Fig. 1). Pelvis onto spine-only records is the safe direction: the sacrum and innominate bones carry no enumeration to get wrong, and quality is reported as held-out performance on the *pelvic_only* records, withheld from training for that purpose. The reverse direction carries the whole problem this dataset addresses, since a pseudolabeled spine must commit to a count and on a transitional vertebra commits to the commoner reading; those 20 records were corrected by hand, tractable at that scale.

II.D. Ribs

No public rib annotation covers this cohort, and the two available automatic sources fail in complementary ways.

Möller’s binary rib network¹⁹ reports high accuracy, and its authors show that stump ribs can be classified from a partial field of view; that result concerns the morphological features computed on a rib once segmented. On this corpus the segmentation itself failed on ribs that were *partially outside* the field of view: a rib crossing the boundary of an abdominal acquisition was liable to be missed altogether rather than segmented to the edge. Its authors do not report this: their evaluation excluded subjects whose last rib was not visible, so it is recorded here as an observation from this cohort. Since this is a colonography cohort, the lower ribs are routinely clipped, and those are precisely the ribs the count depends on.

TotalSegmentator²⁰ does segment clipped ribs and supplies the per-rib numbering that a binary network cannot. Its characteristic failure is at the other end of the bone: the most medial portion of the rib (roughly the last tenth to sixth approaching the costovertebral joint) is frequently absent. A rib truncated at its medial end never reaches the vertebra it articulates with, so the rib–vertebra incidence check (the quality-control gate on the rib class labels, which matches each rib to the vertebra its number implies) has nothing to test.

The two were therefore unioned (Fig. 1): TotalSegmentator supplies numbering and the medial-clipped cases, Möller supplies detection where its output is more complete, and the union carries TotalSegmentator’s per-level identities.

The result is a pseudolabel whose review was triaged rather than exhaustive. A label-

based rule flagged every record in which one rib bone carried more than one identifier, or in which a rib failed to reach the vertebra it articulates with, since a single bone carrying two numbers is a numbering error by construction, whatever the anatomy. The rule selected 152 records; 148 were reviewed and corrected, and 4 were not and are identified in the release. Reviewers were medical students working through a student research consortium²¹, trained against a written labeling protocol, with every correction recorded against the annotator who made it. Records passing the rule were not individually inspected, so the layer only guarantees freedom from the failure modes the rule detects, no more. The residual rib-vertebra offsets reported in Sec. II.G are the independent check on what the rule missed.

II.E. Label scheme and counting anchors

The scheme is VerSe-native: vertebrae keep their VerSe identifiers (C1–C7 = 1–7, T1–T12 = 8–19, L1–L6 = 20–25, sacrum 26), and every non-VerSe structure takes a fixed identifier above that range: S1 carved from the sacrum (29), hips (30–31), femora (32–33), ribs (34–45 left, 46–57 right), lumbar ribs (74–75) and surgical hardware (76–82). Identifiers 58–73 are unassigned: a soft-tissue block once reserved there was never populated and is retired.

Two anchors make the count decidable, and Fig. 2 shows both: the lowest rib-bearing vertebra above, **S1** carved from the sacrum below. Each earns its place by what it supports rather than by being an endpoint. S1 supplies the sacral endplate landmark that sacral slope and pelvic incidence are measured from, and with both brackets present L5 and L6 are distinguishable where a spine-limited view alone would not be.

How S1 is cut. The sacrum’s outer boundary is the source annotation and is never altered; S1 is the cranial part of that same bone, separated by a plane. The plane’s normal is the sacrum’s own cranio-caudal axis, made orthogonal to a left–right axis measured directly rather than assumed, so as to remove the patient’s rotation within the scanner. That rotation is not negligible: median 4.5°, maximum 16.1°, and 508 of 801 records above 3°. The level of the cut preserves the sub-division volume of the preceding release, so the plane changes the shape and orientation of the S1/S2 boundary and not how much of the sacrum is called S1. Per-record geometry ships with the archive.

Neither anchor is novel on its own: TotalSegmentator²⁰ carries a separate S1 class and per-level ribs, and VerSe⁶ carries L6, whose identifier this release keeps, so its L6 *is* VerSe’s.




FIG. 2 The two anchors, from the released labels. Red, lowest rib-bearing vertebra and its rib; blue, S1; yellow, the rib-free vertebrae between. (b) and (c) both count four.

182 L6 is retained because a scheme without it cannot record a six-lumbar spine at all.

183 **A rib on a lumbar body receives its own class**, and this is the one class here with no
 184 published counterpart. A lumbar rib and a stump rib are the same object under two counts,
 185 so assigning it a number would commit the label to one reading of an enumeration anomaly.
 186 A scheme that numbers every rib 1–12 has nowhere to put a thirteenth: the annotator must
 187 either call it rib 12, asserting the vertebra beneath it is thoracic, the very question at issue,
 188 or discard it. TotalSegmentator carries ribs 1–12 per side and no such class, and VerSe’s
 189 T13 is a vertebra rather than a rib. This scheme can record both: VerSe’s T13 (28) for a
 190 thoracic-type thirteenth vertebra with a true rib, and 74/75 for a lumbar-type body bearing
 191 a rudimentary one, so either reading is recorded as itself. No record here carries a T13; the
 192 sixteen lumbar-rib cases were read as lumbar-type bodies. Where a border vertebra could
 193 not be settled by its readers the label stands as read, and reclassifying those records with
 194 a classifier trained on local vertebral morphology (Sec. V) is the intended path for a later
 195 release.

196 Figure 2 shows why the interval is the primitive and the label is derived. Among the
 197 records with four rib-free vertebrae, some reach that count through a lumbar rib and some
 198 without one; the two are indistinguishable by the count alone and are distinguished here
 199 only because rib presence is itself labeled.

200 Sixteen records carry a lumbar rib: thirteen bilateral and three unilateral, all on the
 201 right, a count too small to support any statement about side preference.

202 Eighteen records carry an L6. Fourteen are labeled lumbarization, two sacralization,
 203 one semi-sacralization and one normal, which is worth stating because it shows the two

204 annotation axes disagreeing in the open: a record can carry six lumbar-type vertebrae and
205 still be read as a sacralization, or as unremarkable, depending on where the reader started
206 the count.

207 Both counts are derived from the released label volumes, and so are the manifest fields
208 that report them. That is a correction: through v5, `has_l6` was true for one record that
209 contains no L6 and false for all eighteen that do, `has_lumbar_rib` was false everywhere, and
210 `n_lumbar_labels` read zero in 799 of 802 records. The fields are rebuilt for this release by
211 counting identifiers 25 and 74–75 in each volume, with `lumbar_rib_side` added alongside
212 them.

213 II.F. Computational tools

214 *Access.* Every step is performed by open-source code archived with the dataset, together
215 with the reference loader, the label scheme and the quality-control scripts that generated
216 the tables in this manuscript.

217 *Versions and parameters.* Segmentation of femora, the sacral sub-division and the per-
218 level rib numbering used TotalSegmentator²⁰ (`total` task, release 2.x) on a single graphics
219 processing unit. The binary rib network is Möller’s¹⁹ released weights, applied through the
220 nnU-Net v2 framework²². The pelvic completion for records whose pelvis was absent used
221 a five-fold nnU-Net v2 ensemble, applied out-of-fold: each record is completed by the fold
222 that never trained on it, so no record is completed by a model that has seen it. The five
223 fold checkpoints, with their nnU-Net plans and dataset descriptor, are released alongside the
224 data (<https://huggingface.co/anonymous-mlhc/spinopelvic-seg-checkpoints>). Im-
225 age handling throughout used NiBabel and SciPy; no image is resampled or reoriented when
226 a label is written, so every label shares its image’s grid and affine exactly.

227 *Generative artificial intelligence.* Large language models assisted with drafting and edit-
228 ing text and with analysis and figure code, under author review. They were not used to
229 generate, annotate or interpret imaging data; every number reported here is computed by
230 the released code from the released volumes.

231 II.G. Validation

232 Validation is layered, and the ordering matters: measurements computed on a corpus
233 that has not established internal consistency do not look broken, they look finished.

234 *Geometric invariants.* Every case is checked for label geometry agreeing with its CT in
235 shape, affine and voxel spacing; for identifiers within the published scheme; for a non-empty
236 label; and for sided structures falling on the correct sides, with the left–right axis read from
237 the affine rather than assumed. The check exists because a renumbering pass once wrote
238 a reoriented array under a canonical affine and transposed a label away from its image, a
239 failure invisible downstream.

240 The check compares *each sided pair separately*: testing the ribs alone passed all 802
241 records while four carried `left_hip` on the patient’s right, and pooling all sided structures
242 into one test still missed one of those four, because a large correctly-sided structure masks
243 a smaller transposed one. A gate that passes everything is not evidence of a clean corpus
244 until it has been shown capable of failing.

245 *Rib–vertebra incidence.* Each rib is matched to its nearest vertebra and compared against
246 the vertebra its number implies. Of 11,548 ribs, 5,788 are not evaluable because the expected
247 vertebra lies outside the field of view and 11 have no vertebra within the anchor distance;
248 of the 5,749 evaluable, 5,747 match, two (0.035%) are offset by one level, and no case is
249 misnumbered. The denominator is stated because quoting the two offsets against all 11,548
250 ribs would halve the rate by counting ribs the check never examined. Both offsets are
251 field-of-view truncations on individually characterized cases, one at +1 level and one at −1,
252 reported rather than suppressed.

253 The same check carries an invariant that reads as a null result. Of the 5,749 evaluable
254 ribs, *zero* are assigned to a lumbar vertebra, not because no lumbar ribs exist, since 16
255 records carry one, but because such a rib takes class 74 or 75 and never enters the numbered
256 set the test examines. A numbered rib landing on a lumbar vertebra would be a counting
257 error of exactly the kind this dataset exists to expose.

258 *Agreement with published values.* Derived spinopelvic measures are compared against
259 Vialle *et al.*²³ (Table II). Pelvic incidence is the strongest of these checks because it is a
260 morphological property of the pelvis rather than a posture, and so is directly comparable to
261 a standing reference cohort.

TABLE II Derived measures against a standing reference ($n = 260$)²³; this cohort is supine.

Measure	This dataset (supine)	Reference (standing)
Pelvic incidence	54.7°	54.7 ± 10.6
Pelvic tilt	17.3°	13 ± 6
Sacral slope	36.9°	41.0 ± 8.4
Lumbar lordosis (supine)	52.7°	43 ± 11.2 (standing)
PI – LL mismatch	2.6°	centered near 0

II.H. Castellvi typing

Every record whose vertebral count departs from five carries a Castellvi grade²⁴ in the released manifest: 33 cases, typed I–IV with the *a/b* unilateral–bilateral qualifier. This is released as an annotation layer because the grade and the count are *different axes* and the dataset is built to show that they are: grade IIIb occurs here at rib-free counts of four, five and six alike, and seven of the 33 graded cases carry a normal count of five.

Two radiology resident physicians graded every case independently and resolved their six disagreements by consensus; both reads and the consensus are in the manifest. The distinction the classification turns on, bony fusion against articulation without it², is the one carrying clinical weight, and it is where the disagreements lay: four of the six were III against II.

II.I. Surgical hardware, and why a threshold is not enough

Metal is trivial to detect and hard to interpret, and the difference matters here because **an iatrogenic fusion is indistinguishable from a congenital one to a distance measurement**: a cage-bridged interspace reads as “no gap” exactly as a fused transitional vertebra does. An unrecorded instrumented case is therefore a false positive on the central measure.

II.I.0.a. Detection over-calls by a factor of five. A 2500 HU threshold inside a shell around the labeled skeleton flags 52 of the 802 records. One of the two radiology resident physicians read all 52 and confirmed 11, rejecting 41 (79%) as contrast, calcification and reconstruction artefact. Attenuation does not separate the two groups (both saturate);




FIG. 3 Instrumentation in the release, one exemplar each, drawn through bone. (a) Hip arthroplasty, 80, $n = 8$. (b) Femoral-neck screws, 82, $n = 1$. (c) Sacroiliac screws, 81, $n = 1$. (d) Interbody cages, 77, $n = 1$. Bar, 5 cm.

site and volume do, every confirmed implant at $2,586\text{mm}^3$ or above and every rejection at $1,768\text{mm}^3$ or below. Instrumentation prevalence is 1.4%, not the 6.5% the threshold alone reports.

II.I.0.b. The subtype block had to be extended. Identifiers 76–79 name spinal instrumentation (generic, cage, screw-and-rod, plate) and cannot describe what this cohort holds, so **80 arthroplasty**, **81 sacroiliac screw** and **82 osteosynthesis** were added: osteosynthesis holds parts of one bone together where arthroplasty replaces a joint, and a shape rule calls a femoral stem a rod. The confirmed cases are 8 arthroplasties, one osteosynthesis, one sacroiliac fixation and one pair of interbody cages (Fig. 3).

II.I.0.c. Metal outranks bone, and that is a subtraction. In every confirmed case the implant was already inside a bone label, so naming it reclaims $1,538,852$ voxels rather than adding them. Pelvic incidence and tilt are measured from the femoral head center, and in 8 of these cases that head is an implant: any spinopelvic parameter previously computed from them was measured on metal.

III. DATA FORMAT AND USAGE NOTES

Each record is a gzipped NIfTI image/label pair sharing an identical 4×4 affine, canonicalized to PIR, requiring no resampling. Masks are NIfTI rather than DICOM: the policy prefers DICOM where applicable, and for volumetric segmentation masks NIfTI is what the downstream tooling expects.

Splits are patient-grouped and LSTV-stratified five-fold, frozen and shipped with the

TABLE III Data types and metadata, and the records populating each, from the released manifest.

Field	Records	Available
Identifier, image path, label path, configuration, match type	802	100.0%
Patient position, tube potential, section thickness, kernel	802	100.0%
Transitional label and class	802	100.0%
Spine and pelvic annotation origin	802	100.0%
Instrumentation and partial-annotation flags	802	100.0%
Image/label alignment check	802	100.0%
Spine series identifier; bone-fraction check	782	97.5%
Scanner manufacturer and model	768	95.8%
Sex	749	93.4%
Age	709	88.4%
Pelvic series identifier	713	88.9%
Castellvi grade (consensus of two readers)	33	4.1%

data as `splits_5fold.json`. The five validation folds partition all 802 records, so the release provides cross-validation and *not* a held-out test set; users reporting a single held-out number should carve one out and state which records it holds.

Stratification is on the transitional subtype rather than a binary LSTV flag: every validation fold carries three sacralization and three lumbarization records. With 15 of each across 802 that is the most stratification can guarantee, and it is why a fold-level metric on the rare classes carries an error bar far wider than its own decimal places.

Table III lists the key data types and metadata against the fraction of records for which each is populated.

The archive of record is Zenodo (<https://doi.org/10.5281/zenodo.22139642>). Code and documentation are on GitHub at the tagged release commit; any model-hub copy is a convenience mirror, not the archive.




FIG. 4 Count-free measures. (a) Vertebrae between the lowest rib and the sacrum. (b) Lowest rib length over the rib above. (c) Lowest lumbar transverse span against its gap to the ala; transitional labels marked.

IV. DESCRIPTIVE ANALYSIS

The cohort is a colorectal cancer screening population of 802 records. 738 are recorded female (393) or male (345); 11 carry DICOM’s *other* value, a recorded value and not a missing one, and 53 carry no sex field. Age is present for 709 (median 59, range 50–89). All 802 are counted in totals; the eleven records outside the female/male dichotomy are excluded from sex-stratified measures rather than folded into “unrecorded”. The lower age bound is the screening guideline, not a selection applied here.

Count-free measures (Fig. 4). Five rib-free vertebrae above the sacrum is typical; four and six are where transitional anatomy sits, and which of the two is present does not by itself determine what to call it. The lowest-rib length ratio is bimodal.

Level gradients and spinopelvic measures (Fig. 5). Vertebral dimensions increase caudally in step with the load each level carries. Pelvic incidence does not vary with age, while pelvic tilt increases, the descriptive signature of a pelvis retroverting as a spine ages.

Opportunistic measures. Every scan carries a vertebral bone density measurement at no additional dose.²⁵ L1 trabecular attenuation is reported at the published standard site.




FIG. 5 Spinopelvic measures against standing reference bands (mean $\pm$ SD)²³.

V. POTENTIAL APPLICATIONS AND LIMITATIONS

The principal use is the *spinopelvic interface*. Spine and pelvis carry radiologist-sourced annotation on one coordinate frame in 802 records, making sacral morphology measurable against the lumbar column rather than in isolation: the sacral endplate and its slope, the alar corridors constraining S2-alar-iliac and iliac screw trajectories, and the L5–S1 geometry a lumbosacral fusion must cross. Pelvic incidence is derived from the segmented sacrum and femoral heads rather than from landmarks placed on a radiograph, so it is reproducible from the labels themselves.

The transitional layer serves that use rather than competing with it: no trajectory can be planned across a junction whose segments cannot be named. Level-numbering benchmarks, variant studies and opportunistic screening follow from the same annotation.

V.A. Three uses the released measurements support

Naming a level from its own geometry. Wrong-level spine surgery runs at roughly one in 3,110 spinal procedures, and the cause most often cited is the transitional anatomy this cohort was assembled around.²⁶ Per-level body height, canal width, endplate width, transverse process span and wedge ratio are released for T11–L5 across all 802 records, and because the labels were assigned against the twelfth rib rather than by enumeration, a model trained on them is not learning the convention that fails.

Reference morphometry with its spread. The values a surgeon consults for vertebral body height, canal dimensions and pedicle width come from cadaveric series of a few dozen specimens, plotted as single values with no indication of spread.¹⁶ They cannot tell a reader




FIG. 6 Morphometry by level: median, interquartile range, 5th–95th percentile, n at right. (a) Body height. (b) Endplate width. (c) Canal. (d) Pedicle width. (e) Disc height. (f) Trabecular attenuation.

whether the patient in front of them is unusual, because they never show what usual looks like. The same measures are given here from 802 records with the distribution attached (Fig. 6), reproducing two properties of the classical figures (body height crosses over, dorsal exceeding ventral at T11–T12 and reversing by L4–L5, and pedicle width widens steeply into the lower lumbar spine) and adding the width of each distribution, which is the part needed to judge an individual. A larger n sharpens a distribution without changing the population it came from, and this one is a screening cohort aged 50 and over.

Patient-specific models. Planning is moving toward patient-specific biomechanical models rather than population norms.²⁷ Such a model needs spine, pelvis and femoral heads in one frame, which is this release’s construction. In 351 patients the annotation sits on two acquisitions in different positions, giving a within-patient change in alignment against which a predicted postural response can be checked.

Its limitations are stated at the level of detail a reader would need to catch them independently.




FIG. 7 Records carrying each vertebral level, cranial to caudal. Counts per identifier in Table S1.

Coverage. Thoracic ground truth is field-of-view limited (Fig. 7) and does not extend to T1. Postural angles are supine; pelvic incidence is not postural and needs no such caveat. The cohort is a colorectal screening population aged 50 and over, so its distributions should not be read as representative of a surgical one.

Declared but empty, and confirmed so. The partial-annotation sentinel (255) is declared in the scheme and occurs in no record, established by counting identifiers across all 802 released volumes rather than asserted. It is retained because the protocol is part of the scheme and a later release may ship a partially traced record. The hardware identifiers were in this position until v6 and are no longer: 76–82 are populated in the eleven instrumented records described in Sec. II.I.

Strength of the labels varies by structure, and the release does not average over that. Vertebral labels derive from radiologist-supervised source annotations, corrected where those sources were wrong. Pelvic labels on records that lacked one are pseudolabeled. The rib layer is a pseudolabel whose human review was triaged by an automated rule rather than exhaustive, so its guarantee is the absence of the failure modes that rule detects. Transitional labels derive from two independent sources and are not uniformly adjudicated, which is precisely why the measures reported here are count-free, and the Castellvi grades are a

consensus of two radiology resident physicians. The sacral sub-division inherits its level from an automatic method, and in 100 records that level places S1 above half the sacrum or below fifteen percent of it, which no first sacral segment is; those records are flagged in the released quality-control table and should be filtered before any measurement depending on the sacral endplate.

Structures are not universally present. The same census shows the release is not uniformly complete. Sacrum, hips and femora are present in all 802 records. One record carries no S1, one lacks T12, and nine lack an L5 identifier, in every case because the structure lies outside the field of view or, for L5, because the lowest lumbar segment is labeled L6 or incorporated into the sacrum. Users filtering on the presence of the caudal anchor should filter explicitly rather than assume it.

VI. DISCUSSION

Comparison with related material. Against the collections in Table I, the contribution is the crosswalk placing spine and pelvis on the same series, together with classes for the anatomy an enumeration anomaly actually produces. VerSe⁶ comes closest in intent: it also enriched its cohort for transitional anatomy and graded each case by Castellvi. The difference is what a graded case leaves behind. VerSe names the anomaly but does not segment the sacrum, so the fused or articulating structure the grade describes has no mask to go with it. This release segments that structure directly, as L6 or as a separately carved S1, so a user has the anatomy itself to measure rather than only a grade that describes it. TotalSegmentator²⁰ segments far more of the body than any source used here but has no class for a sixth lumbar vertebra or a rib on a lumbar body; what this release adds is not more anatomy per scan but the classes an enumeration anomaly needs in order to be recorded accurately.

Limitations. The cohort comes from one acquisition protocol of a colorectal-screening population aged 50 and over, so how well it generalizes to a pathologic population is untested. The rib layer’s human review was triaged by an automated rule rather than performed on every bone, so it guarantees only the absence of the failure modes that rule detects.

Future directions. The version history shows the scheme being extended more than once: femora and the S1 anchor, then per-level ribs, then lumbar ribs and surgical hardware, were

each folded into the existing release rather than requiring a new dataset. The same path is open going forward: imaging from beyond this cohort’s single protocol, and the C2 anchor this acquisition cannot supply, are the natural next additions.

VII. CONCLUSION

CTSpinoPelvic1K places spine, pelvis, per-level ribs and femora on one coordinate frame in 802 records, and gives the anatomy that makes lumbar numbering ambiguous a class of its own rather than recording it as something it is not. Because both counting anchors are explicit, a level can be identified from a field of view that cannot support the conventional count downward from C2, which is every record here.

DATA AVAILABILITY

The release described here is v8, permanently archived at <https://doi.org/10.5281/zenodo.22293878>; the concept DOI <https://doi.org/10.5281/zenodo.22139642> resolves to the current version. The archive carries the loader, the label scheme, the quality-control tables and the analysis code under an open-source licence. The repository URL identifies the authors; it is on the title page and will be restored here for the camera-ready version.

ACKNOWLEDGEMENTS

Acknowledgements are given on the title page, per the journal’s double-anonymized review policy. This work received no funding.

CONFLICT OF INTEREST

The authors have no relevant conflicts of interest to disclose.

ETHICS

This work uses publicly available, de-identified imaging and annotations derived from it. The authors’ institutional review board determined that it is not human participant research

under 45 CFR 46 and requires no oversight.

REFERENCES

- ¹J. Lian, N. Levine, and W. Cho, A review of lumbosacral transitional vertebrae and associated vertebral numeration, *Eur. Spine J.* **27**, 995–1004 (2018), doi:10.1007/s00586-018-5554-8.
- ²G. P. Konin and D. M. Walz, Lumbosacral transitional vertebrae: classification, imaging findings, and clinical relevance, *AJNR Am. J. Neuroradiol.* **31**, 1778–1786 (2010), doi:10.3174/ajnr.A2036.
- ³Y. Deng et al., CTSpine1K: a large-scale dataset for spinal vertebrae segmentation in computed tomography, *Mach. Learn. Biomed. Imaging* **3**, 824–832 (2025), doi:10.59275/j.melba.2025-gf84.
- ⁴P. Liu et al., Deep learning to segment pelvic bones: large-scale CT datasets and baseline models, *Int. J. Comput. Assist. Radiol. Surg.* **16**, 749–756 (2021), doi:10.1007/s11548-021-02363-8.
- ⁵H. Möller, H. Schoen, R. Graf, et al., VERIDAH: solving enumeration anomaly aware vertebra labeling across imaging sequences, 2026, arXiv:2601.14066. doi:10.48550/arXiv.2601.14066.
- ⁶A. Sekuboyina et al., VerSe: a vertebrae labelling and segmentation benchmark for multi-detector CT images, *Med. Image Anal.* **73**, 102166 (2021), doi:10.1016/j.media.2021.102166.
- ⁷H. Liebl et al., A computed tomography vertebral segmentation dataset with anatomical variations and multi-vendor scanner data, *Sci. Data* **8**, 284 (2021), doi:10.1038/s41597-021-01060-0.
- ⁸Y. Otake, S. Schafer, J. W. Stayman, et al., Automatic localization of vertebral levels in x-ray fluoroscopy using 3D-2D registration: a tool to reduce wrong-site surgery, *Phys. Med. Biol.* **57**, 5485–5508 (2012), doi:10.1088/0031-9155/57/17/5485.
- ⁹S.-F. L. Lo, Y. Otake, V. Puvanesarajah, et al., Automatic localization of target vertebrae in spine surgery: clinical evaluation of the LevelCheck registration algorithm, *Spine (Phila. Pa. 1976)* **40**, E476–E483 (2015), doi:10.1097/BRS.0000000000000814.

¹⁰T. De Silva, S.-F. L. Lo, N. Aygun, et al., Utility of the LevelCheck algorithm for decision support in vertebral localization, *Spine (Phila. Pa. 1976)* **41**, E1249–E1256 (2016), doi:10.1097/BRS.0000000000001589.

¹¹K. Smith et al., Data from CT COLONOGRAPHY, The Cancer Imaging Archive, 2015, doi:10.7937/K9/TCIA.2015.NWTESAY1.

¹²K. Clark et al., The Cancer Imaging Archive (TCIA): maintaining and operating a public information repository, *J. Digit. Imaging* **26**, 1045–1057 (2013), doi:10.1007/s10278-013-9622-7.

¹³M. R. Zindrick et al., Analysis of the morphometric characteristics of the thoracic and lumbar pedicles, *Spine (Phila. Pa. 1976)* **12**, 160–166 (1987), doi:10.1097/00007632-198703000-00012.

¹⁴M. M. Panjabi et al., Thoracic human vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **16**, 888–901 (1991), doi:10.1097/00007632-199108000-00006.

¹⁵M. M. Panjabi et al., Human lumbar vertebrae: quantitative three-dimensional anatomy, *Spine (Phila. Pa. 1976)* **17**, 299–306 (1992), doi:10.1097/00007632-199203000-00010.

¹⁶E. C. Benzell, *Biomechanics of Spine Stabilization*, Thieme, New York, 3rd edition, 2015.

¹⁷R. J. Hughes and A. Saifuddin, Numbering of lumbosacral transitional vertebrae on MRI: role of the iliolumbar ligaments, *AJR Am. J. Roentgenol.* **187**, W59–W65 (2006), doi:10.2214/AJR.05.0415.

¹⁸L. Jin et al., RibSeg v2: a large-scale benchmark for rib labeling and anatomical centerline extraction, *IEEE Trans. Med. Imaging* **43**, 570–581 (2024), doi:10.1109/TMI.2023.3313627.

¹⁹H. Möller et al., Automated thoracolumbar stump rib detection and analysis in a large CT cohort, 2025, arXiv:2505.05004. doi:10.48550/arXiv.2505.05004.

²⁰J. Wasserthal et al., TotalSegmentator: robust segmentation of 104 anatomic structures in CT images, *Radiol. Artif. Intell.* **5**, e230024 (2023), doi:10.1148/ryai.230024.

²¹[Reference to the authors’ own work, removed for double-anonymized review.]

²²F. Isensee, P. F. Jaeger, S. A. A. Kohl, J. Petersen, and K. H. Maier-Hein, nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation, *Nat. Methods* **18**, 203–211 (2021), doi:10.1038/s41592-020-01008-z.

²³R. Vialle, N. Levassor, L. Rillardon, A. Templier, W. Skalli, and P. Guigui, Radiographic analysis of the sagittal alignment and balance of the spine in asymptomatic subjects, *J.*

496 Bone Joint Surg. Am. **87**, 260–267 (2005), doi:10.2106/JBJS.D.02043.

497 ²⁴A. E. Castellvi, L. A. Goldstein, and D. P. K. Chan, Lumbosacral transitional vertebrae
 498 and their relationship with lumbar extradural defects, Spine (Phila. Pa. 1976) **9**, 493–495
 499 (1984), doi:10.1097/00007632-198407000-00014.

500 ²⁵P. J. Pickhardt, B. D. Pooler, T. Lauder, A. M. del Rio, R. J. Bruce, and N. Binkley,
 501 Opportunistic screening for osteoporosis using abdominal computed tomography scans
 502 obtained for other indications, Ann. Intern. Med. **158**, 588–595 (2013), doi:10.7326/0003-
 503 4819-158-8-201304160-00003.

504 ²⁶N. E. Epstein, A perspective on wrong level, wrong side, and wrong site spine surgery,
 505 Surg. Neurol. Int. **12**, 286 (2021), doi:10.25259/SNI.402_2021.

506 ²⁷E. Beaulieu et al., A review of finite element analysis in spine surgery decision-making, J.
 507 Clin. Med. **15**, 2584 (2026), doi:10.3390/jcm15072584.