

Host–Guest Systems | Hot Paper |

From Packed “Sandwich” to “Russian Doll”: Assembly by Charge-Transfer Interactions in Cucurbit[10]uril

WanJun Gong,^[a] Xiran Yang,^[a] Peter Y. Zavalij,^[c] Lyle Isaacs,^{*,[c]} Zhiyong Zhao,^[a] and Simin Liu^{*,[a, b]}

Abstract: As the host possessing the largest cavity in the cucurbit[n]uril (CB[n]) family, CB[10] has previously displayed unusual recognition and assembly properties with guests but much remains to be explored. Herein, we present the recognition properties of CB[10] toward a series of bipyridinium guests including the tetracationic cyclophane known as blue box along with electron-rich guests and detail the influence of encapsulation on the charge-transfer interactions between guests. For the mono-bipyridinium guest (methylviologen, MV^{2+}), CB[10] not only forms 1:1 and 1:2 inclusion complexes, but also enhances the charge-transfer interac-

tions between methylviologen and dihydroxynaphthalene (HN) by mainly forming the 1:2:1 packed “sandwich” complex (CB[10]·2 MV^{2+} ·HN). For guest 1 with two bipyridinium units, an interesting conformational switching from linear to “U” shape is observed by adding catechol to the solution of CB[10] and the guest. For the tetracationic cyclophane-blue box, CB[10] forms a stable 1:1 inclusion complex; the two bipyridinium units tilt inside the cavity of CB[10] according to the X-ray crystal structure. Finally, a supramolecular “Russian doll” was built up by threading a guest through the cavities of both blue box and CB[10].

Introduction

Simultaneous encapsulation of multiple components in a molecular container is an interesting topic because the reactivity and properties of molecules are often changed when they are trapped within well-defined nano-spaces.^[1] Another attractive feature of multi-component assemblies is that ordered sequences and conformations can be controlled.^[2] Among these multi-component assemblies, those that exhibit a “Russian doll” geometry are rare, but are particularly intriguing.^[3]

Cucurbit[n]uril (CB[n], $n = 5, 6, 7, 8, 10, 14$) (Figure 1a) is a family of macrocyclic hosts,^[4] the supramolecular and host–guest chemistry of which has been developing rapidly. With a hydrophobic cavity and $2n$ carbonyl oxygen atoms on the portals, CB[n] show selective molecular recognition properties towards cationic and neutral guests. There are numerous review papers focusing on different aspects of the CB[n] chemistry (general introduction,^[5] synthesis,^[6] high affinity,^[7]




Figure 1. Structures of: a) CB host, b) bipyridinium guests, and c) electron-rich aromatic guests used in this study.

[a] W. Gong, X. Yang, Dr. Z. Zhao, Prof. S. Liu
School of Chemistry and Chemical Engineering
Wuhan University of Science and Technology, Wuhan 430081 (P.R. China)
E-mail: liusimin@wust.edu.cn

[b] Prof. S. Liu
The State Key Laboratory of Refractories and Metallurgy
Wuhan University of Science and Technology, Wuhan 430081 (P.R. China)

[c] P. Y. Zavalij, Prof. L. Isaacs
Department of Chemistry and Biochemistry
University of Maryland, College Park, MD 20742 (USA)
E-mail: LIsaacs@umd.edu

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/chem.201604149>.

nanoreactor,^[8] conformational control,^[9] supramolecular polymer^[10]). Most recently, Scherman published a comprehensive introduction of CB[n] ($n=5, 6, 7, 8$) host-guest chemistry.^[11] The development of methods for the synthesis of monofunctionalized CB,^[12] broadens the scope of potential applications and enables workers in adjacent scientific fields to easily incorporate CB[n] hosts into their systems.

With the discovery of twisted CB[14] in 2013,^[4d] CB[10] is no longer the CB with the largest number of glycoluril units. However, CB[10] possesses the D_{nh} -symmetry of the smaller CB[n] and displays the largest cavity among the CB[n] family with the portal diameter of 9.5–10.6 Å and volume of ~ 870 Å³.^[13] This voluminous binding pocket endows it with unique molecular recognition and assembly properties, such as encapsulating calix[4]arene (including allosteric control),^[13b] non-natural oligomers,^[14] metalloporphyrin,^[15] or anti-cancer drugs.^[16]

Among the guests used in CB studies, methylviologen (MV^{2+}) has been established as a versatile probe to understand the connection between molecular recognition and redox properties due to its innate electrochemical properties by both the Kaifer and Kim groups. MV^{2+} forms a 1:1 inclusion complex with CB[7] and CB[8],^[17] and interestingly a 1:2 complex ($CB[8] \cdot 2MV^{2+}$) forms when MV^{2+} is chemically or electrochemically reduced to MV^{+} .^[17c] Moreover, CB[8] forms a stable 1:1:1 ternary complex with MV^{2+} and 2,6-dihydroxynaphthalene (HN) featuring charge-transfer interactions between the electron-rich and electron-deficient molecule.^[18] By taking advantage of this special recognition phenomenon, many groups made very interesting molecular machines and devices^[19] as well as supramolecular polymers.^[10b, 20]

Compared to the smaller CB[n] ($n=5, 6, 7, 8$) homologues, the explorations of the recognition properties of CB[10] are still in the very early stages. Herein, we report the encapsulation/assembly of bipyridinium molecules (including MV^{2+} , a dibipyridinium guest and the tetracationic cyclophane known as blue box (Figure 1b)) and electron-rich compounds (Figure 1c) inside CB[10], and the influence of encapsulation on the charge-transfer interactions in the cavity of CB[10]. Finally we demonstrate that a "Russian doll" pseudorotaxane can be assembled by threading a guest through the cavities of both the blue box and CB[10].

Results and Discussion

Encapsulation of MV^{2+} and HN inside CB[10]

Unlike other CB[n] compounds that are slightly soluble in water, CB[10] is extremely insoluble in pure H₂O, but dissolution is often observed when it forms inclusion complexes with water-soluble guests. In some cases, the complex is also insoluble in aqueous solution. Encapsulation of MV^{2+} (chloride salt) inside CB[10] was established based on the observed solubilisation of CB[10] and was also verified by ¹H NMR titration experiments. The signals for α - and β -protons on the pyridinium units of MV^{2+} undergo significant upfield shifts ($\Delta\delta=0.3$ and 0.5 ppm, respectively), whereas the CH₃ resonance remains constant (Figure 2a,b). However, the binding between MV^{2+}




Figure 2. Partial ¹H NMR spectra (600 MHz, D₂O) of: a) free MV^{2+} (1.0 mM); b) MV^{2+} (1.0 mM) and excess solid CB[10]; c) a 2:1 mixture of MV^{2+} (1.0 mM) and HN; and d) a 2:1 mixture of MV^{2+} (1.0 mM) and HN, and excess CB[10].

and CB[10] exhibits fast exchange kinetics on a ¹H NMR time-scale. Based on the integrals of the resonances in the ¹H NMR spectra shown in Figure 2b, we determined that the ratio between MV^{2+} and CB[10] is $\sim 1:0.9$, which implies the binding mode between MV^{2+} and CB[10] is probably 1:1.

As early as 2001, the Kim group reported that due to electrostatic repulsion CB[8] only binds one molecule of MV^{2+} , even though the cavity of CB[8] is big enough to encapsulate two MV^{2+} molecules.^[17c] In contrast, we found CB[10] can accommodate two molecules of MV^{2+} and this was clearly observed by ESI-MS. As shown in Figure 3a, besides the ion at m/z : 923.3, which corresponds to the 1:1 complex ($[CB[10] + MV^{2+}]^{2+} = 923.3$), ions at m/z : 508.2 and 689.6 were observed that correspond to the 1:2 complex ($[CB[10] + 2MV^{2+}]^{4+} = 508.2$; $[CB[10] + 2MV^{2+} + Cl]^{3+} = 689.6$). The formation of a 1:2 complex may be due to CB[10]'s large cavity, which minimizes electrostatic repulsion between the two MV^{2+} molecules. We are aware that ESI-MS results obtained in gas phase may not quantitatively reflect the situation in solution, but




Figure 3. ESI-MS spectra of: a) MV^{2+} with excess CB[10]; b) a 2:1 mixture of MV^{2+} and HN, and excess CB[10]; and c) a 1:2 mixture of MV^{2+} and HN, and excess CB[10].

have found that ESI-MS is a very powerful tool in CB host-guest systems.^[21]

As mentioned above, CB[8] forms a charge-transfer 1:1:1 ternary complex with MV^{2+} and 2,6-dihydroxynaphthalene (HN). In our case, we also observed that hetero-guest pairs bind into CB[10]. When a clear solution of MV^{2+} and HN was treated with solid CB[10], the solution immediately turned red. NMR titrations showed changes in the chemical shift of MV^{2+} and HN that are very similar to those observed upon addition of CB[8] (Figure 2c,d).^[18] When excess solid CB[10] was added to a solution of a 2:1 mixture of MV^{2+} (1.0 mM) and HN, we found the ratio between MV^{2+} and CB[10] is $\sim 1:0.6$ based on 1H NMR integration. When 1:1 or 1:2 mixtures of MV^{2+} (1.0 mM) and HN were treated similarly, the ratio between MV^{2+} and CB[10] is 1:0.7–0.8. Although the binding process displays fast kinetics of exchange on the 1H NMR timescale, these results imply that a 1:2:1 quaternary complex CB[10]·2 MV^{2+} ·HN was the major complex. ESI-MS (Figure 3b) clearly revealed the presence of the quaternary complex based on ions at m/z : 548.2, 730.9, 742.6, and 1132.4. In terms of the geometry of the charge-transfer complex, an alternating “packed sandwich” in which HN is in the middle and two MV^{2+} are on the sides is proposed (Scheme 1), similar to the alternating aromatic stacks found in Fujita’s cage.^[22] Although the 1:1:1 ternary complex was found (m/z peak at 1003.8), its intensity was quite low in the ESI-MS spectrum.

Interestingly, when a sample containing more HN (MV^{2+} and HN with a ratio of 1:2, and excess CB[10]) was examined by ESI-MS, we observed a 1:2:2 pentamolecular complex (m/z peak at 587.95) (Figure 3c)! According to the charge-transfer interaction and MMFF calculation, we believe that in this complex two HN molecules are located between two MV^{2+} molecules (Scheme 1a). Although, we hypothesized that a 1:1:2 quaternary complex with composition CB[10]· MV^{2+} ·2HN could also form, ESI-MS analysis did not support this expectation.


UV/Vis spectroscopy showed that in the presence of CB[10], the charge-transfer interaction between MV^{2+} and HN has been significantly enhanced, which is reflected in the increased charge-transfer absorption at ≈ 490 nm (Figure S1 in the Sup-

porting Information). As observed previously for CB[8],^[18] the addition of CB[10] resulted in a large quenching of HN fluorescence in the presence of equimolar MV^{2+} (Figure S2 in the Supporting Information).

Inducing conformational change from linear to “U” shaped

Inspired by the formation of the 2:1 complex of MV^{2+} and CB[10], we wondered whether a covalently linked MV^{2+} dimer would bind to CB[10] with a “U”-shaped conformation? Accordingly, we synthesized compound **1** (bromide salt) by the methylation of the intermediate used in the synthesis of blue box (**2**).^[23]

Unfortunately **1** itself did not form a “U” shape inside CB[10], even though the sample with a precise ratio of 1:1 was prepared. This guest did form a 1:2 complex with CB[10] in the presence of excess CB[10], as determined by the 1H NMR integrals for peaks of guest and host (Figure 4c) and ESI-MS. In the ESI-MS spectrum (Figure S3a and Table S1, Supporting Information), no ions were observed that could be ascribed to 2:2 complexes or supramolecular polymers. For comparison, we also tested the binding of guest **1** with CB[8]. 1H NMR titration showed that **1** could also form 1:1 and 1:2 complexes with CB[8] (Figures S4 and S5, Supporting Information). In the 1:1 complex, CB[8] was located in the central region (e.g., phenyl ring) of the guest and the 1:1 binding exhibits slow exchange on the 1H NMR timescale. The 1H NMR spectrum of the 1:2 complex is very similar to the Figure 4c. Interestingly, addition of catechol (1 equiv) to a solution of **1** and CB[10] (1:2) resulted in rapid formation of a precipitate. After the solid was removed by centrifugation, the solution had a pale-red color and the ratio of **1**/CB[10] changed from 1:1.9 (Figure 4c) to $\approx 1:1.1$ (Figure 4d) according to the 1H NMR integration (peaks for catechol could not be found due to broadening). The signals for H₁ on the phenyl group and H_{4/5} on the bipyridinium group underwent significant downfield and upfield shifts, respective-



Scheme 1. Schematic illustration of the formation of an inclusion complex between guests and CB[10].




Figure 4. Partial 1H NMR spectra (600 MHz, D_2O) of: a) free **1** (1.0 mM); b) a 1:1 mixture of **1** (1.0 mM) and catechol (resonance signals of catechol are circled); c) **1** (1.0 mM) and excess CB[10]; and d) **1** (1.0 mM) and excess CB[10] with addition of 1.0 equiv catechol (precipitate was observed).

ly. The ESI-MS spectrum (Figure S3b and Table S1, Supporting Information) provided strong evidence for the formation of a 1:1:1 ternary complex. In contrast, there were no obvious changes in chemical shift for a mixture of **1** and catechol (Figure 4a,b) in the absence of CB[10]. When **HN** was used instead of catechol, similar chemical shifts were observed but all peaks on **1** were broad, and precipitation occurred more slowly. Based on the combined inference of the above experiments, we are confident that the addition of catechol guest induced the linear shape of **1** to a "U"-shaped conformation (Scheme 1b) due to the enhancing charge-transfer interaction between **1** and catechol by CB[10]. Structurally related U-shaped complexes inside CB[8] have previously been observed by Kim and by us,^[24] and external stimuli-responsive versions have been created by Schalley.^[19b,21]

Encapsulation of **2** within CB[10]

The formation of the 1:2 complex between CB[10] and **MV**²⁺ also lead us to consider whether a smaller macrocyclic compound containing two bipyridinium groups would be encapsulated inside the cavity of CB[10]. Among these types of cyclic compounds, **2** (blue box) is a known building block pioneered by the Stoddart group to create mechanically interlocked molecular machines^[25] based on its redox properties and its ability to bind to electron-rich aromatic compounds through charge-transfer interactions. Mixing CB[10] and **2** (chloride salt) resulted in the formation of the soluble CB[10]·**2** complex as established by ¹H NMR spectroscopy (Figure 5a–c). The 1:1 inclusion complex CB[10]·**2** exhibits slow kinetics of exchange on the ¹H NMR timescale. The complexation-induced changes in chemical shift allow us to conclude that the two bipyridinium moieties are located inside the cavity of CB[10], whereas the linking *p*-phenylene units are outside of the cavity of CB[10]. In theory, *p*-xylylenedication could compete with bipyridinium group for cavity inclusion (a quick test shows that *p*-xylylenediamine can make CB[10] soluble in H₂O), but this does not occur in practice. We think the main reason is that the bipyridinium group on **2** is too long to reside at the portal of CB[10], if the *p*-xylylene unit were located inside. Another reason is probably because the bipyridinium group binds to CB[10] stronger than *p*-xylylenedications. As an important control experiment, we verified that **2** is incapable of undergoing cavity inclusion within the less spacious cavity of CB[8].

To gauge the strength of the interaction between CB[10] and **2** we performed ¹H NMR competition experiments as described previously.^[24a] For this purpose we added CB[5] (0.25 mM) to the aqueous solution of CB[10]·**2** (0.25 mM) and followed the composition of the mixture over time by ¹H NMR. Equilibrium was reached only after several days^[24a,26] and we used the known concentrations of the various species and the mass balance relationships to calculate a relative binding constant (*K*_{rel}) in the usual manner. The fact that the binding constant for tetracationic **2** is comparable to that of neutral CB[5] (*K*_{rel} = 2.2) was somewhat surprising and probably reflects the complementarity between the external surface of CB[5] and the internal surface of CB[10]. We performed dilution experi-




Figure 5. ¹H NMR spectra (600 MHz, D₂O) of: a) free **2** (1.0 mM); b) a 2:1 mixture of **2** and CB[10] (resonances of free **2** are marked with *); c) CB[10]·**2**; d) CB[10]·**2**·BHEEN; e) **2**·BHEEN; and f) free BHEEN (1.0 mM).

ments (1.0 mM to 10 μM CB[10]·**2**, Figure S6, Supporting Information) and did not observe any signals that could be assigned to free guest **2**. Assuming that we could detect 5% free **2** allows us to place a lower limit on *K*_a for CB[10]·**2** as > 10⁷ M^{−1}.^[27] We also compared the change in photophysical and electrochemical characteristics between free and bound **2**. Adding one equivalent of CB[10] to a solution of **2** leads to a dramatic decrease in the current levels in the cyclic voltammogram. Our interpretation is that encapsulation of **2** inside CB[10] reduced the diffusion coefficient of the complex (Figures S7 and S8 in the Supporting Information). In contrast, for CB[7]·**MV**²⁺, the current level continuously decreased even when the amount of CB[7] added increased to three equivalents.^[16b,17a] For CB[8]·**MV**²⁺, due to the formation of CB[8]·**2MV**⁺, two additional redox waves appeared when the amount of CB[8] was added to 0.25 equivalents.^[17b]

X-ray crystal structure of CB[10]·**2**

Luckily, single crystals of CB[10]·**2** that were suitable for structure determination were obtained from aqueous solution assisted by the addition of a small amount of KI.^[28] In agreement with the conclusions based on the ¹H NMR data, the crystal structure directly shows that the bipyridinium units of **2** were located inside the cavity of CB[10] but are tilted with respect to the potential 10-fold symmetry axis (Figure 6). The dihedral angle between the mean plane of **2** and the equatorial plane of CB[10] is ~59.6°, which is very similar to the dihedral angle observed for CB[10]·CB[5] (~62.7°).^[13a] The stability of the CB[10]·**2** complex can be attributed to the ion-dipole interactions between C=O...N⁺ (the shortest distance between portal O on CB[10] and N on **2** is 2.93 Å), the presence of multiple




Figure 6. X-ray crystal structure of the inclusion complex CB[10]·2 (left: side view; middle: channel view; right: packing view).

C=O...H–C close contacts (the shortest distance between portal C=O on CB[10] and C–H on **2** is ~ 2.27 Å), and of course the hydrophobic effect. Figure 6b nicely illustrates that, even though **2** is encapsulated by CB[10], there is still a rectangular channel through the two molecules of approximate 3.0×4.6 Å dimensions. Interestingly in the packing structure, the heads of three guests are very close to each other and the angle between them is almost 120° (accurately 119.86°). Moreover, the triangular structure of three CB[10], which was stabilized by multiple C=O...H–C close contacts (the shortest distance between portal O on CB[10] and H–C on the adjacent CB[10] is ~ 2.33 Å) is quite similar to that of reported free CB[10].^[6a] It seems that inclusion of **2** inside CB[10] does not significantly perturb the packing preferences of CB[10] in the solid state (more packing views can be found in Figure S9 in the Supporting Information).

Binding of electron-rich aromatics inside **2** or CB[10]·2

Blue box **2** is a known host for various molecules containing electron-rich aromatic groups.^[29] Previously, we have shown that **2** can form an encapsulation complex with *ns*-CB[10],^[30] but we never checked the binding ability of **2** within the *ns*-CB[10]·**2** complex because we thought the two “inverted” CH₂ bridges in the middle of *ns*-CB[10] would occupy the cavity of **2**. As described above, the CB[10]·**2** complex exhibits a sizable rectangular channel through two hosts. Along with the importance of the synergistic effect in the molecular recognition of CB,^[13b,31] we were interested to explore the synergistic effect involved when a host binds inside another larger host. Several aromatic guest molecules containing benzene or naphthalene rings with different substituents were selected for the study (Figure 1c).

Similar to a previous study,^[29] a visible absorption band arises when a solution of either **2** or CB[10]·**2** was titrated with a guest, due to charge-transfer interactions. Figure 7 shows the charge-transfer absorption band that appeared when adding 1,5-DAN to a solution of CB[10]·**2**. The equilibrium binding constants can be determined by fitting the change in absorbance of CB[10]·**2** as a function of [guest]. The K_a values of **2** or CB[10]·**2** towards guests are given in Table 1. In the




Figure 7. Plot of the absorbance at 658 nm as a function of 1,5-DAN concentration, the solid line is the curve fitting of the experimental data to the 1:1 binding model, $K_a = 2.3 \times 10^4 \text{ M}^{-1}$ (insert: UV/Vis spectra recorded during the titration of CB[10]·**2** (0.1 mM) with 1,5-DAN (0–1.8 mM) in Tris-HCl buffer (50 mM, pH 7.0), showing the charge-transfer band at $\lambda_{\text{max}} = 658$ nm).

case of small guest catechol, its binding constant is five-fold higher with CB[10]·**2** than with **2**, probably because CB[10] provides secondary binding interactions that increase the hydrophobic effect for catechol. When indole is used, a slight increase in K_a of 36% is observed. Similarly, when moderate-sized guests like 1-NCA and 2-NCA are used, the binding constants toward **2** or CB[10]·**2** are comparable. We know CB[*n*] likes to bind to cationic species through the ion-dipole interactions, so we thought the binding constant would increase if an amino compound was used as a guest. Disappointingly, when di-amino compound 1,5-DAN was used as the guest, the binding constant of 1,5-DAN toward CB[10]·**2** was only slightly higher than that with **2**. We hypothesize that **2** has already satisfied the electrostatically negative portals of CB[10] such that the 1,5-DAN does not engender additional ion-dipole interactions with CB[10] in the CB[10]·**2**·1,5-DAN complex. When bigger and longer guests like DL-Trp and DA are used, the binding constant decrease in the presence of CB[10] might be

Table 1. Values of K_a [M^{-1}] measured for the interaction of electron-rich aromatic guests with **2**/CB[10]·**2** in Tris-HCl buffer (50 mM, pH 7.0) at 25 °C.

Guest	2 [M^{-1}] (λ_{max} [nm])	CB[10]· 2 [M^{-1}] (λ_{max} [nm])
catechol ^[a]	$(2.1 \pm 0.1) \times 10^3$ (448)	$(1.0 \pm 0.1) \times 10^4$ (455)
indole ^[a]	$(1.4 \pm 0.1) \times 10^4$ (468)	$(1.9 \pm 0.2) \times 10^4$ (416)
1-NCA ^[a]	$(1.5 \pm 0.1) \times 10^4$ (423)	$(1.8 \pm 0.1) \times 10^4$ (438)
2-NCA ^[a]	$(9.8 \pm 1.1) \times 10^3$ (409)	$(1.0 \pm 0.1) \times 10^4$ (409)
1,5-DAN ^[a]	$(1.4 \pm 0.1) \times 10^4$ (686)	$(2.3 \pm 0.3) \times 10^4$ (658)
DL-Trp ^[b]	$(8.7 \pm 0.2) \times 10^2$ (412)	$(4.3 \pm 0.2) \times 10^2$ (401)
DA ^[b]	$1.1 \times 10^{3[c]}$	$(1.7 \pm 0.2) \times 10^2$ (487)

[a] Determined by adding the concentrated guest solution in DMSO.

[b] Determined by adding the aqueous guest solution. [c] From ref. [29].

due to steric hindrance caused by a mismatch between the geometrical preferences of the **2**-guest and CB[10]·**2**-guest complexes (curve fitting for the UV/Vis titration was shown in Figure S10, Supporting Information). Based on the above results, we conclude that encapsulation of **2** inside CB[10] does not influence its recognition properties significantly. The reasons are probably because the binding is dynamic and the encapsulation of compound **2** in CB[10] did not change the relatively rigid structure of **2**.

"Russian doll" assembly

The first "Russian doll"-like assembly based on CB was reported by the Kim group.^[3a] Threading a long-chain guest through both hosts comprising the CB[10]·**2** complex would allow us to prepare a new example of the type of intriguing supramolecular architecture. For this purpose, we selected **BHEEN** (1,5-bis[2-(2-hydroxyethoxy)ethoxy]naphthalene), which is an ideal guest because it forms a pseudorotaxane with **2** with a high $K_a \sim 1.5 \times 10^6 M^{-1}$ in aqueous solution, driven by charge-transfer interactions and the hydrophobic effect.^[32] In Figure 5 e, observation of two sets of host signals is due to the time-averaged C_{2h} symmetry of the complex **2**·**BHEEN** and the slow exchange on the NMR timescale, which is in agreement with previous results.^[33] When CB[10] was added to the solution of **2**·**BHEEN**, as shown in Figure 5 d, dramatic upfield shifts of the $H_{2/6r}$, $H_{3/7r}$, and $H_{4/8}$ protons on the naphthalene unit and $H_{d/d'}$, $H_{c'}$ on bipyridinium units indicate that both **BHEEN** and **2** are encapsulated by CB[10] to form the "Russian doll" assembly CB[10]·**2**·**BHEEN**. Molecular modeling shows that **BHEEN** does thread through the cavity of two hosts (Figure 8). To our surprise, however, both NMR data (compare Figure 5 e with d, H_c on bipyridinium unit slight shifts downfield) and modeling (dihedral angle between the planes of **2** and CB[10] is $\sim 68.3^\circ$, compared to $\sim 59.6^\circ$ in the CB[10]·**2**) show that **2** in this complex prefers to close the portal entrance of CB[10], instead of decreasing the dihedral angle for the guest threading.

Conclusion

We have presented the binding of CB[10] with bipyridinium-containing guests. With mono-bipyridinium (methylviologen)




Figure 8. Modeling of CB[10]·**2**·**BHEEN** (MMFF).

as the guest, CB[10] not only forms 1:1 and 1:2 inclusion complexes, but also enhances the charge-transfer interactions between MV^{2+} and **HN** by forming a 1:2:1 "sandwich" complex. When bis(bipyridinium) compound **1** was used as the guest we observed an interesting conformational switching from linear to "U"-shaped upon addition of catechol to a solution of CB[10] and **1**. We found that the tetracationic cyclophane blue box forms a stable 1:1 inclusion complex with CB[10]. Interestingly, two bipyridinium units tilt inside the cavity of CB[10] according to the X-ray crystal structure. The encapsulation of the blue box does not significantly affect its host-guest recognition properties.

Compared to other CB[n]s ($n=5, 6, 7$), the larger cavity of CB[8] is more useful in the fabrication of supramolecular polymers^[10] or as a nano-reactor.^[8] The capacity of CB[10] for the simultaneous encapsulation of multiple guests makes it attractive as a building block to create 3D multi-functional supramolecular materials or to promote multi-component reactions. Furthermore, encapsulation of a host in a bigger host and the assembly of a "Russian doll" expand the use of CB[10] as a fascinating new building block to create new molecular machines and devices.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (No. 21472143), Thousand Youth Talents Program of China (No. D1118031) (to S. L.), and the US National Science Foundation (CHE-1404911 to L. I.). We also thank Joshua A. Wilhide at the University of Maryland Baltimore County for the ESI measurement.

Keywords: charge-transfer interactions • cucurbituril • host-guest systems • Russian dolls • supramolecular chemistry

- [1] M. Yoshizawa, J. K. Klosterman, M. Fujita, *Angew. Chem. Int. Ed.* **2009**, *48*, 3418–3438; *Angew. Chem.* **2009**, *121*, 3470–3490.
- [2] J. K. Klosterman, Y. Yamauchi, M. Fujita, *Chem. Soc. Rev.* **2009**, *38*, 1714–1725.
- [3] a) S.-Y. Kim, I.-S. Jung, E. Lee, J. Kim, S. Sakamoto, K. Yamaguchi, K. Kim, *Angew. Chem. Int. Ed.* **2001**, *40*, 2119–2121; *Angew. Chem.* **2001**, *113*, 2177–2179; b) A. Shivanyuk, *J. Am. Chem. Soc.* **2007**, *129*, 14196–14199; c) S. J. Dalgarno, J. L. Atwood, C. L. Raston, *Chem. Commun.* **2006**, 4567–4574; d) Y. Han, T.-L. Liang, X. Hao, C.-F. Chen, *CrystEngComm* **2016**, *18*, 4900–4904.
- [4] a) W. A. Freeman, W. L. Mock, N. Y. Shih, *J. Am. Chem. Soc.* **1981**, *103*, 7367–7368; b) J. Kim, I.-S. Jung, S.-Y. Kim, E. Lee, J.-K. Kang, S. Sakamoto, K. Yamaguchi, K. Kim, *J. Am. Chem. Soc.* **2000**, *122*, 540–541; c) A. I. Day, A. P. Arnold, R. J. Blanch, B. Snushall, *J. Org. Chem.* **2001**, *66*, 8094–8100; d) X.-J. Cheng, L.-L. Liang, K. Chen, N.-N. Ji, X. Xiao, J.-X. Zhang, Y.-Q. Zhang, S.-F. Xue, Q.-J. Zhu, X.-L. Ni, Z. Tao, *Angew. Chem. Int. Ed.* **2013**, *52*, 7252–7255; *Angew. Chem.* **2013**, *125*, 7393–7396.
- [5] a) B. Han, Y. Liu, *Chin. J. Org. Chem.* **2003**, *23*, 139–149; b) J. W. Lee, S. Samal, N. Selvapalam, H.-J. Kim, K. Kim, *Acc. Chem. Res.* **2003**, *36*, 621–630; c) J. Lagona, P. Mukhopadhyay, S. Chakrabarti, L. Isaacs, *Angew. Chem. Int. Ed.* **2005**, *44*, 4844–4870; *Angew. Chem.* **2005**, *117*, 4922–4949; d) E. Masson, X. Ling, R. Joseph, L. Kyeremeh-Mensah, X. Lu, *RSC Adv.* **2012**, *2*, 1213–1247.
- [6] a) L. Isaacs, *Chem. Commun.* **2009**, 619–629; b) K. Kim, N. Selvapalam, Y. H. Ko, K. M. Park, D. Kim, J. Kim, *Chem. Soc. Rev.* **2007**, *36*, 267–279.
- [7] a) K. I. Assaf, W. M. Nau, *Chem. Soc. Rev.* **2015**, *44*, 394–418; b) D. Shetty, J. K. Khedkar, K. M. Park, K. Kim, *Chem. Soc. Rev.* **2015**, *44*, 8747–8761.
- [8] B. C. Pemberton, R. Raghunathan, S. Volla, J. Sivaguru, *Chem. Eur. J.* **2012**, *18*, 12178–12190.
- [9] a) A. E. Kaifer, *Acc. Chem. Res.* **2014**, *47*, 2160–2167; b) L. Isaacs, *Acc. Chem. Res.* **2014**, *47*, 2052–2062.
- [10] a) S. Gürbüz, M. Idris, D. Tuncel, *Org. Biomol. Chem.* **2015**, *13*, 330–347; b) Y. Liu, H. Yang, Z. Wang, X. Zhang, *Chem. Asian J.* **2013**, *8*, 1626–1632.
- [11] S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio, O. A. Scherman, *Chem. Rev.* **2015**, *115*, 12320–12406.
- [12] a) D. Lucas, T. Minami, G. Iannuzzi, L. Cao, J. Wittenberg, P. Anzenbacher, L. Isaacs, *J. Am. Chem. Soc.* **2011**, *133*, 17966–17976; b) B. Vinciguerra, L. Cao, J. R. Cannon, P. Y. Zavalij, C. Fenselau, L. Isaacs, *J. Am. Chem. Soc.* **2012**, *134*, 13133–13140; c) N. Zhao, G. O. Lloyd, O. A. Scherman, *Chem. Commun.* **2012**, *48*, 3070–3072; d) Y. Ahn, Y. Jang, N. Selvapalam, G. Yun, K. Kim, *Angew. Chem. Int. Ed.* **2013**, *52*, 3140–3144; *Angew. Chem.* **2013**, *125*, 3222–3226; e) M. M. Ayhan, H. Karoui, M. Hardy, A. Rockenbauer, L. Charles, R. Rosas, K. Udachin, P. Tordo, D. Bardelang, O. Ouari, *J. Am. Chem. Soc.* **2015**, *137*, 10238–10245.
- [13] a) A. I. Day, R. J. Blanch, A. P. Arnold, S. Lorenzo, G. R. Lewis, I. Dance, *Angew. Chem. Int. Ed.* **2002**, *41*, 275–277; *Angew. Chem.* **2002**, *114*, 285–287; b) S. Liu, P. Y. Zavalij, L. Isaacs, *J. Am. Chem. Soc.* **2005**, *127*, 16798–16799.
- [14] S. Liu, P. Y. Zavalij, Y.-F. Lam, L. Isaacs, *J. Am. Chem. Soc.* **2007**, *129*, 11232–11241.
- [15] S. Liu, A. D. Shukla, S. Gadde, B. D. Wagner, A. E. Kaifer, L. Isaacs, *Angew. Chem. Int. Ed.* **2008**, *47*, 2657–2660; *Angew. Chem.* **2008**, *120*, 2697–2700.
- [16] a) M. J. Pisani, Y. Zhao, L. Wallace, C. E. Woodward, F. R. Keene, A. I. Day, J. G. Collins, *Dalton Trans.* **2010**, 39, 2078–2086; b) F. Li, M. Feterl, J. M. Warner, A. I. Day, F. R. Keene, J. G. Collins, *Dalton Trans.* **2013**, 42, 8868–8877.
- [17] a) H.-J. Kim, W. S. Jeon, Y. H. Ko, K. Kim, *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 5007–5011; b) W. Ong, M. Gómez-Kaifer, A. E. Kaifer, *Org. Lett.* **2002**, *4*, 1791–1794; c) W. S. Jeon, H. J. Kim, C. Lee, K. Kim, *Chem. Commun.* **2002**, 1828–1829.
- [18] H.-J. Kim, J. Heo, W. S. Jeon, E. Lee, J. Kim, S. Sakamoto, K. Yamaguchi, K. Kim, *Angew. Chem. Int. Ed.* **2001**, *40*, 1526–1529; *Angew. Chem.* **2001**, *113*, 1574–1577.
- [19] a) W. S. Jeon, A. Y. Ziganshina, J. W. Lee, Y. H. Ko, J. K. Kang, C. Lee, K. Kim, *Angew. Chem. Int. Ed.* **2003**, *42*, 4097–4100; *Angew. Chem.* **2003**, *115*, 4231–4234; b) Y. H. Ko, K. Kim, J.-K. Kang, H. Chun, J. W. Lee, S. Sakamoto, K. Yamaguchi, J. C. Fetting, K. Kim, *J. Am. Chem. Soc.* **2004**, *126*, 1932–1933.
- [20] a) Y. J. Jeon, P. K. Bharadwaj, S. Choi, J. W. Lee, K. Kim, *Angew. Chem. Int. Ed.* **2002**, *41*, 4612–4612; *Angew. Chem.* **2002**, *114*, 4807–4807; b) J. Zhang, R. J. Coulston, S. T. Jones, J. Geng, O. A. Scherman, C. Abell, *Science* **2012**, *335*, 690–694; c) E. A. Appel, X. J. Loh, S. T. Jones, F. Biedermann, C. A. Dreiss, O. A. Scherman, *J. Am. Chem. Soc.* **2012**, *134*, 11767–11773; d) R. M. Parker, J. Zhang, Y. Zheng, R. J. Coulston, C. A. Smith, A. R. Salmon, Z. Yu, O. A. Scherman, C. Abell, *Adv. Funct. Mater.* **2015**, *25*, 4091–4100; e) Y. Zheng, Z. Yu, R. M. Parker, Y. Wu, C. Abell, O. A. Scherman, *Nat. Commun.* **2014**, *5*, 5772–5780; f) J. Tian, T.-Y. Zhou, S.-C. Zhang, S. Aloni, M. V. Altoe, S.-H. Xie, H. Wang, D.-W. Zhang, X. Zhao, Y. Liu, Z.-T. Li, *Nat. Commun.* **2014**, *5*, 5574–5584.
- [21] L. Cera, C. A. Schalley, *Chem. Sci.* **2014**, *5*, 2560–2567.
- [22] M. Yoshizawa, J. Nakagawa, K. Kumazawa, M. Nagao, M. Kawano, T. Ozeki, M. Fujita, *Angew. Chem. Int. Ed.* **2005**, *44*, 1810–1813; *Angew. Chem.* **2005**, *117*, 1844–1847.
- [23] P. L. Anelli, P. R. Ashton, R. Ballardini, V. Balzani, M. Delgado, M. T. Gandolfi, T. T. Goodnow, A. E. Kaifer, D. Philp, M. Pietraszkiewicz, L. Prodi, M. V. Reddington, A. M. Z. Slawin, N. Spencer, J. F. Stoddart, C. Vicent, D. J. Williams, *J. Am. Chem. Soc.* **1992**, *114*, 193–218.
- [24] a) S. Liu, C. Ruspici, P. Mukhopadhyay, S. Chakrabarti, P. Y. Zavalij, L. Isaacs, *J. Am. Chem. Soc.* **2005**, *127*, 15959–15967; b) Y. H. Ko, H. Kim, Y. Kim, K. Kim, *Angew. Chem. Int. Ed.* **2008**, *47*, 4106–4109; *Angew. Chem.* **2008**, *120*, 4174–4177.
- [25] a) A. C. Fahrenbach, S. Sampath, D. J. Late, J. C. Barnes, S. L. Kleinman, N. Valley, K. J. Hartlieb, Z. Liu, V. P. Dravid, G. C. Schatz, R. P. V. Duyne, J. F. Stoddart, *ACS Nano* **2012**, *6*, 9964–9971; b) Z. Zhu, A. C. Fahrenbach, H. Li, J. C. Barnes, Z. Liu, S. M. Dyar, H. Zhang, J. Lei, R. Carmieli, A. A. Sarjeant, C. L. Stern, M. R. Wasielewski, J. F. Stoddart, *J. Am. Chem. Soc.* **2012**, *134*, 11709–11720; c) J. C. Barnes, A. C. Fahrenbach, D. Cao, S. M. Dyar, M. Frascioni, M. A. Giesener, D. Benitez, E. Tkatchouk, O. Chernyashevskyy, W. H. Shin, H. Li, S. Sampath, C. L. Stern, A. A. Sarjeant, K. J. Hartlieb, Z. Liu, R. Carmieli, Y. Y. Botros, J. W. Choi, A. M. Slawin, J. B. Ketterson, M. R. Wasielewski, W. A. Goddard III, J. F. Stoddart, *Science* **2013**, *339*, 429–433; d) A. C. Fahrenbach, C. J. Bruns, H. Li, A. Trabolsi, A. Coskun, J. F. Stoddart, *Acc. Chem. Res.* **2014**, *47*, 482–493; e) C. J. Bruns, J. F. Stoddart, *Acc. Chem. Res.* **2014**, *47*, 2186–2199; f) E. J. Dale, N. A. Vermeulen, M. Juriček, J. C. Barnes, R. M. Young, M. R. Wasielewski, J. F. Stoddart, *Acc. Chem. Res.* **2016**, *49*, 262–273.
- [26] M. H. Tootoonchi, S. Yi, A. E. Kaifer, *J. Am. Chem. Soc.* **2013**, *135*, 10804–10809.
- [27] K_a calculation: since no obviously free guest signals were observed by diluting 1.0 mM CB[10]·2 down to 0.01 mM, we assume that in the solution the amount of free guest was less than 5%. A K_a larger than 10^7 M^{-1} can be obtained by calculation.
- [28] Crystal data for $(\text{C}_{60}\text{H}_{60}\text{N}_{40}\text{O}_{20})\cdot(\text{C}_{36}\text{H}_{32}\text{N}_4)_4\cdot 58\text{H}_2\text{O}$: $M_r = 3734.65$; rhombohedral; space group: $R\bar{3}c$; $a = 26.654(6)$, $b = 26.654(6)$, $c = 118.53(3)$ Å; $\alpha = \beta = 90^\circ$, $\gamma = 120^\circ$; $Z = 18$; $V = 72930(40)$ Å³; $\rho_{\text{calcd}} = 1.531 \text{ g cm}^{-3}$; $\mu = 0.877 \text{ mm}^{-1}$; $\theta_{\text{max}} = 16.50^\circ$; $F(000) = 34704$ e; reflections collected/unique, 57285/4338 ($R_{\text{int}} = 0.1403$); final R indices [$I > 2\sigma(I)$] $R_1 = 0.1504$, R indices (all data) $wR_2 = 0.3434$; GOF = 1.470 for all data. CCDC-1434842 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [29] A. R. Bernardo, J. F. Stoddart, A. E. Kaifer, *J. Am. Chem. Soc.* **1992**, *114*, 10624–10631.
- [30] W.-H. Huang, S. Liu, P. Y. Zavalij, L. Isaacs, *J. Am. Chem. Soc.* **2006**, *128*, 14744–14745.
- [31] a) A. L. Koner, C. Márquez, M. H. Dickman, W. M. Nau, *Angew. Chem. Int. Ed.* **2011**, *50*, 545–548; *Angew. Chem.* **2011**, *123*, 567–571; b) N. Baroah, M. Sundararajan, J. Mohanty, A. C. Bhasikuttan, *J. Phys. Chem. B* **2014**, *118*, 7136–7146.
- [32] M. Bria, G. Cooke, A. Cooper, J. F. Garety, S. G. Hewage, M. Nutley, G. Rabani, P. Woisel, *Tetrahedron Lett.* **2007**, *48*, 301–304.
- [33] a) M. Asakawa, W. Dehaen, G. Labbe, S. Menzer, J. Nouwen, F. M. Raymo, J. F. Stoddart, D. J. Williams, *J. Org. Chem.* **1996**, *61*, 9591–9595; b) C.-H. Sue, S. Basu, A. C. Fahrenbach, A. K. Shveyd, S. K. Dey, Y. Y. Botros, J. F. Stoddart, *Chem. Sci.* **2010**, *1*, 119–125.

Received: September 1, 2016

Published online on November 8, 2016