

## Peskin Solutions: Chapter 9

Jinchen

### I. HOW TO USE THE FUNCTIONAL METHOD TO GET PROPOGATOR.

According to (9.34), generating functional  $Z[J] = \int \mathcal{D}\phi [\exp(i \int d^4x \mathcal{L}) \cdot \exp(i \int d^4x J(x)\phi(x))]$ . Then change the variable to get  $Z[J] = \int \mathcal{D}\phi' [\exp(i \int d^4x \mathcal{L}') \cdot \exp(-\frac{i}{2} \int d^4x J \hat{O}^{-1} J)]$ . Here the current term is irrelevant to  $\phi'$ , so  $Z[J] = Z_0 \cdot \exp(-\frac{i}{2} \int d^4x J \hat{O}^{-1} J)$ , and the functional derivatives will be applied on the current term.

### II. PROBLEM 9.1

(a)

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^2 + (\partial_\mu \phi^* - ieA_\mu \phi^*)(\partial^\mu \phi + ieA^\mu \phi) - m^2 \phi^* \phi = \mathcal{L}_A + \mathcal{L}_\phi + \mathcal{L}_I$$

The  $\mathcal{L}_A$  is just free E-M field, so the propagator is the propagator of photon.

The  $\mathcal{L}_\phi = \partial_\mu \phi^* \partial^\mu \phi - m^2 \phi^* \phi = \partial_\mu (\phi^* \partial^\mu \phi) - \phi^* \partial_\mu (\partial^\mu \phi) - m^2 \phi^* \phi$ , because the differential term in the Lagrangian density makes no difference, we got  $\mathcal{L}_\phi = -\phi^* \partial_\mu (\partial^\mu \phi) - m^2 \phi^* \phi = \phi^* (-\partial^2 - m^2) \phi = \phi^* \hat{T} \phi$ .

With generating functional method, we have  $\mathcal{L}_\phi + \eta^* \phi + \phi^* \eta$  in the  $Z[J]$ , then do a shift  $\phi \rightarrow \phi' = \phi + \hat{T}^{-1} \eta$ , we got  $\mathcal{L}_\phi + \eta^* \phi + \phi^* \eta = \mathcal{L}_{\phi'} - \eta^* \hat{T}^{-1} \eta$ . If  $G$  is the Green function of  $\hat{T}$ , then  $\mathcal{L}_{\phi'} - \eta^* \hat{T}^{-1} \eta = \mathcal{L}_{\phi'} - \eta^* (iG * \eta)$ ,

$$Z[\eta, \eta^*] = Z_0 \cdot \exp[-i \int d^4x d^4y \eta^*(x) iG(x-y) \eta(y)]$$

$$\text{prop} = -\frac{\delta}{\delta \eta^*} \frac{\delta}{\delta \eta} \exp[-i \int d^4x d^4y \eta^*(x) iG(x-y) \eta(y)] = -G$$

After two functional derivatives, we will find the propagator is exactly the  $-G$ .

So the propagator of  $\phi$  and  $\phi^*$  is  $\frac{i}{p^2 - m^2 + i\epsilon}$ . (How to calculate the Green function of  $\hat{T}$  - [Check Eq.\(2.57\) in Peskin](#))

$$\hat{T}^{-1} \eta(x) = i \int d^4y G(x-y) \eta(y)$$

$$\hat{T} G(x-y) = (-\partial^2 - m^2) G(x-y) = -i \delta(x-y)$$

FT to get,

$$(p^2 - m^2) \tilde{G}(p) = -i$$

$$-\tilde{G}(p) = \frac{i}{p^2 - m^2}$$

Then comes to vertices,  $\mathcal{H}_I = -\mathcal{L}_I$  (P. 289 in Peskin), theoretically we should check Eq.(4.31) and do the contraction to get Feynman rules, but here we can just look at  $\exp[i \int \mathcal{L}_I]$ , here  $\mathcal{L}_I = ie g^{\mu\nu} (\partial_\mu \phi^* A_\nu \phi - A_\mu \phi^* \partial_\nu \phi) + e^2 g^{\mu\nu} A_\mu \phi^* A_\nu \phi$ , then  $i\mathcal{L}_I = -ie g^{\mu\nu} (-i\partial_\mu \phi^* A_\nu \phi + A_\mu \phi^* i\partial_\nu \phi) + ie^2 g^{\mu\nu} A_\mu \phi^* A_\nu \phi$ .

There are three terms, let's throw those fields away and turn  $i\partial\phi$  to  $p_\phi\phi$ ,  $-i\partial\phi^*$  to  $p_{\phi^*}\phi^*$ , here  $p$ 's are along particle/anti-particle lines, besides, the third term has two  $A$  fields, which are commutative, so there should be a factor 2 for the  $AA\phi^*\phi$  vertex.

So,

$$\text{For } \phi^* A \phi : -ie(p + p')^\mu$$

$$\text{For } AA\phi^*\phi : 2ie^2 g^{\mu\nu}$$

Theoretically,

$$\langle \phi\phi^* | S | \gamma \rangle = \langle \phi\phi^* | T \int d^4x i\mathcal{L}_I | \gamma \rangle = \langle \phi\phi^* | T \int d^4x (-ie) g^{\mu\nu} (-i\partial_\mu \phi^* A_\nu \phi + A_\mu \phi^* i\partial_\nu \phi) | \gamma \rangle$$

and

$$\langle \phi\phi^* | S | \gamma\gamma \rangle = \langle \phi\phi^* | T \int d^4x i\mathcal{L}_I | \gamma\gamma \rangle = \langle \phi\phi^* | T \int d^4x (ie^2) g^{\mu\nu} A_\mu \phi^* A_\nu \phi | \gamma\gamma \rangle$$

give the Feynman rules of two kinds of vertex with contractions.

(b)

With Eq.(4.84),  $m_e$  is ignored, then,

$$\left(\frac{d\sigma}{d\Omega}\right)_{c.m.} = \frac{|\vec{p}_\phi|}{32(2\pi)^2 E_e^2 \cdot 2E_e} \frac{1}{4} \Sigma |\mathcal{M}(ee \rightarrow \phi^* \phi)|^2$$

The outlines of  $\phi$  and  $\phi^*$  are 1, the Feynman diagram looks similar to the diagram in P.131.

$$i\mathcal{M} = (-ie)^2 \bar{v}(k_2) \gamma^\mu u(k_1) \frac{-ig_{\mu\nu}}{s + i\epsilon} (p_1 - p_2)^\nu = ie^2 \bar{v}(k_2) (\not{p}_1 - \not{p}_2) u(k_1) \frac{1}{s + i\epsilon}$$

$$\frac{1}{4} \sum_{\text{spin}} |\mathcal{M}|^2 = \sum_{\text{spin}} \frac{e^4}{4s^2} \bar{v}(k_2) (\not{p}_1 - \not{p}_2) u(k_1) \bar{u}(k_1) (\not{p}_1 - \not{p}_2) v(k_2)$$

$$= \sum_{\text{spin}} \frac{e^4}{4s^2} \text{tr}(v(k_2) \bar{v}(k_2) (\not{p}_1 - \not{p}_2) u(k_1) \bar{u}(k_1) (\not{p}_1 - \not{p}_2))$$

$$= \frac{e^4}{4s^2} \text{tr}(\not{k}_2 (\not{p}_1 - \not{p}_2) \not{k}_1 (\not{p}_1 - \not{p}_2))$$

$$= \frac{e^4}{4s^2} \left[ 8(k_1 \cdot p_1 - k_1 \cdot p_2)(k_2 \cdot p_1 - k_2 \cdot p_2) - 4(k_1 \cdot k_2)(p_1 - p_2)^2 \right]$$

Choose a specific frame,

$$k_1 = (E, 0, 0, E), \quad p_1 = (E, p \sin \theta, 0, p \cos \theta)$$

$$k_2 = (E, 0, 0, -E), \quad p_2 = (E, -p \sin \theta, 0, -p \cos \theta)$$

$$\frac{1}{4} \sum_{\text{spin}} |\mathcal{M}|^2 = \frac{e^4 p^2}{2E^2} \sin^2 \theta$$

$ee \rightarrow \mu\mu$  is [Eq.\(5.11\)](#)

So,

$$\left( \frac{d\sigma}{d\Omega} \right)_{\text{c.m.}} = \frac{1}{2(2E)^2} \frac{p}{8(2\pi)^2 E} \left( \frac{1}{4} \sum |\mathcal{M}|^2 \right) = \frac{\alpha^2}{8s} \left( 1 - \frac{m^2}{E^2} \right)^{3/2} \sin^2 \theta$$

(c)

Two diagrams because there are two kinds of vertex which are listed in (a). Because the minus signs in the  $\mathcal{L}_I$  are all absorbed in vertex, and there is no Fermion field, the sign between two vertices is  $+$ . That's why the two diagrams should be added.

$$i\Pi_1^{\mu\nu} = e^2 \int \frac{d^4 k}{(2\pi)^4} (2k+q)^\mu \frac{1}{k^2 - m^2 + i\epsilon} (2k+q)^\nu \frac{1}{(k+q)^2 - m^2 + i\epsilon}$$

$$i\Pi_2^{\mu\nu} = -2e^2 g^{\mu\nu} \int \frac{d^4 k}{(2\pi)^4} \frac{1}{(k+q)^2 - m^2 + i\epsilon}$$

add together, get

$$i\Pi^{\mu\nu} = -e^2 \int \frac{d^4 k}{(2\pi)^4} \frac{2g^{\mu\nu}(k^2 - m^2) - (2k+q)^\mu (2k+q)^\nu}{(k^2 - m^2)((k+q)^2 - m^2)}$$

$$\frac{1}{(k^2 - m^2)((k+q)^2 - m^2)} = \int_0^1 dx \frac{1}{[(k + (1-x)q)^2 + xq^2 - x^2q^2 - m^2]^2}$$

change the variable,  $l = k + (1-x)q$ , with [Eq.\(6.45\)](#),

$$\text{numerator} = g^{\mu\nu} l^2 + 2g^{\mu\nu} (1-x)^2 q^2 - 2g^{\mu\nu} m^2 - (2x-1)^2 q^\mu q^\nu$$

do the Wick rotation,  $l^0 = il_E^0$  and  $l^i = l_E^i$ , so we have  $d^4 l = id^4 l_E$  and  $l^2 = -l_E^2$ ,

$$i\Pi^{\mu\nu} = -ie^2 \int_0^1 dx \int \frac{d^4 l_E}{(2\pi)^4} \frac{-g^{\mu\nu} l_E^2 + 2g^{\mu\nu} (1-x)^2 q^2 - 2g^{\mu\nu} m^2 - (2x-1)^2 q^\mu q^\nu}{[l_E^2 + m^2 + x^2 q^2 - xq^2]^2}$$

$$= -ie^2 \int_0^1 dx \int \frac{d^4 l_E}{(2\pi)^4} \left[ \frac{-g^{\mu\nu} l_E^2}{(l_E^2 + \Delta)^2} + \frac{2g^{\mu\nu}(1-x)^2 q^2 - 2g^{\mu\nu} m^2 - (2x-1)^2 q^\mu q^\nu}{(l_E^2 + \Delta)^2} \right]$$

use dimensional regularization, with Eq.(7.85) and Eq.(7.86),

$$i\Pi^{\mu\nu} = -ie^2 \int_0^1 dx [(2g^{\mu\nu}(1-x)^2 q^2 - 2g^{\mu\nu} m^2 - (2x-1)^2 q^\mu q^\nu) \frac{1}{(4\pi)^{d/2}} \frac{\Gamma(2-\frac{d}{2})}{\Gamma(2)} \left(\frac{1}{\Delta}\right)^{2-\frac{d}{2}} \\ - g^{\mu\nu} \frac{1}{(4\pi)^{d/2}} \frac{d}{2} \frac{\Gamma(2-\frac{d}{2}-1)}{\Gamma(2)} \left(\frac{1}{\Delta}\right)^{2-\frac{d}{2}-1}]$$

$$= -ie^2 \int_0^1 dx \frac{1}{(4\pi)^{d/2}} \left(\frac{1}{\Delta}\right)^{2-d/2} \Gamma(2-d/2) [(2g^{\mu\nu}(1-x)^2 q^2 - (2x-1)^2 q^\mu q^\nu) - g^{\mu\nu} \frac{d}{2-d} (x^2 q^2 - x q^2)]$$

set  $d = 4 - \epsilon$  with  $\epsilon \rightarrow 0$ ,

$$i\Pi^{\mu\nu} = \frac{-ie^2}{(4\pi)^2} \int_0^1 dx \left(\frac{\epsilon}{2} - \log \Delta - \gamma + \log(4\pi)\right) [(g^{\mu\nu}(2x-2)(2x-1)q^2 - (2x-1)^2 q^\mu q^\nu)]$$

Because  $\int_0^1 dx (\frac{\epsilon}{2} - \log \Delta - \gamma + \log(4\pi))(2x-1) = \int_0^1 dx \frac{\epsilon}{2}(2x-1) = 0$ , we have

$$i\Pi^{\mu\nu} = \frac{-ie^2}{(4\pi)^2} \int_0^1 dx \left(\frac{\epsilon}{2} - \log \Delta - \gamma + \log(4\pi)\right) (2x-1)^2 [(g^{\mu\nu} q^2 - q^\mu q^\nu)]$$

with MS-bar scheme,

$$\Pi(q^2) = \frac{-\alpha}{4\pi} \int_0^1 dx (-\log \Delta) (2x-1)^2$$

If we adopt  $-q^2 \gg m^2$ ,

$$\Pi(q^2) = \frac{-\alpha}{4\pi} \int_0^1 dx (-\log(x-x^2) - \log(-q^2)) (2x-1)^2 \rightarrow \frac{-\alpha}{12\pi} \log(-q^2)$$

while looking at Eq.(7.90),  $\int_0^1 dx x(1-x) = \frac{1}{6}$ , we know for  $e+e-$  pair,

$$\Pi(q^2) \rightarrow \frac{-\alpha}{3\pi} \log(-q^2)$$

which is four times as our results.

### III. PROBLEM 9.2

(a)

---