



On each attack day, write the intensity from 1 to 10 at the bottom of the box and tick “Med” if you took an acute medication. You can also jot down a word in the box: period day, bad night, stress...

More than 10 days with acute medication this month? Talk to your doctor: that is the medication-overuse threshold.



Attack log

One row per attack. For each medication or relief, tick whether it helped. Print this page as many times as you need.

Date	Start and end	Intensity	Symptoms	Possible triggers	Medication and time	Drug-free relief
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No
		/10			Helped: ☐ Yes ☐ A bit ☐ No	Helped: ☐ Yes ☐ A bit ☐ No

Common triggers Stress · Lack of sleep · Skipped meal · Period · Weather · Screens · Alcohol · Dehydration · Caffeine · Bright light

Common symptoms Nausea · Light sensitivity · Noise sensitivity · Visual aura · Dizziness · Throbbing pain

Common relief Rest in the dark · Cold on the forehead · Slow breathing · Sleep · Drinking water · Eating a little



Mellow Take back control of your migraines with the app

mellowmigraine.com