

Universidade de São Paulo  
Instituto de Física

# Vínculos Cosmológicos com o Auxílio do Efeito Sachs-Wolfe Integrado

Arthur Diniz Meirelles

Orientador: Prof. Dr. Edivaldo Moura Santos  
Coorientador: Prof. Dr. Ronaldo Carlotto Batista

Dissertação de mestrado apresentada ao Instituto de Física da Universidade de São Paulo, como requisito parcial para a obtenção do título de Mestre(a) em Ciências.

Banca Examinadora:

Prof(a). Dr. Edivaldo Moura Santos - Orientador (IF-USP)

Prof(a). Dr(a). Nome do(a) Professor(a) (instituição de trabalho)

Prof(a). Dr(a). Nome do(a) Professor(a) (instituição de trabalho)

São Paulo  
2024

## FICHA DA BIBLIOTECA

University of São Paulo  
Physics Institute

# Cosmological Constraints using the Sachs-Wolfe Integrated Effect

Arthur Diniz Meirelles

Supervisor: Prof. Dr. Edivaldo Moura Santos  
Co-supervisor: Prof. Dr. Ronaldo Carlotto Batista

Dissertation submitted to the Physics Institute of the  
University of São Paulo in partial fulfillment of the  
requirements for the degree of Master of Science.

Examining Committee:

Prof. Dr. Edivaldo Moura Santos - Supervisor (IF-USP)

Prof. Dr. Name (institution)

Prof. Dr. Name (institution)

São Paulo  
2024

*Frase bonita*

# Acknowledgements

testando

testando mais

# Abstract

ABSTRACT AQUI

**Keywords:**

# Resumo

RESUMO AQUI

**Palavras-chave:**

# List of Figures

1.1	Each image represents the same 10 square degree patch of the sky measured by the three telescopes shown above each image, the improvement in the resolution of each map is very noticeable. Credit: NASA/LAMBDA Science Team. . . . .	2
2.1	A representation of the time evolution of the Universe, starting from the left (the Big Bang) with time passing towards the right. The radius of each section indicates the size of the Universe at that time; during inflation this radius increases very rapidly, indicating an accelerated expansion during this period, while closer to the present the slope of increase in size shows a slight acceleration, indicating the current dark energy accelerated period. Briefly after the inflation period (in cosmological time), the photons decouple from baryons, leaving a radiation field in the Universe, resulting in the CMB, represented in the image by the afterglow of inflation. After this, the only radiation in the Universe is this afterglow for many years, during that time structure starts to form from the small inhomogeneities left by quantum fluctuations in the very early stages of the Universe, forming the first stars and then the first galaxies. Credit: NASA/WMAP Science Team. . . . .	11
2.2	Planck 2018 CMB temperature power spectrum: the blue line corresponds to a $\Lambda$ CDM cosmology best-fit to the data points in orange. For $\ell < 30$ a logarithmic scale was used and the data points correspond to the $D_\ell^{TT}$ for each value of $\ell$ , while for $\ell \geq 30$ a linear scale was used and the data points correspond to binned values for the data for $D_\ell^{TT}$ . The error bars show $\pm 1\sigma$ uncertainties, including cosmic variance. Extracted from [6]. . . . .	18
2.3	Matter power spectrum calculated as a function of $k/h$ , where $h = H_0/(100 \text{ km/s/Mpc})$ , using the HALOFIT model with cosmological parameters according to Planck's best fit results[6]. . . . .	21
2.4	CMB autocorrelation comparison for the late-ISW term and the full spectrum. Calculations were made using CAMB (for the full spectrum) and codes developed during the project. The parameters used are those corresponding to the Planck best fit parameters[6]. . . . .	22
2.5	Galaxy autocorrelation spectrum (left) and late-ISW contribution to the galaxy-CMB cross-correlation (right). The selection function used to calculate these spectra was parametrized to be compatible with 2MASS' galaxy redshift distribution[16, 44]. . . . .	23
2.6	The solid lines in the plots correspond to theoretical predictions for the CMB temperature autocorrelation spectra corresponding to different parameter values of the dark energy equation of state, the black data points correspond to Planck's data[6]. Figures 2.6a, 2.6b and 2.6c show how the spectrum changes for a constant value of $w_a$ shown on the top left of the axis and varying speeds of sound, while the other three figures show how the curve behaves with varying values of $w_a$ and a constant speed of sound. All theoretical calculations were done using CAMB. . . . .	25



LIST OF FIGURES

---

3.1	Selection function calculated for the 4 bands of the 2MASS catalog according to equation (3.1). . . . .	27
3.2	Selection function calculated for various parameter values according to the parametrization (3.1). Each plot is made with constant values for two parameters, corresponding to the band 2 of the 2MASS catalog (see Table 3.1), and the parameter on the right is varied. . . . .	27

# List of Tables

2.1	Equation of state parameters $w$ , energy density functions $\rho(a)$ and energy density fractions for each known component of the Universe according to the $\Lambda$ CDM model. Most data were adapted from Planck[6], the radiation data was calculated based on [28]. . . . .	10
2.2	Marginalized values and 68% confidence limits for cosmological parameters, assuming parametrization (2.53), obtained by combining Planck temperature and polarization autocorrelation and cross-correlation spectra[50], type IA supernovae data from the Joint Light Curve Analysis (JLA)[51] and BAO data from the Sloan Digital Sky Survey (SDSS)[52]. The $\Delta\chi^2$ value was computed with respect to the $\Lambda$ CDM best fits computed from the same data set combination. .	24
3.1	Parameter values for the 4 bands of the 2MASS catalog. Values extracted from [39]. . . . .	26

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Cosmology's Standard Model</b>	<b>4</b>
2.1	The Homogeneous Background . . . . .	4
2.1.1	Fundamental Observations and Principles . . . . .	4
2.1.2	Friedmann Equations . . . . .	6
2.1.3	Energy Content . . . . .	7
2.1.4	A Brief History of the Universe . . . . .	9
2.2	Inhomogeneities and Anisotropies . . . . .	11
2.2.1	Perturbed Space-Time . . . . .	12
2.2.2	The Boltzmann Equation . . . . .	13
2.3	The Cosmic Microwave Background . . . . .	16
2.3.1	CMB Autocorrelation Spectrum . . . . .	16
2.3.2	Analytical Approximation . . . . .	18
2.4	The Integrated Sachs Wolfe Effect . . . . .	20
2.4.1	The Matter Power Spectrum . . . . .	20
2.4.2	The Galaxy-CMB Correlation Spectra . . . . .	20
2.4.3	Dependence on Dark Energy Models . . . . .	23
<b>3</b>	<b>Optimizing the Selection Function</b>	<b>26</b>
3.1	Parametrization of the Selection Function . . . . .	26
3.2	Maximizing the Cross-Correlation Signal . . . . .	28
<b>4</b>	<b>Extracting Correlations from Data</b>	<b>30</b>
4.1	Two Point Functions . . . . .	30
4.2	Gibbs Sampling . . . . .	31
4.3	Data Sets and Analysis . . . . .	31
<b>5</b>	<b>Results</b>	<b>32</b>
<b>6</b>	<b>Conclusions</b>	<b>33</b>
<b>A</b>	<b>Fourier Space</b>	<b>35</b>
<b>B</b>	<b>Monte Carlo Markov Chains</b>	<b>36</b>

# 1 | Introduction

The discovery of the Cosmic Microwave Background (CMB) is a fine example of serendipity, Arno A. Penzias and Robert W. Wilson found an unexpected excess temperature while measuring the effective zenith noise temperature in 1964, and after some investigation, they attributed this excess temperature to an isotropic radiation field that today we call the CMB [1]. The CMB has since been used as one of the main probes to studying cosmology, being able to provide various tests and insights into cosmological models [2–4] and measurements of cosmological parameters [5, 6].

In the standard cosmological model, commonly called the  $\Lambda$ CDM model, the CMB is a radiation field that has permeated the Universe since the Big Bang, its photons have been freely traveling the Universe since their decoupling from electrons and protons during the very early stages of the Universe’s history [7]. The current measurement for the black body spectrum temperature of the CMB is 2.725 48(57) K [8], and despite being isotropic to many orders of precision, it does contain anisotropies in the order of  $10^{-5}$  K, which is expected in the  $\Lambda$ CDM model [9]. These fluctuations are the result of many physical effects that contribute to the CMB’s emission or are acting on the photons as they travel through the Universe, making them an excellent probe to constrain cosmological parameters.

Some mappings of the CMB were carried out in the last few decades, the first satellite launched specifically dedicated to doing this was the Cosmic Background Explorer (COBE) satellite with a  $7^\circ$  angular resolution [10], followed by the Wilkinson Microwave Anisotropy Probe (WMAP) with a  $0.3^\circ$  angular resolution, which was also able to show that CMB photons are polarized, a property related to other physical effects that assisted cosmologists in further understanding the history of the Universe[11–13]. In 2009, the Planck Satellite was launched, improving upon WMAP’s precision to an impressive value of  $5'$  ( $\sim 0.083^\circ$ ) angular resolution [14], allowing cosmologists to access the finest details in the CMB anisotropies to this day. Figure 1.1 shows a comparison between the resolution of these maps.

Amongst the many effects that affect the CMB, the Integrated Sachs-Wolfe (ISW) Effect is one of the most important ones. First described by Rainer K. Sachs and Arthur M. Wolfe in 1967, the ISW effect is a mechanism that causes photons (including CMB photons) to blueshift



Figure 1.1: Each image represents the same 10 square degree patch of the sky measured by the three telescopes shown above each image, the improvement in the resolution of each map is very noticeable. Credit: NASA/LAMBDA Science Team.

in cosmological periods when gravitational potentials change rapidly. One such period is the time between the current dark energy dominated era and the dark matter dominated era before it. During this transition, gravitational potentials decay (or "flatten") over time, causing CMB photons that entered potential wells with a certain gravitational blueshift to leave these wells with a redshift that does not compensate for the initial blueshift, leaving a net blueshift in these photons, creating patterns that can be observed in the CMB temperature map and spectrum[15].

#### ILUSTRAÇÃO DO ISW?

There are two contributions to the ISW effect: The early-time ISW effect occurs immediately after the last scattering surface (COMPLEMENTAR AQUI DEPOIS), and the late-time ISW effect, which is more recent, occurring in the current dark energy dominated era.

From this association between blueshifted CMB photons and gravity wells, it is natural to assume that the matter distribution of the Universe should hold a statistical cross-correlation between with the CMB temperature map. When looking at the theoretical model, we will see that this cross-correlation is expected, and many works in the past have already searched and found such cross-correlation (EXAMPLE REFS) (COMPLETAR AQUI, VOU COMPLEMENTAR NISSO? APRESENTAR ALGO NOVO? ETC).

The present work expands on a method (what characterizes this method and sets it apart from other ones?) presented in [16] to estimate these cross-correlations, using the models and some techniques presented there, with a more thorough selection of the data that is to be used in

the analysis, using the newer Planck Satellite data instead of the WMAP data, and applying the method to alternative dark energy models, making it possible to compare them with the  $\Lambda$ CDM model and providing insights about the nature of dark energy.

The rest of this dissertation will be organized as follows: DESCRIBE HOW THE SECTIONS ARE ORGANIZED WHEN THE REST OF THE DISSERTATION IS DONE.

## 2 | Cosmology's Standard Model

The advent of general relativity in the early 20th century allowed physicists to conceptualize mathematical models of the Universe. Albert Einstein proposed a model for a static Universe using his theory[17], and other Universe models were proposed in the following years. To start this chapter, the fundamental principles of cosmology will be discussed and used to setup a simple background for the Universe, and the specificities of the currently most accepted model for the Universe – the  $\Lambda$ CDM model – will be discussed. After this, we will complement this simple background with finer details and provide a statistical description of cosmological observables that can be used to test our models.

### 2.1 The Homogeneous Background

#### 2.1.1 Fundamental Observations and Principles

One of the fundamental principles of cosmology is called the Cosmological Principle: at large scales, the Universe is both homogeneous and isotropic. This principle can be used to simplify the description of the energy distribution in the Universe, but it also simplifies the possible metrics that can describe the Universe to those that have constant curvature (a quantity denoted as  $k$ ), and it can be shown that any space with constant curvature is locally isometric to: The Euclidian flat space ( $k = 0$ ), a hyperspherical space ( $k = +1$ ) or a hyperbolic space ( $k = -1$ )[18]. The Cosmological Principle has support from a good amount of cosmological observations, such as the CMB[10], the galaxy sky distribution[19, 20], amongst others.

In the period from 1927 to 1929, Edwin Hubble and Georges Lemaître, amongst other scientists, greatly contributed to the current understanding of cosmology: Observations were carried out and showed that most galaxies and nebulae data show a redshift proportional to their distances relative to the earth[21], these results were studied using general relativity and cosmological models at the time[22]. It was then established another principle that physicists were avoiding in their theoretical models for quite some time: The Universe is not static, in fact it is expanding, and for objects not too far (REFERÊNCIA), there is a linear relationship between the redshift of an observed object and its distance to the observer.

To describe the Universe, a coordinate system is embedded in the entire Universe, and to describe its dynamical nature, the scale factor  $a$  is introduced to the theoretical model:  $a$  is set to 1 at the present as a convention (usually called  $a_0$ ), and its value at any given time  $a(t)$  is the size of the Universe relative to its present size. A simple relation can be established between the scale factor of an observed object and its redshift  $z$ :

$$1 + z = \frac{\lambda_{\text{obs}}}{\lambda_{\text{emit}}} = \frac{a_{\text{obs}}}{a_{\text{emit}}} = \frac{1}{a_{\text{emit}}} \quad (2.1)$$

where  $\lambda$  and  $a$  are the wavelength of the radiation and the scale factor, respectively, when emitted ( $\lambda_{\text{emit}}$  and  $a_{\text{emit}}$ ) and when observed ( $\lambda_{\text{obs}}$  and  $a_{\text{obs}}$ ).

$$H(t) = \frac{1}{a} \frac{da}{dt} \quad (2.2)$$

These properties are not exclusive to a particular cosmological model, instead being present in all of them. The standard model of cosmology, called  $\Lambda$ CDM, usually takes a few assumptions on top of the ones described previously:

- The Universe emerged from a space-time singularity that started expanding several billion years ago, in an event called the Big Bang;
- Aside from baryonic matter, radiation and neutrinos, there are two other main components in the Universe: non-relativistic (cold) dark matter and an energy component with constant density called dark energy, which is related to a cosmological constant  $\Lambda$ ;
- During the early stages of the Universe's history, it went through a period of rapid acceleration driven by a scalar quantum field.

Based on the Cosmological Principle, most theoretical models first set up a homogeneous and isotropic background description of the Universe, and then introduce perturbations to this background, allowing for a detailed statistical description of its evolution. In the rest of this section, we will use the principles and assumptions just presented to set this background.



### 2.1.2 Friedmann Equations

As already mentioned, the main theoretical framework to describe the Universe is the theory of general relativity, since the most relevant force in the scale of the Universe is the gravitational one. We then use Einstein's field equations with a cosmological constant  $\Lambda$ :

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad (2.3)$$

where  $R_{\mu\nu}$  is the Ricci curvature tensor for the Universe's metric  $g_{\mu\nu}$ ,  $R \equiv g_{\mu\nu}R^{\mu\nu}$  is the Ricci scalar,  $G$  is the gravitational constant and  $T_{\mu\nu}$  is the energy-momentum tensor.

In equation (2.3) and throughout this text (unless otherwise stated), we will use units such that the Boltzmann constant ( $k_b$ ), the speed of light in a vacuum ( $c$ ) and the reduced Planck constant ( $\hbar$ ) are all equal to one. Furthermore, Greek letter tensor indices indicate that the time component of the tensor is included in the equation and Latin letter tensor indices indicate that only spatial components are included. Current results strongly indicate the metric  $g_{\mu\nu}$  of the Universe to be flat ( $k = 0$ )[6], thus justifying the use of an Euclidian metric scaled by  $a(t)$ , and we will assume a homogeneous and isotropic distribution of energy throughout the Universe to construct an energy-momentum tensor with constant energy density  $\rho_0$  and constant pressure  $P$ [23]:

$$g_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & a^2 & 0 & 0 \\ 0 & 0 & a^2 & 0 \\ 0 & 0 & 0 & a^2 \end{pmatrix} \quad T_{\mu\nu} = \begin{pmatrix} -\rho_0 & 0 & 0 & 0 \\ 0 & P & 0 & 0 \\ 0 & 0 & P & 0 \\ 0 & 0 & 0 & P \end{pmatrix}. \quad (2.4)$$

The metric  $g_{\mu\nu}$  described in equation (2.4) is called the Friedmann-Lemaître-Robertson-Walker (FLRW) metric[24]. In order to expand any equation from (2.3), it is necessary to determine  $R = g^{\mu\nu}R_{\mu\nu}$ . By expanding the expression using the matrices defined in (2.4), the result is

$$R = 6 \left[ \frac{\ddot{a}}{a} + \left( \frac{\dot{a}}{a} \right)^2 \right]. \quad (2.5)$$

From all the equations in (2.3), if the one corresponding to  $\mu = \nu = 0$  is expanded using the

result (2.5), one obtains

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{1}{3}(8\pi G\rho_0 + \Lambda), \quad (2.6)$$

which relates the relative rate at which the Universe is expanding with the energy content of the Universe. In the  $\Lambda$ CDM model,  $\Lambda$  is assumed to be the source of the current accelerated expansion of the Universe, so an energy component related to this constant is necessary, and its energy density can be defined as

$$\rho_\Lambda = \frac{\Lambda}{8\pi G}, \quad (2.7)$$

such that

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho, \quad (2.8)$$

where  $\rho = \rho_\Lambda + \rho_0$ . This is called Friedmann's first equation.

When any of the space-space equations of (2.3) is expanded and the resulting equation is combined with equation (2.8), Friedmann's second equation is obtained:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3P). \quad (2.9)$$

This equation is a relation between the acceleration of the Universe and its energy content. It is clear that  $P < -\frac{1}{3}\rho$  indicates a Universe expanding with a positive acceleration, which is highly unexpected since gravity is an attractive force, but in section 2.1.3 the case of the cosmological constant will be discussed, in which  $\Lambda$  yields  $P_\Lambda = -\rho_\Lambda$ , being the reason it contributes to an accelerated expansion.

### 2.1.3 Energy Content

To understand the contribution of each component of the Universe to its dynamics, a differential equation for each component has to be determined, we do this using the energy-momentum conservation equation:

$$\nabla_\mu T_\nu^\mu \equiv \partial_\mu T_\nu^\mu + \Gamma_{\alpha\mu}^\mu T_\nu^\alpha - \Gamma_{\nu\mu}^\alpha T_\alpha^\mu = 0. \quad (2.10)$$

Again using the energy-momentum tensor defined in (2.4), the following equation is obtained:

$$\frac{\partial \rho}{\partial t} + \frac{\dot{a}}{a}(3\rho + 3P) = 0. \quad (2.11)$$

To solve this, a relation between density and pressure for each component of the Universe is needed, an equation of state. It is commonly assumed that, for a given component  $s$  of the Universe, its equation of state takes the form

$$P_s = w_s \rho_s, \quad (2.12)$$

where  $w_s$  is called the equation of state parameter. Using that equation and solving for  $\rho_s(a)$  on equation (2.11) yields

$$\rho_s(a) \propto \exp \left[ -3 \int_1^a \frac{da'}{a'} [1 + w_s(a')] \right]. \quad (2.13)$$

This solution ignores further interaction between components, it would apply to a single-component Universe, but it is also a useful first approximation in epochs of dominance of a single component. In the  $\Lambda$ CDM model, it is assumed that all components of the Universe possess a constant value for  $w_s$ , in that case the solution is simple:

$$\rho_s(a) \propto a^{-3(1+w_s)}. \quad (2.14)$$

This represents the evolution of the density parameter of any component as a function of the scale parameter, and thus as a function of cosmological time, since  $a(t)$  is monotonically increasing (the Universe has not reduced at any point in its history so far).

The rigorous form of determining  $w_s$  for some components of the Universe requires the usage of statistical physics. In cases where we have limited information on the physical properties of a certain component, such as dark energy, theoretical frameworks and assumptions have to be used, as well as the analysis of data.

In the case of dark energy, assuming its origin is related to the cosmological constant, the data strongly suggests a value of  $w_\Lambda = -1$ , which also indicates a constant density for dark energy. From the quantum field theory point of view, the cosmological constant energy density would be associated to zero point energy of a quantum field. However, so far, no calculation has been

successful in predicting a zero point energy whose density is consistent with the cosmological observations[25]. Many models propose alternative possibilities, such as dynamical dark energy in which  $w_{DE}$  is a function of  $a$ [26], which would still be able to explain the accelerated expansion of the Universe as long as  $w_{DE} < -\frac{1}{3}$ , these are discussed in more detail in section 2.4.3. Table 2.1 summarizes the equation of state properties of each known component in the Universe.

It is very useful to also define the energy density fraction of each component of the Universe using

$$\Omega_s = \frac{\rho_s(a_0)}{\rho_{\text{crit}}}, \quad (2.15)$$

where  $\rho_{\text{crit}}$  is the so called critical density, defined as

$$\rho_{\text{crit}} \equiv \frac{3H_0^2}{8\pi G} \approx 8.5 \times 10^{-27} \text{ kg/m}^3, \quad (2.16)$$

where  $H_0$  is the Hubble constant, the value of the Hubble parameter  $H(t)$  defined in (2.2) at the present.

This energy density makes it possible to determine what is the curvature of the Universe: if the total sum of energy densities in the Universe is equal to the critical density, the Universe is flat; if this sum is higher than  $\rho_{\text{crit}}$  the Universe would have positive curvature; and if this sum is lower than  $\rho_{\text{crit}}$ , the Universe would have negative curvature[27].

Table 2.1 summarizes the current estimated values for these energy fractions for the Universe in the present. It is important to note that commonly in cosmology (and throughout this text), the classification of baryons includes electrons, even though, according to the standard model of particle physics, electrons belong, together with their associated neutrinos, to one of the three families of leptons.

### 2.1.4 A Brief History of the Universe

We commonly categorize different stages of the Universe according to important events that happened during its history. The  $\Lambda$ CDM model assumes the Universe started as a singularity; as it started expanding, quantum fluctuations at that time would eventually result in inhomogeneities thought to be the source for the formation of structures.

Component	$\rho(a)$	$w$	$\Omega$
Baryonic matter	$\rho_0 a^{-3}$	0	0.05
Cold dark matter	$\rho_0 a^{-3}$	0	0.26
Radiation	$\rho_0 a^{-4}$	$\frac{1}{3}$	$5.29 \times 10^{-5}$
Dark energy	$\rho_0$	-1	0.68
Curvature	$\rho_0 a^{-2}$	$-\frac{1}{3}$	0.001

Table 2.1: Equation of state parameters  $w$ , energy density functions  $\rho(a)$  and energy density fractions for each known component of the Universe according to the  $\Lambda$ CDM model. Most data were adapted from Planck[6], the radiation data was calculated based on [28].

The next important period of the Universe's evolution was a highly accelerated expansion epoch driven by an effect called cosmic inflation, first proposed as a solution to the horizon problem[29, 30], related to how can the CMB temperature differences across the sky be so small ( $\sim 10^{-5}$  K), even between points apparently causally disconnected during all epochs of the Universe's thermal history. The  $\Lambda$ CDM model assumes inflation was driven by a quantum scalar field[23]. Definitive evidence of the inflationary model may be reached with the detection of a certain type of polarization of the CMB called the B-modes, which have not been detected yet, but have reached good constraints.[13]

Briefly after inflation, baryons and photons are still coupled through Compton scattering, thus photons are still unable to travel freely. The Universe at this point is composed of photons coupled with baryons, helium nuclei, and trace amounts of lithium. With the decrease in temperature, hydrogen atoms can form without being quickly ionized, Compton scattering eventually can no longer maintain baryons and photons coupled, resulting in a decoupling of these components. The photons are released to travel across the Universe, the radiation field formed from this period is what we currently measure as the CMB.

After this, an epoch called "the Dark Ages" starts, in which the photons in the Universe are the result of the decoupling described before and stars are yet to be formed, which is the reason why it is called a dark age. Matter dominates this period, greatly decreasing the rate of expansion of the Universe. As a result of the quantum fluctuations in the early Universe, stars, galaxies, and other structures form. The study of the formation of structure is completely dependent on the study of the inhomogeneities of the Universe at small scales, where the cosmological principle is no longer expected to be valid. Some details of this theoretical framework will be covered in the next section.

Figure 2.1 shows a summarized and visual representation of some of the events described in

this section.




Figure 2.1: A representation of the time evolution of the Universe, starting from the left (the Big Bang) with time passing towards the right. The radius of each section indicates the size of the Universe at that time; during inflation this radius increases very rapidly, indicating an accelerated expansion during this period, while closer to the present the slope of increase in size shows a slight acceleration, indicating the current dark energy accelerated period. Briefly after the inflation period (in cosmological time), the photons decouple from baryons, leaving a radiation field in the Universe, resulting in the CMB, represented in the image by the afterglow of inflation. After this, the only radiation in the Universe is this afterglow for many years, during that time structure starts to form from the small inhomogeneities left by quantum fluctuations in the very early stages of the Universe, forming the first stars and then the first galaxies. Credit: NASA/WMAP Science Team.

## 2.2 Inhomogeneities and Anisotropies

The existence of structures – nebulae, stars, galaxies, clusters – in the Universe indicates that, throughout its history, it has been inhomogeneous and anisotropic at some scale. These perturbations to the uniform background are relatively small though, take the CMB for instance: CMB anisotropies are around 5 orders of magnitude lower than its average blackbody temperature[5].

In this scenario, there is no known theoretical model that can predict the exact behavior of every particle in the Universe. Therefore, we turn to a statistical prediction taking into account

the properties of space-time and the components of the Universe. In this section, the first order equations that govern these properties will be shown by introducing perturbations both to the metric and to the stress-energy tensor.

### 2.2.1 Perturbed Space-Time

In the  $\Lambda$ CDM model, to account for perturbations in the uniform background previously set, we use the following new form for the metric:

$$\begin{cases} g_{00} = -1 - 2\Psi(\mathbf{x}, t), \\ g_{0i} = g_{i0} = 0, \\ g_{ij} = a^2(t)\delta_{ij}[1 + 2\Phi(\mathbf{x}, t)]. \end{cases} \quad (2.17)$$

In equation (2.17),  $\Psi$  corresponds to the Newtonian potential and governs the movement of nonrelativistic objects[23], while  $\Phi$  is a perturbation to the spatial curvature.

With the use of the geodesic equation on this new tensor and using a first order approximation, it is possible to obtain the following results for the motion equations of particles traveling through the perturbed space-time:

$$\frac{dx^i}{dt} = \frac{\hat{p}_i}{a} \frac{p}{E} (1 - \Phi + \Psi), \quad (2.18)$$

$$\frac{dp^i}{dt} = -(H + \dot{\Phi})p^i - \frac{E}{a}\partial_i\Psi - \frac{1}{a}\frac{dp^i}{dE}p^k\partial_k\Phi + \frac{p^2}{aE}\partial_i\Phi. \quad (2.19)$$

We can use

$$\frac{dp}{dt} = \frac{d}{dt}\sqrt{\delta_{ij}p^ip^j} = \delta_{ij}\frac{p^i}{p}\frac{dp^j}{dt} \quad (2.20)$$

and  $\hat{p}^i = p^i/p$  in combination with equation (2.19) to find separate equations for  $p$  and  $\hat{p}^i$ :

$$\frac{dp}{dt} = -[H + \dot{\Phi}]p - \frac{E}{a}\hat{p}^i\partial_i\Psi, \quad (2.21)$$

$$\frac{d\hat{p}^i}{dt} = \frac{E}{ap}[\delta^{ik} - \hat{p}^i\hat{p}^k]\partial_k\left(\frac{p^2}{E^2}\Phi - \Psi\right). \quad (2.22)$$

As for the dynamics of the field perturbations, it is possible to use Einstein's field equations

to obtain, in Fourier space (see Appendix A), the equations for scalar perturbations on the gravitational potentials[23]

$$k^2\Phi + 3H(\dot{\Phi} - H\Psi) = -4\pi Ga^2(\rho_{DM}\delta_{DM} + \rho_b\delta_b + 4\rho_\gamma\Theta_0 + 4\rho_\nu\mathcal{N}_0), \quad (2.23)$$

$$k^2(\Phi + \Psi) = -32\pi Ga^2(\rho_\gamma\Theta_2 + \rho_\nu\mathcal{N}_2). \quad (2.24)$$

Here,  $\rho$  indicates the average density of a component,  $\delta$  represents the perturbation to these densities, and the indexes  $DM$ ,  $b$ ,  $\gamma$  and  $\nu$  indicate the quantities related to dark matter, baryons, radiation and neutrinos respectively. The terms  $\Theta_0$ ,  $\Theta_2$ ,  $\mathcal{N}_0$  and  $\mathcal{N}_2$  are obtained in the calculation of the perturbed energy-momentum tensor, they are defined as

$$F_\ell = \frac{1}{(-i)^\ell} \int_{-1}^1 \frac{\mathcal{P}_\ell(\mu)F(\mu)}{2} d\mu, \quad (2.25)$$

where  $F$  can either represent  $\Theta$  or  $\mathcal{N}$ ;  $\mu = \hat{k} \cdot \hat{p}$  is the cosine between the momentum direction of the photon and the Fourier space wavevector;  $\mathcal{P}_\ell$  are Legendre polynomials; the functions  $\Theta(\mu)$  and  $\mathcal{N}(\mu)$  are dimensionless fields that quantify the perturbations of photon and neutrino distributions with respect to the homogeneous background; and  $i$  is the imaginary unit  $i^2 = -1$ .

### 2.2.2 The Boltzmann Equation

The Boltzmann equation is a fundamental component in the study of every component of the Universe. One general form of writing the Boltzmann equation is

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x^i} \cdot \frac{dx^i}{dt} + \frac{\partial f}{\partial p} \frac{dp}{dt} + \frac{\partial f}{\partial \hat{p}^i} \cdot \frac{d\hat{p}^i}{dt} = C[f], \quad (2.26)$$

where  $f$  is the distribution function of the component to be analyzed and  $C[f]$  is the collision term, a factor that is calculated by considering the interactions of such components.

Using equations (2.21) and (2.22) with the Boltzmann equation and assuming  $E = p$ , i.e. assuming a relativistic particle such as photons, the full first order Boltzmann equation for photons is



$$\begin{aligned} \frac{df}{dt} = & \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x^i} \frac{\hat{p}^i}{a} (1 - \Phi - \Psi) - \frac{\partial f}{\partial p} \left\{ [H + \dot{\Phi}]p + \frac{1}{a} \hat{p}^i \partial_i \Psi \right\} \\ & + \frac{\partial f}{\partial \hat{p}^i} \frac{1}{a} \left[ \partial_i (\Phi - \Psi) - \hat{p}^i \hat{p}^k \partial_k (\Phi - \Psi) \right]. \end{aligned} \quad (2.27)$$

The first order approximation is being used, and so it is possible to simplify this expression further. For instance, the distribution  $f$  for the homogeneous background takes the form of Bose-Einstein's distribution, which means there is no dependence on  $\hat{p}^i$  and  $x^i$  in this case. It is commonly assumed that the deviations from this distribution are of the same order of the perturbations  $\Phi$  and  $\Psi$ , so both  $\frac{\partial f}{\partial x^i}$  and  $\frac{\partial f}{\partial \hat{p}^i}$  are first order factors, multiplied by  $\Phi$  or  $\Psi$  result in second order terms which can be neglected, resulting in

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \frac{\hat{p}^i}{a} \frac{\partial f}{\partial x^i} - \left[ H + \dot{\Phi} + \frac{1}{a} \hat{p}^i \partial_i \Psi \right] p \frac{\partial f}{\partial p}. \quad (2.28)$$

This is an important equation for the description of the CMB, but the description of anisotropies for massive particles is still important, for which the same assumptions are made, resulting in

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \frac{p}{Ea} \hat{p}^i \frac{\partial f}{\partial x^i} - \left[ H + \dot{\Phi} + \frac{E}{ap} \hat{p}^i \partial_i \Psi \right] p \frac{\partial f}{\partial p}. \quad (2.29)$$

The only difference between these equations is the term  $E/p$  which, in the case of radiation, is equal to 1. The first order approximation for non-relativistic matter breaks down in the late Universe, a non-linear model is necessary in this scenario.

To arrive at a differential equation, a function to describe the perturbations needs to be defined, and so does the distribution function. Taking the CMB as an example, the perturbed form of the temperature distribution for CMB photons in general can be defined as

$$T(\mathbf{x}, \hat{\mathbf{p}}, t) = \bar{T}(t)[1 + \Theta(\mathbf{x}, \hat{\mathbf{p}}, t)], \quad (2.30)$$

where  $\bar{T}$  is the average temperature in a given time and  $\Theta$  is the same field presented in equation (2.25), that quantifies the perturbations of the CMB distribution with respect to its average value. This perturbation field can be expressed in terms of the set of functions  $\Theta_\ell$  defined in (2.25) by the following equation in Fourier space:

$$\Theta(\hat{\mathbf{k}}, \mu) = \sum_{\ell=0}^{\infty} (2\ell+1)(-i)^\ell \Theta_\ell(\hat{\mathbf{k}}) \mathcal{P}_\ell(\mu). \quad (2.31)$$

If we use the Bose-Einstein distribution for the temperature defined in (2.30) and expand it to its first order, we obtain

$$f(\mathbf{x}, \mathbf{p}, t) = f^{(0)}(p, t) - p \frac{\partial f^{(0)}(p, t)}{\partial p} \Theta(\mathbf{x}, \mathbf{p}, t), \quad (2.32)$$

where

$$f^{(0)}(p, t) = \left[ \exp\left(\frac{p}{T}\right) - 1 \right]^{-1} \quad (2.33)$$

We can then expand the Boltzmann equation using the results in (2.32) and (2.33) and take into account Compton scattering to calculate the collision term  $C[f]$  to obtain a differential equation for the CMB photons. We can use similar procedures, using adequate distribution functions, and obtain a set of coupled differential equations to describe the evolution of the perturbations of every known component of the Universe in Fourier space[23]:

$$\dot{\Theta} + ik\mu\Theta = -\dot{\Phi} - ik\mu\Psi - \dot{\tau} \left[ \Theta_0 - \Theta + \mu v_b - \frac{1}{2} \mathcal{P}_2(\mu)(\Theta_2 + \Theta_{P2} + \Theta_{P0}) \right], \quad (2.34)$$

$$\dot{\Theta}_P + ik\mu\Theta_P = -\dot{\tau} \left\{ -\Theta_P + \frac{1}{2} [1 - \mathcal{P}_2(\mu)] (\Theta_2 + \Theta_{P2} + \Theta_{P0}) \right\}, \quad (2.35)$$

$$\dot{\delta} + ikv = -3\dot{\Phi}, \quad (2.36)$$

$$\dot{v} + Hv = -ik\Psi, \quad (2.37)$$

$$\dot{v}_b + Hv_b = -ik\Psi + \frac{4\rho_\gamma}{3\rho_b} \dot{\tau} (v_b + 3i\Theta_1), \quad (2.38)$$

$$\dot{\mathcal{N}} + ik\mu\mathcal{N} = -\dot{\Phi} - ik\mu\Psi. \quad (2.39)$$

Here,  $\Theta_P$  is the polarization field and  $\Theta_{P\ell}$  is defined by equation (2.25) using  $F = \Theta_P$ ;  $\delta$  and  $\delta_b$  are the perturbations to the matter and baryon distributions respectively;  $v$  and  $v_b$  are the bulk velocities of the fluids in the stress-energy tensor for matter and baryons respectively;  $\rho_b$  and  $\rho_\gamma$  are the baryon and radiation energy densities respectively; and  $\tau$  is the optical depth of the Universe.

These equations, alongside equations (2.23) and (2.24), are very complex to be solved analytically, so it is common to evolve them numerically, and there are a few codes developed specifically to do this. One of them, that has been used in this project, is CAMB.[31, 32]

## 2.3 The Cosmic Microwave Background

With the dynamical equations for photons determined, one can obtain the temperature perturbation distribution  $\Theta$  if initial conditions are provided. With the equations already presented, it is possible to show that the initial conditions at the early Universe depend only on the initial conditions for  $\Phi$ [23], which are then evolved through the inflationary period assuming the existence of a quantum scalar field  $\phi(\mathbf{x}, t)$  that drives a brief accelerated expansion.

With time, particles start to form: Quarks, hadrons, neutrinos, amongst others, at different times. Eventually baryons have formed and the photons are tightly coupled to them, forming a baryon-photon fluid that can be described by the differential equations (2.34) - (2.39), the initial distribution obtained for this fluid can then be used as the initial condition and evolved to predict the present anisotropies to the uniform background. To analyze these predictions, the statistical behavior of each component of the Universe is taken: either their spectral densities (see Appendix A) – usually called their power spectra – or what is known as their correlation spectra.

### 2.3.1 CMB Autocorrelation Spectrum

Equation (2.30) explicitly assumes a dependence in position, time, and momentum direction. Measurements made on Earth can assume a fixed value for time since our measurements related to the Universe are taken in a period negligible in the cosmological scale, but the translational movement of the Earth is relevant to our measurements: measurements of the CMB made by a non-comoving observer, when projected onto a sphere, lead to a strong dipole pattern in the temperature distribution, modulation of the multipole terms and a small relativistic aberration at angular scales of a few arcminutes[33–35].

Despite these effects, measurements of the temperature distribution can be projected onto a sphere independent of the Earth's position, making it possible to describe this distribution simply using two angle variables, or simply  $\hat{\mathbf{p}}$ .

We can expand the temperature perturbations in spherical harmonics:

$$\Theta(\mathbf{x}, \hat{\mathbf{p}}, t) = \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m}(\mathbf{x}, t) Y_{\ell m}(\hat{\mathbf{p}}). \quad (2.40)$$

All the information about temperature maps of the CMB measured on Earth are contained in the set of  $a_{\ell m}$  values. Given the limited number of pixels  $N_{\text{pix}}$  of any of these maps, there is a maximum value of  $\ell$  ( $\ell_{\text{max}}$ ) up to which statistically meaningful information can be retrieved. For each value of  $\ell$ , there are  $2\ell + 1$  values of  $m$  – and  $a_{\ell m}$  – in the series, a hypothetical observed map with precision  $p_r$  in squared radians would have  $4\pi/p_r$  pixels, setting this number of pixels equal to the number of  $a_{\ell m}$  used to describe this map gives a reasonable estimation of  $\ell_{\text{max}}$  [23], so

$$\sum_{\ell=0}^{\ell_{\text{max}}} (2\ell + 1) = (\ell_{\text{max}} + 1)^2 = N_{\text{pix}} = \frac{4\pi}{p_r}. \quad (2.41)$$

Using the orthogonality property of the spherical harmonics, one can obtain a direct expression for  $a_{\ell m}$  in terms of the angular distribution of the perturbations:

$$a_{\ell m}(\mathbf{x}, t) = \int \frac{d^3 k}{(2\pi)^3} e^{i\mathbf{k} \cdot \mathbf{x}} \int d\Omega Y_{\ell m}^*(\hat{\mathbf{p}}) \Theta(\mathbf{k}, \hat{\mathbf{p}}, t). \quad (2.42)$$

Again, there are available codes that efficiently and precisely calculate  $a_{\ell m}$  from a pixelized map and vice versa, the one used in this work was HEALPix [36]. Under the hypothesis that the spherical harmonics coefficients  $a_{\ell m}$  are independent Gaussian random variables with zero mean, all its statistical properties are contained in the variance  $C_\ell$ , also known as the power spectrum:

$$\langle a_{\ell m} a_{\ell' m'}^* \rangle = \delta_{\ell \ell'} \delta_{m m'} C(\ell), \quad (2.43)$$

where  $\delta_{mm'}$  is the Kronecker delta taking  $m$  and  $m'$  as arguments.

The fact that for lower values of  $\ell$  there are fewer coefficients  $a_{\ell m}$  naturally leads to lower multipoles  $\ell$  having higher uncertainty, this property is usually called cosmic variance and can be expressed using [23]

$$\left( \frac{\Delta C(\ell)}{C(\ell)} \right)_{\text{cosmic variance}} = \sqrt{\frac{2}{2\ell + 1}}. \quad (2.44)$$

The temperature autocorrelation defined in equation (2.43) is usually expressed as  $C_\ell^{TT}$ , but sometimes a modified spectrum  $D_\ell^{TT} = \frac{\ell(\ell+1)}{2\pi} C_\ell^{TT}$  is used for visualization purposes. Figure 2.2 shows Planck’s CMB power spectrum data and its corresponding best-fit spectrum assuming a  $\Lambda$ CDM cosmology.




Figure 2.2: Planck 2018 CMB temperature power spectrum: the blue line corresponds to a  $\Lambda$ CDM cosmology best-fit to the data points in orange. For  $\ell < 30$  a logarithmic scale was used and the data points correspond to the  $D_\ell^{TT}$  for each value of  $\ell$ , while for  $\ell \geq 30$  a linear scale was used and the data points correspond to binned values for the data for  $D_\ell^{TT}$ . The error bars show  $\pm 1\sigma$  uncertainties, including cosmic variance. Extracted from [6].

### 2.3.2 Analytical Approximation

Although solving equations such as (2.34) numerically gives accurate results, obtaining an analytical approximation to these equations helps us understand the underlying physical effects that most influence the evolution of the perturbations. Working on the tightly coupled limit and assuming recombination to be an instantaneous process happening at conformal time  $\eta = \eta_*$ , one can obtain the following equation[23]:

$$\begin{aligned}
\Theta_\ell(k, \eta_0) \approx & [\Theta_0(k, \eta_*) + \Psi(k, \eta_*)] j_\ell[k(\eta_0 - \eta_*)] \\
& + iv_b(k, \eta_*) \left\{ j_\ell[k(\eta_0 - \eta_*)] - (l+1) \frac{j_\ell[k(\eta_0 - \eta_*)]}{k(\eta_0 - \eta_*)} \right\} \\
& + \int_0^{\eta_0} d\eta e^{-\tau} [\Psi'(k, \eta) - \Phi'(k, \eta)] j_\ell[k(\eta_0 - \eta)],
\end{aligned} \tag{2.45}$$

where  $j_\ell$  are spherical Bessel functions, the conformal time  $\eta$  is defined as

$$\eta = \int_0^t \frac{dt}{a(t)}, \tag{2.46}$$

and  $\eta_*$  is the value of  $\eta$  that maximizes the visibility function  $g(\eta)$ :

$$g(\eta) = -\tau' e^{-\tau}. \tag{2.47}$$

With these conventions, the derivative with respect to  $\eta$  is being represented by the apostrophe notation, while the dot notation used in equation (2.34) will still indicate derivatives with respect to time.

There are three main contributors to the solution (2.45):

- The  $(\Theta_0 + \Psi)$  term is usually called the Sachs-Wolfe term, which is associated with Baryon Acoustic Oscillations (BAO), which are fluctuations in the density of the visible baryonic matter of the Universe, caused by acoustic density waves in the primordial plasma of the early Universe[23];
- The  $iv_b$  term shows a contribution to the spectrum related to a Doppler shift of the BAO, due to its dependence on a non-zero bulk velocity of the baryons during recombination;
- The last term, an integral involving the time variations of the gravitational potentials, represents the Integrated Sachs-Wolfe (ISW) effect. Its dependence on the variation of the gravitational potential makes it so that it only has a non-negligible effect on the CMB at cosmological periods with high variation of these potentials: At around  $a \approx 10^{-4}$  the Universe transitioned from a radiation-dominated to a matter-dominated Universe (early ISW), and recently – around  $a \approx 0.61$  or  $z \approx 0.63$ [37] according to the  $\Lambda$ CDM model – the Universe transitioned from a matter-dominated Universe to the current dark energy dominated Universe (late ISW).

## 2.4 The Integrated Sachs Wolfe Effect

It was proposed that the cross-correlation spectrum between CMB maps and matter contrast could be used to detect the ISW effect[38], and since then many works have reported the detection of this signal[39–42]. In this section, a framework for theoretical estimations of the ISW contribution to this cross-correlation will be discussed.

### 2.4.1 The Matter Power Spectrum

While the CMB is usually described as a two dimensional map measured around the Earth, the matter distribution in the Universe is first analyzed taking the distance from the Earth into consideration, so three coordinates are necessary, and then one can take a 2D projection for a certain redshift  $z$ . Again using Fourier space, we first define the 3D matter power spectrum for the distribution of matter perturbations in the Universe  $P(k)$  as

$$\langle \delta(\mathbf{k}, z) \delta^*(\mathbf{k}', z) \rangle = (2\pi)^3 P(k, z) \delta_D(\mathbf{k} - \mathbf{k}'), \quad (2.48)$$

where  $\delta_D$  is the Dirac delta,  $\delta$  is the previously shown matter perturbation distribution in the Universe and the  $*$  superscript indicates the complex conjugate of a quantity. Again, using appropriate initial conditions, one can determine  $\delta(\mathbf{k})$  and thus  $P(k, z)$ , and again these results can be obtained numerically using CAMB, Figure 2.3 shows the results for this calculation using the HALOFIT model[43], a semi-analytical method to determine a non-linear solution for  $P(k, z)$ .

### 2.4.2 The Galaxy-CMB Correlation Spectra

To obtain the autocorrelation for a spherical projection of a matter density distribution on a spherical shell on a certain redshift  $z$ , one has to choose a way to trace the matter density with an observable. One way of doing this is by tracing the mass distribution with galaxy density on a certain redshift. It is common to assume that the galaxy density perturbations, usually called galaxy contrast maps, take the form  $\delta_g = b_g \delta$  where  $b_g$  is a constant value (for each redshift  $z$ ) called the bias factor. From this point, the same process needed to calculate (2.43) can be applied:




Figure 2.3: Matter power spectrum calculated as a function of  $k/h$ , where  $h = H_0/(100 \text{ km/s/Mpc})$ , using the HALOFIT model with cosmological parameters according to Planck's best fit results[6].

Expand the distribution in spherical harmonics with factors  $a_{\ell m}^g$  and calculate the variance of these variables to obtain an autocorrelation spectrum  $C^{gg}(\ell)$ .

Similarly, instead of calculating an autocorrelation spectrum, one can calculate a cross-correlation between two perturbation distributions, which is analogous to definition (2.43):

$$\langle a_{\ell m}^t a_{\ell' m'}^g \rangle = C^{tg}(\ell) \delta_{\ell \ell'} \delta_{m m'}. \quad (2.49)$$

One can determine a general and direct equation to calculate these spectra. In this work we have used the form shown in [16], which is specific to the ISW contribution to these correlation spectra:

$$C_\ell^{xy} = \frac{2}{\pi} \int dk k^2 W_\ell^x(k) W_\ell^y(k) P(k), \quad (2.50)$$

where  $x$  and  $y$  can represent  $g$  or  $t$ , and  $W_\ell$  are called the kernel functions, which can be calculated using

$$W_\ell^t = -3\Omega_m \left( \frac{H_0}{k} \right)^2 \int dz \frac{d[(1+z)D(z)]}{dz} j_\ell[k\chi(z)], \quad (2.51)$$



$$W_\ell^g = \int dz b_g(z) \frac{dN}{dz} D(z) j_\ell[k\chi(z)], \quad (2.52)$$

where  $D(z)$  is the linear growth function (normalized to 1 at  $z = 0$ );  $\chi(z)$  is the comoving distance; and  $\frac{dN}{dz}$  is the normalized galaxy redshift distribution, also called the selection function[39].

With equation (2.50), we can determine the theoretical spectra for CMB and galaxy autocorrelations and CMB-galaxy cross-correlation directly. Figure 2.4 shows the total CMB autocorrelation and the ISW contribution, both at  $z = 0$ .




Figure 2.4: CMB autocorrelation comparison for the late-ISW term and the full spectrum. Calculations were made using CAMB (for the full spectrum) and codes developed during the project. The parameters used are those corresponding to the Planck best fit parameters[6].

Figure 2.5 shows the galaxy autocorrelation spectrum and the galaxy-CMB cross-correlation spectrum. All of these spectra have been integrated numerically.

These plots, show very clearly that the ISW effect has a lot more relevance in the lower multipoles, where cosmic variance has a significant influence. The selection function is an important aspect of this signal, in chapter 3 the matter of how to optimize the signal with the use of an idealized selection function will be discussed.




Figure 2.5: Galaxy autocorrelation spectrum (left) and late-ISW contribution to the galaxy-CMB cross-correlation (right). The selection function used to calculate these spectra was parametrized to be compatible with 2MASS' galaxy redshift distribution[16, 44].

### 2.4.3 Dependence on Dark Energy Models

Many approaches to study dark energy have been proposed, each of them with different properties, ranging from parameterization of the equation of state[45, 46] to alternative fluid models for dark energy[47] and modifications to general relativity[48].

The simplest parametrization proposed for dark energy's equation of state is called the Chevallier-Polarski-Linder (CPL) parametrization[26, 45]:

$$w(z) = w_0 + \left(1 - \frac{1}{1+z}\right) w_a. \quad (2.53)$$

So far, this parametrization has not solved some of the current limitations being faced by the  $\Lambda$ CDM model, such as the Hubble tension [49] and the  $\sigma_8$  tension [6], it has not improved over the  $\Lambda$ CDM predictions, but it shows good compatibility with CMB, type IA supernovae and baryon acoustic oscillations (BAO) data[6]. Planck results using this parametrization are shown in Table 2.2.

Another important parameter related to dark energy is its speed of sound, defined as[53]:

$$c_s^2 = \frac{\delta P_{DE}}{\delta \rho_{DE}} \quad (2.54)$$

where  $\rho_{DE}$  and  $P_{DE}$  are the dark energy's density and pressure respectively, and  $\delta$  indicates the perturbation of these quantities.

Parameter	Marginalized Value
$w_0$	$-0.957 \pm 0.080$
$w_a$	$-0.29^{+0.32}_{-0.26}$
$H_0[\text{km s}^{-1}\text{Mpc}^{-1}]$	$68.31 \pm 0.82$
$\sigma_8$	$0.820 \pm 0.011$
$\Delta\chi^2$	-1.4

Table 2.2: Marginalized values and 68% confidence limits for cosmological parameters, assuming parametrization (2.53), obtained by combining Planck temperature and polarization autocorrelation and cross-correlation spectra[50], type IA supernovae data from the Joint Light Curve Analysis (JLA)[51] and BAO data from the Sloan Digital Sky Survey (SDSS)[52]. The  $\Delta\chi^2$  value was computed with respect to the  $\Lambda$ CDM best fits computed from the same data set combination.

The value of this parameter influences, for instance, the evolution of the gravitational potentials[54], and many other factors that relate to the evolution of the universe, thus having an impact on observables.

Figure 2.6 shows the influence of different parameter values, using the CPL parametrization, on the CMB temperature autocorrelation spectrum, and compares the expected theoretical spectra with Planck's data.



Figure 2.6: The solid lines in the plots correspond to theoretical predictions for the CMB temperature autocorrelation spectra corresponding to different parameter values of the dark energy equation of state, the black data points correspond to Planck's data[6]. Figures 2.6a, 2.6b and 2.6c show how the spectrum changes for a constant value of  $w_a$  shown on the top left of the axis and varying speeds of sound, while the other three figures show how the curve behaves with varying values of  $w_a$  and a constant speed of sound. All theoretical calculations were done using CAMB.

### 3 | Optimizing the Selection Function

Equation (2.52) introduced the selection function  $\frac{dN}{dz}$ , which simply corresponds to the redshift distribution of galaxy number  $N$  and can vary depending on the galaxy catalog used, thus having an influence on the expected signal that can be extracted from that catalog. As shown in Figure 2.5, the ISW signal is mostly present in low multipoles, where cosmic variance has a strong influence, this motivated us to study the possibility of optimizing the cross-correlation signal by determining a selection function that could theoretically maximize it, this study will be presented in this chapter.

#### 3.1 Parametrization of the Selection Function

Following [39], we are assuming the selection function can be parametrized as

$$\frac{dN}{dz}(z|\lambda, \beta, z_0) dz = \frac{\beta}{\Gamma(\lambda)} \left(\frac{z}{z_0}\right)^{\beta\gamma-1} \exp\left[-\left(\frac{z}{z_0}\right)^\beta\right] d\left(\frac{z}{z_0}\right) \quad (3.1)$$

where the parameters  $\beta$ ,  $\lambda$  and  $z_0$  are all positive. The values used to calculate the spectra in 2.5 are those corresponding to the so called band 1 of the 2MASS catalog. Table 3.1 shows the parameter values for each band of this catalog, while Figure 3.1 shows the plots for their corresponding selection functions. The details on what each of these bands represent will be presented in chapter 4.

Band	$z_0$	$\beta$	$\lambda$
1	0.043	1.825	1.524
2	0.054	1.800	1.600
3	0.067	1.765	1.636
4	0.084	1.723	1.684

Table 3.1: Parameter values for the 4 bands of the 2MASS catalog. Values extracted from [39].

To start the search for an ideal selection, a good first guess would be to find functions that favor galaxies at redshifts close to the period of transition to the current dark energy dominated universe, which has been stated before in this text to be  $z^* \approx 0.63$ , a value that I calculated inputting the  $\Lambda$ CDM values for cosmological parameter in equation[37]



Figure 3.1: Selection function calculated for the 4 bands of the 2MASS catalog according to equation (3.1).

$$z^* = \left[ -(1 + 3w_{DE}) \frac{\Omega_{DE}}{\Omega_m} \right]^{-\frac{1}{3w_{DE}}} - 1, \quad (3.2)$$

which assumes a constant equation of state parameter  $w_{DE}$  for dark energy.

To determine the region in the parameter space that favors this redshift, a study of the influence of each parameter in the selection function was necessary. Figure 3.2 shows how the selection function varies with changes in each of its parameters.




Figure 3.2: Selection function calculated for various parameter values according to the parametrization (3.1). Each plot is made with constant values for two parameters, corresponding to the band 2 of the 2MASS catalog (see Table 3.1), and the parameter on the right is varied.

Figure 3.2 shows that increasing the value of  $\lambda$  seems to shift the peak of the selection function towards higher redshifts without significantly changing the width of the curve, while increasing  $\beta$  seems to narrow the curve without significantly changing the position of the peak, and increasing  $z_0$  seems to increase both the position of the maximum and the width of the curve.

These observations make it easier to change the parameters and see their influence on the cross-correlation signal and helps choose the starting region of the parameter space to explore, but to do so, the matter of how to quantify the quality of the signal has to be discussed.

## 3.2 Maximizing the Cross-Correlation Signal

To maximize the signal we are in search of, we compare it with the null hypothesis that there is no cross-correlation induced by the ISW effect, i.e  $C^{tg}(\ell) = 0$  for every multipole. To do this is, we need to estimate the probability distribution of  $C^{tg}$  for each multipole  $\ell$  extracted from an uncorrelated map.

In this part of the project, the HEALPix software was used extensively, and to validate the procedure for the search of a probability distribution for the null hypothesis, we first tested how the software would be able to retrieve the signal of a non-null cross-correlation spectrum. To do this, the full autocorrelation spectrum in 2.4 calculated using CAMB was used as well as the cross-correlations 2.5 as inputs for a HEALPix function that can sample values of  $a_{\ell m}$  for the temperature and galaxy contrast maps that follow both the autocorrelation functions given, as well as the cross-correlation function, and a map representation can be calculated with this using another HEALPix functionality.

Figure ?? illustrates one galaxy contrast map and one CMB temperature map calculated using this procedure.

### FIGURA DO MAPA HEALPIX

It is necessary that multiple maps are synthesized this way in order to use these maps to actually retrieve the correlation spectra properly. During our tests, the autocorrelations quickly converged to their corresponding theoretical spectra, but not the cross-correlation. Figure ?? shows a comparison between the theoretical cross-correlation spectrum and the one estimated from ten thousand synthesized maps, which was the value we considered a reasonable tradeoff between quality and time efficiency for this project.

### FIGURA DO CTG SINTÉTICO+TEÓRICO

The same process can be done using  $C^{tg} = 0$  for every multipole, giving us an estimation of the distribution of  $C^{tg}(\ell)$  that can be used as a probability distribution for the null hypothesis. Figure ?? shows two histograms representing this distribution.

### FIGURA DOS HISTOGRAMAS

Upon thorough inspection of the histograms, we decided to fit each histogram using the gaussian distribution function, and using that as the distribution function of  $C^{tg}$  for that multipole, each of them will be indicated by  $f_\ell[C^{tg}(\ell)]$ .

With that, we can finally define the probability  $P_{\text{null}}$  that a correlation spectrum  $C^{tg}$  follows the null hypothesis:

$$P_{\text{null}} = \prod_{l=2}^{\ell_{\text{max}}} f_l [C^{tg}(\ell)] \quad (3.3)$$

With this, we can then explore the parameter space  $(z_0, \beta, \lambda)$  and estimate the theoretical cross-correlation spectrum for each point in it, and determine  $P_{\text{null}}$  for each of them. If we are able to find the point in this parameter space that minimizes  $P_{\text{null}}$ , that should contribute to the improvement of the cross-correlation signal and reduce the dependence on cosmic variance.

## 3.3 Results

SHOW THE HEAT PLOTS, THE MINIMIZER FINAL RESULT AND DISCUSS HOW IT DOESN'T SEEM TO HAVE A RELEVANT EFFECT ON THE FINAL RESULTS



## 4 | Extracting Correlations from Data

CMB experiments do not directly measure correlation spectra, instead the data consists of time ordered data streams, each stream with information of the direction of observation and the sky temperature measured by that detector, which is contained within an array of detectors.

In this chapter, the process of converting this signal, taking into account the effects of detector noise and environmental variables such as the galactic background, will be discussed, not only for CMB but also for galaxy maps. A Monte Carlo Markov Chain (MCMC) algorithm capable of efficiently extracting these correlations, including the galaxy-CMB cross-correlation, will be presented.

### 4.1 Two Point Functions

To estimate any correlation spectrum, we first need to estimate the values of the multipole terms  $a_{\ell m}$ , but applying equation (Cl definition) using these does not yield accurate results, one has to introduce factors such as noise and map masks first.

In general, it is common to introduce a weight function  $W(\hat{\mathbf{n}})$  to account for various possible effects of the map measured, such as galactic contamination, edge effects, and down-weighting more noisy pixels (CITAR HIVON 2002). If we ignore other effects for now (such as noise), the multipole terms  $\tilde{a}_{\ell m}$  as defined in (ref. alm definition), assuming a high number of small pixels to approximate a sum with an integral, becomes

$$\tilde{a}_{\ell m} = \int d\Omega \Theta(\hat{\mathbf{n}}) W(\hat{\mathbf{n}}) Y_{\ell m}^*(\hat{\mathbf{n}}). \quad (4.1)$$

With this, one can define the so-called pseudo-power spectrum  $\tilde{C}_\ell$ :

$$\tilde{C}_\ell = \frac{1}{2\ell + 1} \sum_{m=-\ell}^{\ell} |\tilde{a}_{\ell m}|^2. \quad (4.2)$$

(para ctg, posso trocar  $|alm|^2$  por  $(alm^t)(alm^{g*})$  por exemplo?)

Still ignoring noise and other factors, one can relate the ensemble average of the pseudo-power spectrum with the true power spectrum with a simple system of equations

$$\langle \tilde{C}_\ell \rangle = \sum_{\ell'} M_{\ell\ell'} \langle C_{\ell'} \rangle, \quad (4.3)$$

where  $M_{\ell\ell'}$  is a matrix that describes the coupling between modes, called the multipole mixing matrix. For a map masking the CMB map (for instance, the galactic contamination map) with multipole functions  $\mathcal{W}_\ell$ , the mixing matrix corresponds to

$$M_{\ell\ell'} = \frac{2\ell' + 1}{4\pi} \sum_{\ell''} (2\ell'' + 1) \mathcal{W}_\ell'' \begin{pmatrix} \ell & \ell' & \ell'' \\ 0 & 0 & 0 \end{pmatrix}^2. \quad (4.4)$$

(como pode elevar essa matriz ao quadrado see o numero de linhas é diferente do número de colunas?)

To determine the full form of the system of equations used to determine the real power spectrum, one has to take noise into consideration and the so-called beam pattern.

(EXPLICAR O QUE É UM "BEAM" E EXPLICAR O BEAM PATTERN COM IMAGEM DE EXEMPLO)

If  $B_\ell$  is the multipole function of the  $\ell$  multipole, and given a noise map  $N$  that can also be expanded in multipoles  $N_\ell$ , the correlation functions can be extracted by solving the following system of equations:

$$\langle \tilde{C}_\ell \rangle = \sum_{\ell'} M_{\ell\ell'} B_{\ell'}^2 \langle C_{\ell'} \rangle + \langle N_\ell \rangle. \quad (4.5)$$

## 4.2 Gibbs Sampling

## 4.3 Data Sets and Analysis

2MASS CATALOG, INCLUDE EXPLANATION OF 2MASS BANDS, PLANCK DATA, ETC

## 5 | Results

## 6 | Conclusions

Artigo do Edivaldo: [16]

ISW artigo original: [15]

The GNU parallel software was used: [55]

Cobaya: [56, 57]

Gibbs Sampling: [58]

# Appendices

# A | Fourier Space

Falar sobre espaço de fourier

Falar do conceito de spectral density/power spectrum

## B | Monte Carlo Markov Chains

# Bibliography

- [1] A. A. Penzias and R. W. Wilson. “A Measurement of Excess Antenna Temperature at 4080 Mc/s.” In: *Astrophysical Journal* 142 (July 1965), pp. 419–421. DOI: 10.1086/148307.
- [2] Ivan Agullo, Dimitrios Kranas, and V Sreenath. “Large scale anomalies in the CMB and non-Gaussianity in bouncing cosmologies”. In: *Classical and Quantum Gravity* 38.6 (Feb. 2021), p. 065010. DOI: 10.1088/1361-6382/abc521. URL: <https://dx.doi.org/10.1088/1361-6382/abc521>.
- [3] Xingang Chen. “Primordial Non-Gaussianities from Inflation Models”. In: *Adv. Astron.* 2010 (2010), p. 638979. DOI: 10.1155/2010/638979. arXiv: 1002.1416 [astro-ph.CO].
- [4] Robert G. Crittenden and Neil Turok. “Looking for a Cosmological Constant with the Rees-Sciama Effect”. In: *Phys. Rev. Lett.* 76 (4 Jan. 1996), pp. 575–578. DOI: 10.1103/PhysRevLett.76.575. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.76.575>.
- [5] G. Hinshaw et al. “FIVE-YEAR WILKINSON MICROWAVE ANISOTROPY PROBE\* OBSERVATIONS: DATA PROCESSING, SKY MAPS, AND BASIC RESULTS”. In: *The Astrophysical Journal Supplement Series* 180.2 (Feb. 2009), p. 225. DOI: 10.1088/0067-0049/180/2/225. URL: <https://dx.doi.org/10.1088/0067-0049/180/2/225>.
- [6] Planck Collaboration et al. “Planck 2018 results - VI. Cosmological parameters (Corrigendum)”. In: *Astronomy & Astrophysics* 652 (2021), p. C4. DOI: 10.1051/0004-6361/201833910e. URL: <https://doi.org/10.1051/0004-6361/201833910e>.
- [7] P. J. E. Peebles and J. T. Yu. “Primeval adiabatic perturbation in an expanding universe”. In: *Astrophys. J.* 162 (1970), pp. 815–836. DOI: 10.1086/150713.
- [8] D. J. Fixsen. “THE TEMPERATURE OF THE COSMIC MICROWAVE BACKGROUND”. In: *The Astrophysical Journal* 707.2 (Nov. 2009), p. 916. DOI: 10.1088/0004-637X/707/2/916. URL: <https://dx.doi.org/10.1088/0004-637X/707/2/916>.
- [9] Wayne Hu and Naoshi Sugiyama. “Anisotropies in the Cosmic Microwave Background: an Analytic Approach”. In: *The Astrophysical Journal* 444 (May 1995), p. 489. DOI: 10.1086/175624. arXiv: astro-ph/9407093 [astro-ph].
- [10] D. J. Fixsen et al. “Calibration of the COBE FIRAS Instrument”. In: *The Astrophysical Journal* 420 (Jan. 1994), p. 457. DOI: 10.1086/173577.
- [11] Marc Kamionkowski, Arthur Kosowsky, and Albert Stebbins. “A Probe of Primordial Gravity Waves and Vorticity”. In: *Phys. Rev. Lett.* 78 (11 Mar. 1997), pp. 2058–2061. DOI: 10.1103/PhysRevLett.78.2058. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.78.2058>.



- [12] Silvia Galli et al. “CMB polarization can constrain cosmology better than CMB temperature”. In: *Phys. Rev. D* 90 (6 Sept. 2014), p. 063504. DOI: 10.1103/PhysRevD.90.063504. URL: <https://link.aps.org/doi/10.1103/PhysRevD.90.063504>.
- [13] P. A. R. Ade et al. “Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season”. In: *Phys. Rev. Lett.* 127 (15 Oct. 2021), p. 151301. DOI: 10.1103/PhysRevLett.127.151301. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.127.151301>.
- [14] The Planck Collaboration. *The Scientific Programme of Planck*. 2006. arXiv: astro-ph/0604069 [astro-ph].
- [15] R. K. Sachs and A. M. Wolfe. “Perturbations of a cosmological model and angular variations of the microwave background”. In: *Astrophys. J.* 147 (1967), pp. 73–90. DOI: 10.1007/s10714-007-0448-9.
- [16] E. Moura-Santos et al. “A Bayesian Estimate of the CMB–large-scale Structure Cross-correlation”. In: *The Astrophysical Journal* 826.2 (July 2016), p. 121. DOI: 10.3847/0004-637X/826/2/121. URL: <https://dx.doi.org/10.3847/0004-637X/826/2/121>.
- [17] Albert Einstein. “Cosmological Considerations in the General Theory of Relativity”. In: *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys. )* 1917 (1917), pp. 142–152.
- [18] Y. Choquet-Bruhat. *General Relativity and the Einstein Equations*. Oxford Mathematical Monographs. OUP Oxford, 2008. ISBN: 9780191578854. URL: <https://books.google.com.br/books?id=UjHbm5rfpi8C>.
- [19] Suman Sarkar, Biswajit Pandey, and Rishi Khatri. “Testing isotropy in the Universe using photometric and spectroscopic data from the SDSS”. In: *Monthly Notices of the Royal Astronomical Society* 483.2 (Nov. 2018), pp. 2453–2464. ISSN: 0035-8711. DOI: 10.1093/mnras/sty3272. eprint: <https://academic.oup.com/mnras/article-pdf/483/2/2453/27184728/sty3272.pdf>. URL: <https://doi.org/10.1093/mnras/sty3272>.
- [20] Biswajit Pandey and Suman Sarkar. “Testing homogeneity of the galaxy distribution in the SDSS using Renyi entropy”. In: *Journal of Cosmology and Astroparticle Physics* 2021.07 (July 2021), p. 019. DOI: 10.1088/1475-7516/2021/07/019. URL: <https://dx.doi.org/10.1088/1475-7516/2021/07/019>.
- [21] Edwin Hubble. “A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae”. In: *Proceedings of the National Academy of Science* 15.3 (Mar. 1929), pp. 168–173. DOI: 10.1073/pnas.15.3.168.
- [22] Georges Lemaitre. “A Homogeneous Universe of Constant Mass and Growing Radius Accounting for the Radial Velocity of Extragalactic Nebulae”. In: *Annales Soc. Sci. Bruxelles A* 47 (1927), pp. 49–59. DOI: 10.1007/s10714-013-1548-3.
- [23] S. Dodelson and F. Schmidt. *Modern Cosmology*. Elsevier Science, 2020. ISBN: 9780128159484. URL: <https://books.google.com.br/books?id=GGjfywEACAAJ>.

- [24] Fulvio Melia. “The Friedmann–Lemaître–Robertson–Walker metric”. In: *Modern Physics Letters A* 37.03 (2022), p. 2250016. DOI: 10.1142/S021773232250016X. eprint: <https://doi.org/10.1142/S021773232250016X>. URL: <https://doi.org/10.1142/S021773232250016X>.
- [25] Steven Weinberg. “The cosmological constant problem”. In: *Rev. Mod. Phys.* 61 (1 Jan. 1989), pp. 1–23. DOI: 10.1103/RevModPhys.61.1. URL: <https://link.aps.org/doi/10.1103/RevModPhys.61.1>.
- [26] Verónica Motta et al. “Taxonomy of Dark Energy Models”. In: *Universe* 7.6 (2021), p. 163. DOI: 10.3390/universe7060163. arXiv: 2104.04642 [astro-ph.CO].
- [27] B. Schutz. *A First Course in General Relativity*. Cambridge University Press, 2009. ISBN: 9780521887052.
- [28] Ofer Lahav and Andrew R Liddle. *The Cosmological Parameters 2014*. 2014. arXiv: 1401.1389 [astro-ph.CO].
- [29] Alan H. Guth. “Inflationary universe: A possible solution to the horizon and flatness problems”. In: *Phys. Rev. D* 23 (2 Jan. 1981), pp. 347–356. DOI: 10.1103/PhysRevD.23.347. URL: <https://link.aps.org/doi/10.1103/PhysRevD.23.347>.
- [30] A.D. Linde. “A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems”. In: *Physics Letters B* 108.6 (1982), pp. 389–393. ISSN: 0370-2693. DOI: [https://doi.org/10.1016/0370-2693\(82\)91219-9](https://doi.org/10.1016/0370-2693(82)91219-9). URL: <https://www.sciencedirect.com/science/article/pii/0370269382912199>.
- [31] Antony Lewis, Anthony Challinor, and Anthony Lasenby. “Efficient computation of CMB anisotropies in closed FRW models”. In: *ApJ* 538 (2000), pp. 473–476. DOI: 10.1086/309179. arXiv: astro-ph/9911177 [astro-ph].
- [32] Antony Lewis and Sarah Bridle. “Cosmological parameters from CMB and other data: A Monte Carlo approach”. In: *PRD* 66 (2002), p. 103511. DOI: 10.1103/PhysRevD.66.103511. arXiv: astro-ph/0205436 [astro-ph].
- [33] D. J. Fixsen et al. “Calibration of the COBE FIRAS Instrument”. In: *The Astrophysical Journal* 420 (Jan. 1994), p. 457. DOI: 10.1086/173577.
- [34] Planck Collaboration et al. “Planck 2013 results. XXVII. Doppler boosting of the CMB: Eppur si muove”. In: *Astronomy & Astrophysics* 571 (2014), A27. DOI: 10.1051/0004-6361/201321556. URL: <https://doi.org/10.1051/0004-6361/201321556>.
- [35] Esteban Ricardo Chalbaud Mogollón. *Kinematic effects and the break of statistical isotropy of the cosmic microwave background fluctuations*. 2021. DOI: <https://doi.org/10.11606/D.43.2021.tde-31052021-162009>.
- [36] K. M. Górski et al. “HEALPix: A Framework for High-Resolution Discretization and Fast Analysis of Data Distributed on the Sphere”. In: *ApJ* 622 (Apr. 2005), pp. 759–771. DOI: 10.1086/427976. eprint: astro-ph/0409513.
- [37] José Carlos N. de Araujo. “The dark energy–dominated Universe”. In: *Astroparticle Physics* 23.2 (2005), pp. 279–286. ISSN: 0927-6505. DOI: <https://doi.org/10.1016/j.astropartphys.2004.12.004>. URL: <https://www.sciencedirect.com/science/article/pii/S0927650504002087>.

- [38] Robert G. Crittenden and Neil Turok. “Looking for a Cosmological Constant with the Rees-Sciama Effect”. In: *Phys. Rev. Lett.* 76 (4 Jan. 1996), pp. 575–578. DOI: 10.1103/PhysRevLett.76.575. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.76.575>.
- [39] Niayesh Afshordi, Yeong-Shang Loh, and Michael A. Strauss. “Cross-correlation of the cosmic microwave background with the 2MASS galaxy survey: Signatures of dark energy, hot gas, and point sources”. In: *Phys. Rev. D* 69 (8 Apr. 2004), p. 083524. DOI: 10.1103/PhysRevD.69.083524. URL: <https://link.aps.org/doi/10.1103/PhysRevD.69.083524>.
- [40] S. P. Boughn and R. G. Crittenden. “The cross-correlation between the microwave and X-ray backgrounds: foregrounds and systematics”. In: *Monthly Notices of the Royal Astronomical Society* 360.3 (July 2005), pp. 1013–1021. ISSN: 0035-8711. DOI: 10.1111/j.1365-2966.2005.09090.x. eprint: <https://academic.oup.com/mnras/article-pdf/360/3/1013/3178164/360-3-1013.pdf>. URL: <https://doi.org/10.1111/j.1365-2966.2005.09090.x>.
- [41] S.P. Boughn and R.G. Crittenden. “A detection of the integrated Sachs–Wolfe effect”. In: *New Astronomy Reviews* 49.2 (2005). Sources and Detection of Dark Matter and Dark Energy in the Universe, pp. 75–78. ISSN: 1387-6473. DOI: <https://doi.org/10.1016/j.newar.2005.01.005>. URL: <https://www.sciencedirect.com/science/article/pii/S1387647305000096>.
- [42] Planck Collaboration et al. “Planck 2015 results - XXI. The integrated Sachs-Wolfe effect”. In: *A&A* 594 (2016), A21. DOI: 10.1051/0004-6361/201525831. URL: <https://doi.org/10.1051/0004-6361/201525831>.
- [43] A J Mead et al. “hmcode-2020: improved modelling of non-linear cosmological power spectra with baryonic feedback”. In: *Monthly Notices of the Royal Astronomical Society* 502.1 (Jan. 2021), pp. 1401–1422. ISSN: 0035-8711. DOI: 10.1093/mnras/stab082. eprint: <https://academic.oup.com/mnras/article-pdf/502/1/1401/42438464/stab082.pdf>. URL: <https://doi.org/10.1093/mnras/stab082>.
- [44] Niayesh Afshordi, Yeong-Shang Loh, and Michael A. Strauss. “Cross-correlation of the cosmic microwave background with the 2MASS galaxy survey: Signatures of dark energy, hot gas, and point sources”. In: *Phys. Rev. D* 69 (8 Apr. 2004), p. 083524. DOI: 10.1103/PhysRevD.69.083524. URL: <https://link.aps.org/doi/10.1103/PhysRevD.69.083524>.
- [45] Eric V. Linder. “Exploring the Expansion History of the Universe”. In: *Phys. Rev. Lett.* 90 (9 Mar. 2003), p. 091301. DOI: 10.1103/PhysRevLett.90.091301. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.90.091301>.
- [46] Chao-Jun Feng et al. “A new class of parametrization for dark energy without divergence”. In: *Journal of Cosmology and Astroparticle Physics* 2012.09 (Sept. 2012), p. 023. DOI: 10.1088/1475-7516/2012/09/023. URL: <https://dx.doi.org/10.1088/1475-7516/2012/09/023>.

- [47] A. Hernández-Almada et al. “Stability analysis and constraints on interacting viscous cosmology”. In: *Phys. Rev. D* 101 (6 Mar. 2020), p. 063516. DOI: 10.1103/PhysRevD.101.063516. URL: <https://link.aps.org/doi/10.1103/PhysRevD.101.063516>.
- [48] A. Amariti et al. “Brane cosmology and the self-tuning of the cosmological constant”. In: *Journal of Cosmology and Astroparticle Physics* 2019.10 (Oct. 2019), p. 007. DOI: 10.1088/1475-7516/2019/10/007. URL: <https://dx.doi.org/10.1088/1475-7516/2019/10/007>.
- [49] Marc Kamionkowski and Adam G. Riess. “The Hubble Tension and Early Dark Energy”. In: *Annual Review of Nuclear and Particle Science* 73.1 (2023), pp. 153–180. DOI: 10.1146/annurev-nucl-111422-024107. eprint: <https://doi.org/10.1146/annurev-nucl-111422-024107>. URL: <https://doi.org/10.1146/annurev-nucl-111422-024107>.
- [50] Planck Collaboration et al. “Planck 2018 results - V. CMB power spectra and likelihoods”. In: *Astronomy & Astrophysics* 641 (2020), A5. DOI: 10.1051/0004-6361/201936386. URL: <https://doi.org/10.1051/0004-6361/201936386>.
- [51] M. Betoule et al. “Improved cosmological constraints from a joint analysis of the SDSS-II and SNLS supernova samples”. In: *Astronomy & Astrophysics* 568, A22 (Aug. 2014), A22. DOI: 10.1051/0004-6361/201423413. arXiv: 1401.4064 [astro-ph.CO].
- [52] Julian E. Bautista et al. “Measurement of baryon acoustic oscillation correlations at  $z = 2.3$  with SDSS DR12 Ly $\alpha$ -Forests”. In: *Astronomy & Astrophysics* 603, A12 (June 2017), A12. DOI: 10.1051/0004-6361/201730533. arXiv: 1702.00176 [astro-ph.CO].
- [53] Mark S. Linton et al. “Variable sound speed in interacting dark energy models”. In: *Journal of Cosmology and Astroparticle Physics* 2018.04 (Apr. 2018), p. 043. DOI: 10.1088/1475-7516/2018/04/043. URL: <https://dx.doi.org/10.1088/1475-7516/2018/04/043>.
- [54] Mark S Linton et al. “Variable sound speed in interacting dark energy models”. In: *J. Cosmol. Astropart. Phys.* 2018.04 (Apr. 2018), pp. 043–043.
- [55] O. Tange. “GNU Parallel 2018”. In: (Mar. 2018). URL: <https://doi.org/10.5281/zenodo.1146014>.
- [56] Jesús Torrado and Antony Lewis. *Cobaya: Bayesian analysis in cosmology*. Astrophysics Source Code Library, record ascl:1910.019. Oct. 2019. ascl: 1910.019.
- [57] Jesús Torrado and Antony Lewis. “Cobaya: code for Bayesian analysis of hierarchical physical models”. In: *Journal of Cosmology and Astroparticle Physics* 2021.05 (May 2021), p. 057. DOI: 10.1088/1475-7516/2021/05/057. URL: <https://doi.org/10.1088/1475-7516/2021/05/057>.
- [58] George Casella and Edward I. George. “Explaining the Gibbs Sampler”. In: *The American Statistician* 46.3 (1992), pp. 167–174. ISSN: 00031305. URL: <http://www.jstor.org/stable/2685208> (visited on 08/07/2023).