

1. Consider the curve $f(x) = x^3$. Let $y = mx + b$ be a tangent.

(a) Expand $(x-A)(x-B)^2$.

(b) Find the x -coordinate of the point of contact in terms of m .

(c) Hence, find the derivative of $f(x) = x^3$.

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

JE MATHS

$$\begin{aligned}
 1. \quad (a) \quad (x-A)(x-B)^2 &= (x-A)(x^2 - 2Bx + B^2) \\
 &= x^3 - 2Bx^2 + B^2x - Ax^2 + 2ABx - AB^2 \\
 &= x^3 + (-A-2B)x^2 + (2AB+B^2)x - AB^2
 \end{aligned}$$


(b) Solve $\begin{cases} y = x^3 \\ y = mx + b \end{cases}$ for the x -coordinate of point of contact of the tangent and parabola.

$$x^3 = mx + b \Rightarrow x^3 - mx - b = 0$$

The solution of the equation $x^3 - mx - b = 0$ gives the x -coordinate of point of contact.

The equation $x^3 - mx - b = 0$ can be expressed in the form of $(x-A)(x-B)^2 = 0$ as $x^3 - mx - b = x^3 + (-A-2B)x^2 + (2AB+B^2)x - AB^2$ by finding proper values of coefficients A and B .

Hence, the solution of $x^3 - mx - b = 0$, which representing the x -coordinate of point of contact, can be obtained from the solution of the equation $(x-A)(x-B)^2 = 0$, where $x = B$ is the x -coordinate of point of contact and $x = A$ is the x -coordinate of point of intersection between the tangent and parabola.



$$\text{Let } x^3 - mx - b = x^3 + (-A - 2B)x^2 + (2AB + B^2)x - AB^2.$$

By matching coefficients,

$$-A - 2B = 0 \quad (1)$$

$$2AB + B^2 = -m \quad (2)$$

$$AB^2 = -b \quad (3)$$

From (1),

$$A = -2B$$

Sub $A = -2B$ into (2):

$$-4B^2 + B^2 = -m$$

$$B^2 = \frac{m}{3}$$

$$x = B = \pm \sqrt{\frac{m}{3}}$$

$$(c) \quad x = \pm \sqrt{\frac{m}{3}} \Rightarrow x^2 = \frac{m}{3}$$

$$\Rightarrow m = 3x^2$$

$$\Rightarrow f'(x) = 3x^2$$

