Sarcoma epigenomic data has accumulated rapidly, and the modality mix is broadening

87% of all deposits fall in the last ten years and 50% in the last five. Assays other than ChIP-seq and methylation arrays rose from 6% of deposits in 2011–2015 to 33% in 2021–2026.



Cumulative count by the deposition year of the containing GEO series, de-duplicated. 2026 is a partial year, so every curve is flat at its right edge by construction.

Entities with any epigenomic data: 8 of 45 by 2010, 21 by 2015, 44 by 2020, 44 today.

Year each modality first passed 50 deposited samples: ChIP-seq / ChIP-chip 2007; DNA methylation 2010; Accessibility (ATAC / DNase) 2011; 3D genome 2014; CUT&RUN / CUT&Tag 2021; Single-cell epigenome 2024.

The six families cover 15,950 of the 16,337 dated samples; the remaining 387 are Repli-seq, which belongs to none of them.

Controlled-access deposits are not in this figure: EGA and dbGaP do not publish a comparable deposition timeline.

Dashed segments project the 6 months after the harvest cutoff (01 September 2026, dotted rule), resampling whole months from the preceding 24; the shaded wedge is the 10th to 90th percentile of 20,000 such draws. Across all six families the projection is 895 further samples, 80% interval 619 to 1,308. The interval is wide because deposits arrive as whole series: the median sarcoma series is 9 samples and the largest 1% hold 16% of everything, so a single large resource landing inside the window moves the total by several hundred.