
Regulatory epigenomics tracks research history, not disease burden
No regulatory epigenomics whatsoever
· Atypical fibroxanthoma / pleomorphic d (1,710/yr)
· Dermatofibrosarcoma protuberans (1,400/yr)
· Angiosarcoma (1,312/yr)
· Myxofibrosarcoma (820/yr)
· Liposarcoma, pleomorphic (185/yr)
· Inflammatory myofibroblastic tumor (175/yr)
· Extraskeletal myxoid chondrosarcoma (75/yr)
· Intimal sarcoma (60/yr)
· Myoepithelial carcinoma of soft tissue (50/yr)
· Angiomatoid fibrous histiocytoma (42/yr)
· Clear cell sarcoma of the kidney (20/yr)
· BCOR::CCNB3 sarcoma (15/yr)
Regulatory epigenomic samples in public archives
(ChIP-seq / CUT&RUN / ATAC / DNase / Hi-C / HiChIP)
10,000
1,000
100
10
1
0
Osteosarcoma
Ewing sarcoma
GIST
Kaposi sarcoma
Fibrosarcoma
Rhabdoid tumor / ATRT
Leiomyosarcoma
ASPS
Undifferentiated pleomorphic sarcoma
Liposarcoma, well-differentiated
Chondrosarcoma
MYOD1-mutant RMS
Infantile fibrosarcoma
Solitary fibrous tumor
PEComa
Desmoid / aggressive fibromatosis
Pediatric-predominant
Both ages
Adult-predominant
ZERO
10
100
1,000
10,000
US cases per year (all ages)