


25 of 45 sarcoma entities have never had a single patient's tumour profiled for active chromatin, chromatin accessibility or 3D genome architecture

Leiomyosarcoma	✗	none, against ~3,310 US cases per year
Desmoid / aggressive fibromatosis	✗	none, against ~1,740 US cases per year
Dermatofibrosarcoma protuberans	✗	none, against ~1,400 US cases per year
Angiosarcoma	✗	none, against ~1,312 US cases per year
Chondrosarcoma	✗	none, against ~1,280 US cases per year
Kaposi sarcoma	✗	none, against ~1,268 US cases per year
Solitary fibrous tumour	✗	none, against ~1,200 US cases per year
Liposarcoma, well-differentiated	✗	none, against ~1,100 US cases per year
Myxofibrosarcoma	✗	none, against ~820 US cases per year
Endometrial stromal sarcoma	✗	none, against ~328 US cases per year
LGFMS / SEF	✗	none, against ~229 US cases per year
Liposarcoma, pleomorphic	✗	none, against ~185 US cases per year
Inflammatory myofibroblastic tumour	✗	none, against ~175 US cases per year
Epithelioid sarcoma	✗	none, against ~156 US cases per year
Epithelioid haemangioendothelioma	✗	none, against ~110 US cases per year
Extraskeletal myxoid chondrosarcoma	✗	none, against ~75 US cases per year
Intimal sarcoma	✗	none, against ~60 US cases per year
PEComa (malignant)	✗	none, against ~25 US cases per year
Clear cell sarcoma of the kidney	✗	none, against ~20 US cases per year
BCOR-CCNB3 sarcoma	✗	none, against ~15 US cases per year
Infantile fibrosarcoma	✗	none, against ~13 US cases per year
Myoepithelial carcinoma of soft tissue	✗	none (no population rate published anywhere)
Ossifying fibromyxoid tumour	✗	none (no population rate published anywhere)
Angiomatoid fibrous histiocytoma	✗	none (no population rate published anywhere)
Atypical fibroxanthoma / pleomorphic dermal sarcoma	✗	none (no population rate published anywhere)



Regulatory epigenomic profiles of PATIENT-DERIVED MATERIAL — tumour tissue, PDX or patient-derived organoids (ChIP-seq / CUT&RUN / ATAC / Hi-C; input and IgG controls excluded)