

EECS C106A/206A

Discussion #2: Forward Kinematics

Agenda

- Logistics
- Lecture Review
 - Homogeneous Transforms
 - Forward Kinematics

Logistics

- Upcoming:
 - Homework 1 due **9/6**
 - Homework 2 due **9/13**
 - Midterm 1 on **9/27**
- Office Hours
 - Started this week!
 - Tuesdays & Thursdays @ 11:30 - 12:30, Locations on Piazza
 - By appointment: brentyi@berkeley.edu


Logistics

- Discussions
 - Non-comprehensive review of lecture material
 - Attendance not required
- New GSI
 - Andrew Barkan
 - Mechanical Engineering PhD Student
 - Running discussions every other week, starting next week
 - He's great and you'll love him!

Homogeneous Transforms

Recall: Rotation Matrices


- Allow us to write a rotation as a matrix multiplication



$$q_A = R_{AB} q_B$$

Rigid Body Motion

- Translation of origin,
relative rotation



$$q_A = R_{AB}q_B + p_{AB}$$

$$SE(3) = \{(p, R) : p \in \mathbb{R}^3, R \in SO(3)\}$$

Homogeneous Coordinates

For all points:

$$\bar{q}_A = \begin{bmatrix} q_A \\ 1 \end{bmatrix}$$

Homogeneous Transforms

Homogeneous coordinates
allow us to represent both a
rotation and a translation
using a single matrix
multiply.

$$q_A = R_{AB}q_B + p_{AB}$$

$$\begin{bmatrix} q_A \\ 1 \end{bmatrix} = \begin{bmatrix} R_{AB} & p_{AB} \\ \vec{0}^T & 1 \end{bmatrix} \begin{bmatrix} q_B \\ 1 \end{bmatrix}$$

$$\bar{q}_A = \begin{bmatrix} R_{AB} & p_{AB} \\ \vec{0}^T & 1 \end{bmatrix} \bar{q}_B$$

Homogeneous Transforms

This can be written
alternatively as:

$$g_{AB} = \begin{bmatrix} R_{AB} & p_{AB} \\ \vec{0}^T & 1 \end{bmatrix}$$

$$\bar{q}_A = g_{AB} \bar{q}_B$$

Chaining Transforms

$$\bar{q}_A = \begin{bmatrix} R_{AB} & p_{AB} \\ \vec{0}^T & 1 \end{bmatrix} \begin{bmatrix} R_{BC} & p_{BC} \\ \vec{0}^T & 1 \end{bmatrix} \bar{q}_C$$

$$\bar{q}_A = g_{AB} g_{BC} \bar{q}_C$$

Inverting Transforms

~~$$g^{-1} = \begin{bmatrix} R & p \\ \vec{0}^T & 1 \end{bmatrix}^{-1} = \begin{bmatrix} R^T & -R^T p \\ \vec{0}^T & 1 \end{bmatrix}$$~~

$$g^{-1} = \begin{bmatrix} R & p \\ \vec{0}^T & 1 \end{bmatrix}^{-1} = \begin{bmatrix} R^T & -R^T p \\ \vec{0}^T & 1 \end{bmatrix}$$


Questions so far?

Forward Kinematics

Given a set of joints and their positions, how do we find the configuration of the end effector?


Joints: Rotational and Prismatic

- *Rotational joints*: second frame moves rotationally relative to first one
- *Prismatic joints*: second frame moves linearly relative to first one




Joints

- What kind of joints does a Sawyer have?
- How many? Why?



Forward Kinematics

- We can describe the configuration of the end effector with a homogeneous transform \mathbf{g}
- **FK:** given joint angles, we want to find a mapping from the initial transform to a final transform



Twists

We can characterize the kinematics of a joint with a twist:

$$\xi = \begin{bmatrix} v \\ \omega \end{bmatrix} \quad \begin{array}{l} v \in \mathbb{R}^3 : \text{linear component} \\ \omega \in \mathbb{R}^3 : \text{angular component} \end{array}$$

Twists

In 106A, we mostly care about two kinds of twists:

- Purely translational:

- Axis \mathbf{v}

$$\xi = \begin{bmatrix} v \\ \vec{0} \end{bmatrix}$$

- Purely rotational:

- Axis of rotation $\boldsymbol{\omega}$

- Any point along $\boldsymbol{\omega}, \mathbf{q}$

$$\xi = \begin{bmatrix} -\boldsymbol{\omega} \times \mathbf{q} \\ \boldsymbol{\omega} \end{bmatrix}$$

- Normalized axes

Twist Operators

We define two operators for use in the exponential:

- “Wedge”


$$\begin{aligned}\hat{\xi} &= \begin{bmatrix} v \\ \omega \end{bmatrix}^{\wedge} \\ &= \begin{bmatrix} \hat{\omega} & v \\ 0 & 0 \end{bmatrix}\end{aligned}$$

- “Vee”

$$(\hat{\xi})^{\vee} = \xi$$

Exponential Representation (Revolute)

Consider point \mathbf{p} , rotating about axis $\boldsymbol{\omega}$.



Exponential Representation (Revolute)

We can describe the motion of \mathbf{p} with:


$$\dot{\mathbf{p}}(t) = \boldsymbol{\omega} \times (\mathbf{p}(t) - \mathbf{q})$$

Which can equivalently be written as:

$$\begin{aligned} \begin{bmatrix} \dot{p} \\ 0 \end{bmatrix} &= \begin{bmatrix} \hat{\boldsymbol{\omega}} & -\boldsymbol{\omega} \times \mathbf{q} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} p \\ 1 \end{bmatrix} \\ &= \hat{\boldsymbol{\xi}} \begin{bmatrix} p \\ 1 \end{bmatrix} \end{aligned}$$

The solution for which is:

$$\bar{\mathbf{p}}(\theta) = e^{\hat{\boldsymbol{\xi}}\theta} \bar{\mathbf{p}}(0)$$



Exponential Representation

Does this form look familiar?

$$p(\theta) = \underline{e^{\hat{\xi}\theta}} p(0)$$

Similar to exponential computed by Rodrigues' Formula:

$$\underline{e^{\hat{\omega}\theta}} = I_3 + \frac{\hat{\omega}}{\|\omega\|_2} \sin(\|\omega\|\theta) + \frac{\hat{\omega}^2}{\|\omega\|_2^2} (1 - \cos(\|\omega\|\theta))$$

...but generalized to all twists instead of just rotations.

Exponential Rigid Body Motion

- Given a homogeneous transform representing the initial configuration of the end effector (**World -> End Effector**):

$$g_{WE}(0) = \begin{bmatrix} R_{WE} & p_{WE} \\ \vec{0}^T & 1 \end{bmatrix}$$


- We can compute a new transform parameterized by our joint angle as:

$$g_{WE}(\theta) = e^{\hat{\xi}\theta} g_{WE}(0)$$

Exponential Representations

- This also works for multiple joints:

$$g(\theta) = e^{\hat{\xi}_1 \theta_1} e^{\hat{\xi}_2 \theta_2} g(0)$$



Exponential Solution

$$e^{\hat{\xi}\theta} = \begin{bmatrix} e^{\hat{\omega}\theta} & (I - e^{\hat{\omega}\theta})(\omega \times v) + \omega\omega^T v\theta \\ \mathbf{0} & 1 \end{bmatrix}$$

$$e^{\hat{\omega}\theta} = I_3 + \frac{\hat{\omega}}{\|\omega\|_2} \sin(\|\omega\|\theta) + \frac{\hat{\omega}^2}{\|\omega\|_2^2} (1 - \cos(\|\omega\|\theta))$$