

- Polyhedral cuts
 - warehouse location
 - node covering


Discrete Optimization

Mixed Integer Programming: Part IV

2

Thursday 11 June 15


Convex Hull



4

Thursday 11 June 15


Convex Hull



5

Thursday 11 June 15


Convex Hull



6

Thursday 11 June 15


Convex Hull



7

Thursday 11 June 15


Convex Hull



8

Thursday 11 June 15


Convex Hull



9


Thursday 11 June 15

Convex Hull




3

Thursday 11 June 15



10

Tuesday, 11 June 19



11

Tuesday, 11 June 19

- **Polyhedral cuts**
 - cuts that represent the facets of the convex hull of the integer solution

12

Tuesday, 11 June 19

Polyhedral Cuts

- **Polyhedral cuts**
 - cuts that represent the facets of the convex hull of the integer solution
- **These cuts are valid**
 - they do not remove any solution

12

Tuesday, 11 June 19

Polyhedral Cuts

- **Polyhedral cuts**
 - cuts that represent the facets of the convex hull of the integer solution
- **These cuts are valid**
 - they do not remove any solution
- **The cuts are as strong as possible**
 - if we have all of them, we could use linear programming to solve the problem

12

Tuesday, 11 June 19

Polyhedral Cuts

- **They exploit the problem structure**
 - they are derived from the structure of constraints
 - not based on information in the tableau

13

Tuesday, 11 June 19

Polyhedral Cuts

- **They exploit the problem structure**
 - they are derived from the structure of constraints
 - not based on information in the tableau
- **They share some of the spirit of syntactic cuts**
 - validity
 - must cut the current basic feasible solution
 - do not need to generate all of them

13

Tuesday, 11 June 19

Polyhedral Cuts

- **They exploit the problem structure**
 - they are derived from the structure of constraints
 - not based on information in the tableau
- **They share some of the spirit of syntactic cuts**
 - validity
 - must cut the current basic feasible solution
 - do not need to generate all of them
- **An application may use multiple cut types**
 - exploit different substructure

13

Tuesday, 11 June 19

What is a Facet?

- **To find an facet in \mathbb{R}^n**
 - find n affinely independent solutions (points)

14

Tuesday, 11 June 19

What is a Facet?

- **To find an facet in \mathbb{R}^n**
– find n affinely independent solutions (points)
- **Affine independence**
 x_1, \dots, x_n are affinely independent iff $(x_1, 1), \dots, (x_n, 1)$ are linearly independent.

14

Facility 11, Lect 10

What is a Facet?

- **To find an facet in \mathbb{R}^n**
– find n affinely independent solutions (points)
- **Affine independence**
 x_1, \dots, x_n are affinely independent iff $(x_1, 1), \dots, (x_n, 1)$ are linearly independent.
- **Linear independence**
 x_1, \dots, x_n are linearly independent iff $\alpha_1 x_1 + \dots + \alpha_n x_n = 0$ implies that $\alpha_i = 0$ for all i .

14

Facility 11, Lect 10

Warehouse Location

$$\begin{aligned} \min \quad & \sum_{w \in W} c_w x_w + \sum_{w \in W, c \in C} t_{w,c} y_{w,c} \\ \text{subject to} \quad & y_{w,c} \leq x_w \quad (w \in W, c \in C) \\ & \sum_{w \in W} y_{w,c} = 1 \quad (c \in C) \\ & x_w \in \{0, 1\} \quad (w \in W) \\ & y_{w,c} \in \{0, 1\} \quad (w \in W, c \in C) \end{aligned}$$

16

Facility 11, Lect 10


Warehouse Location

$$\begin{aligned} \min \quad & \sum_{w \in W} c_w x_w + \sum_{w \in W, c \in C} t_{w,c} y_{w,c} \\ \text{subject to} \quad & y_{w,c} \leq x_w \quad (w \in W, c \in C) \\ & \sum_{w \in W} y_{w,c} = 1 \quad (c \in C) \\ & x_w \in \{0, 1\} \quad (w \in W) \\ & y_{w,c} \in \{0, 1\} \quad (w \in W, c \in C) \end{aligned}$$

16

Facility 11, Lect 10

Warehouse Location



15

Facility 11, Lect 10

Facets for Warehouse Location

- Are these inequalities facets?
 $y_{w,c} \leq x_w$

17

Facility 11, Lect 10

Facets for Warehouse Location

- Are these inequalities facets?
 $y_{w,c} \leq x_w$
- Consider $y_{w,1} \leq x_w$ and the following n points


w	1	2	3	...	n
$y_{w,1}$	0	0	0	...	0
x_w	1	0	0	...	0
$y_{w,2}$	1	1	0	...	0
x_w	1	1	1	...	0
$y_{w,3}$	1	1	1	...	0
x_w	1	1	1	...	0
$y_{w,n}$	1	1	1	...	1
x_w	1	1	1	...	1

17

Facility 11, Lect 10

Node Packing

- Let $G=(V,E)$ be a graph.
– A node packing is a subset W of V such that no two nodes in W are connected by an edge. The goal is to find the node packing of maximal size.




- How do we express it as a MIP?

18

Facility 11, Lect 10

Node Packing

- Let $G=(V,E)$ be a graph.
– A node packing is a subset W of V such that no two nodes in W are connected by an edge. The goal is to find the node packing of maximal size.



- How do we express it as a MIP?


18

Facility 11, Lect 10

Node Packing

► Let $G=(V,E)$ be a graph.

- A node packing is a subset W of V such that no two nodes in W are connected by an edge. The goal is to find the node packing of maximal size.



► How do we express it as a MIP?


18

Forming 11, Lect 10

Node Packing

► Let $G=(V,E)$ be a graph.

- A node packing is a subset W of V such that no two nodes in W are connected by an edge. The goal is to find the node packing of maximal size.



► How do we express it as a MIP?

19

Forming 11, Lect 10

Node Packing

$$\begin{aligned} \max \quad & x_1 + \dots + x_6 \\ \text{s.t.} \quad & x_1 + x_2 \leq 1 \\ & x_1 + x_3 \leq 1 \\ & x_1 + x_4 \leq 1 \\ & x_1 + x_5 \leq 1 \\ & x_2 + x_3 \leq 1 \\ & x_2 + x_6 \leq 1 \\ & x_3 + x_4 \leq 1 \\ & x_4 + x_5 \leq 1 \\ & x_5 + x_6 \leq 1 \\ & x_i \in \{0,1\} \end{aligned}$$

Node Packing

$$\begin{aligned} \max \quad & x_1 + \dots + x_6 \\ \text{s.t.} \quad & x_1 + x_2 \leq 1 \\ & x_1 + x_3 \leq 1 \\ & x_1 + x_4 \leq 1 \\ & x_1 + x_5 \leq 1 \\ & x_1 + x_6 \leq 1 \\ & x_2 + x_3 \leq 1 \\ & x_3 + x_4 \leq 1 \\ & x_4 + x_5 \leq 1 \\ & x_5 + x_6 \leq 1 \\ & 0 \leq x_1 \dots x_6 \leq 1 \end{aligned}$$

20

Forming 11, Lect 10

Node Packing

$$\begin{aligned} \max \quad & x_1 + \dots + x_6 \\ \text{s.t.} \quad & x_1 + x_2 \leq 1 \\ & x_1 + x_3 \leq 1 \\ & x_1 + x_4 \leq 1 \\ & x_1 + x_5 \leq 1 \\ & x_1 + x_6 \leq 1 \\ & x_2 + x_3 \leq 1 \\ & x_3 + x_4 \leq 1 \\ & x_4 + x_5 \leq 1 \\ & x_5 + x_6 \leq 1 \\ & 0 \leq x_1 \dots x_6 \leq 1 \end{aligned}$$

► What does the linear relaxation produce?

$$x_1 = \frac{1}{2}, \dots, x_6 = \frac{1}{2}$$

20

Forming 11, Lect 10

Node Packing

$$\begin{aligned} \max \quad & x_1 + \dots + x_6 \\ \text{s.t.} \quad & x_1 + x_2 \leq 1 \\ & x_1 + x_3 \leq 1 \\ & x_1 + x_4 \leq 1 \\ & x_1 + x_5 \leq 1 \\ & x_1 + x_6 \leq 1 \\ & x_2 + x_3 \leq 1 \\ & x_3 + x_4 \leq 1 \\ & x_4 + x_5 \leq 1 \\ & x_5 + x_6 \leq 1 \\ & 0 \leq x_1 \dots x_6 \leq 1 \end{aligned}$$

► What does the linear relaxation produce?

$$x_1 = \frac{1}{2}, \dots, x_6 = \frac{1}{2}$$

20


Forming 11, Lect 10

How do we Find Facets?

► Find a property of the solution

- constraints satisfied by all solutions

► Consider a clique



21


Forming 11, Lect 10

How do we Find Facets?

► Find a property of the solution

- constraints satisfied by all solutions

► Consider a clique




► How many nodes can be in a packing?

21

Forming 11, Lect 10


26

Tuesday, 11 June 19




27

Tuesday, 11 June 19




27

Tuesday, 11 June 19




27

Tuesday, 11 June 19




28

Tuesday, 11 June 19



28

Tuesday, 11 June 19



- With the best model from before

$$obj = 0.5 \quad \text{Need at least 2 colors,}$$

- Add the 3-Clique

$$\sum_{c \in \{0,3,4\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 3$$

$$obj = 1.0 \quad \text{Still at least 2 colors,}$$

28

Tuesday, 11 June 19

- With the best model from before

$$obj = 0.5 \quad \text{Need at least 2 colors,}$$

- Add the 3-Clique

$$\sum_{c \in \{0,3,4\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 3$$

$$obj = 1.0 \quad \text{Still at least 2 colors,}$$

- Add the 4-Clique

$$\sum_{c \in \{0,2,5,5\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 6$$

28

Tuesday, 11 June 19

- With the best model from before

$$obj = 0.5 \quad \text{Need at least 2 colors,}$$

- Add the 3-Clique

$$\sum_{c \in \{0,3,4\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 3$$

$$obj = 1.0 \quad \text{Still at least 2 colors,}$$

- Add the 4-Clique

$$\sum_{c \in \{0,2,3,5\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 6$$

$$obj = 1.5 \quad \text{Need at least 3 colors!!}$$

► With the best model from before

$$obj = 0.5 \quad \text{Need at least 2 colors.}$$

► Add the 3-Clique

MIP:

Optimal - 9 nodes
Proof - 41 nodes

$$\sum_{c \in \{0,3,4\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 3$$

$$obj = 1.0$$

► Add the 4-Clique

$$\sum_{c \in \{0,2,3,5\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 6$$

$$obj = 1.5 \quad \text{Need at least 3 colors!!!}$$

► With the best model from before

$$obj = 0.5 \quad \text{Need at least 2 colors.}$$

► Add the 3-Clique

MIP:

Optimal - 9 nodes
Proof - 41 nodes

$$\sum_{c \in \{0,3,4\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 3$$

$$obj = 1.0$$

► Add the 4-Clique

$$\sum_{c \in \{0,2,3,5\}} \sum_{v=0}^3 v \times \text{color}_{c,v} \geq 6$$

$$obj = 1.5 \quad \text{Need at least 3 colors!!!}$$