

Some symbolic tools for the Fox H -function

Le Chen

Department of Mathematics and Statistics

Auburn University

le.chen@auburn.edu, chenle02@gmail.com

December 16, 2023

Contents

1	Introduction	1
2	Examples	5
2.1	Example FoxH-2_9_11.wls	5
2.2	Example FoxH-2_9_12.wls	6
2.3	Example FoxH-2_9_13.wls	7
2.4	Example FoxH-2_9_4.wls	8
2.5	Example FoxH-2_9_5.wls	9
2.6	Example FoxH-2_9_6.wls	10
2.7	Example FoxH32-21.wls	11
2.8	Example FoxH32-21-Y.wls	12
2.9	Example FoxH32-21-Z.wls	13
2.10	Example FoxH32-21-Z-Star.wls	15
2.11	Example FoxH-Bessel-J_2_9_18.wls	16
2.12	Example FoxH-Bessel-K_2_9_19.wls	17
2.13	Example FoxH-Bessel-Y_2_9_20.wls	18
2.14	Example FoxH-Cos.wls	19
2.15	Example FoxH-H.G_2_9_14.wls	20
2.16	Example FoxH-H.G_2_9_15.wls	21
2.17	Example FoxH-Lommel_2_9_22.wls	22
2.18	Example FoxH-Mittag-Leffler.wls	24
2.19	Example FoxH-Sin.wls	25
2.20	Example FoxH-Whittaker_2_9_21.wls	26
	References	27

1 Introduction

In this note, we explain the code for checking the conditions of the Fox H -function [Fox61]. Here we follow the notation from Kilbas and Saigo [KS].

Let m, n, p, q be configure integers such that

$$0 \leq m \leq q \quad \text{and} \quad 0 \leq n \leq p.$$

Let $a_i, b_j \in \mathbb{C}$ and $\alpha_i, \beta_j \in \mathbb{R}_+$ be the parameters given below:

$\in (\mathbb{C}, \mathbb{R}_+)$	Front list	Rear list	
p	$(a_1, \alpha_1), \dots, (a_n, \alpha_n)$	$(a_{n+1}, \alpha_{n+1}), \dots, (a_p, \alpha_p)$	Upper list
q	$(b_1, \beta_1), \dots, (b_m, \beta_m)$	$(b_{m+1}, \beta_{m+1}), \dots, (b_q, \beta_q)$	Lower list

and denote

$$\mathcal{H}_{p,q}^{m,n}(s) := \frac{\prod_{i=1}^n \Gamma(1 - a_i - \alpha_i s)}{\prod_{j=n+1}^p \Gamma(a_j + \alpha_j s)} \times \frac{\prod_{j=1}^m \Gamma(b_j + \beta_j s)}{\prod_{j=m+1}^q \Gamma(1 - b_j - \alpha_j s)}. \quad (1)$$

Then the Fox H -function $H_{2,3}^{2,1} \left(z \left| \begin{array}{c} \dots \\ \dots \end{array} \right. \right)$ is defined by a Mellin-Barnes type integral of the form

$$H_{m,n}^{p,q} \left(z \left| \begin{array}{c} (a_1, \alpha_1), \dots, (a_p, \alpha_p) \\ (b_1, \beta_1), \dots, (b_q, \beta_q) \end{array} \right. \right) := \frac{1}{2\pi i} \int_{\mathcal{L}} H_{p,q}^{m,n}(s) z^{-s} ds. \quad (2)$$

The basic assumption for the well-posedness of the Fox H -function is that two sets of poles do not overlap, i.e.,

$$\left\{ b_{j\ell} = \frac{-b_j - \ell}{\beta_j}, \ell = 0, 1, \dots \right\} \cap \left\{ a_{ik} = \frac{1 - a_i + k}{\alpha_i}, k = 0, 1, \dots \right\} = \emptyset. \quad (3)$$

The contour \mathcal{L} in (2) is given by one of the following three cases:

1. $\mathcal{L} = \mathcal{L}_{-\infty}$ is a left loop situated in a horizontal strip starting at point $-\infty + i\phi_1$ and terminating at point $-\infty + i\phi_2$ for some $-\infty < \phi_1 < \phi_2 < \infty$;
2. $\mathcal{L} = \mathcal{L}_{+\infty}$ is a right loop situated in a horizontal strip starting at point $+\infty + i\phi_1$ and terminating at point $+\infty + i\phi_2$ for some $-\infty < \phi_1 < \phi_2 < \infty$;
3. $\mathcal{L} = \mathcal{L}_{i\gamma\infty}$ is a contour starting at point $\gamma - i\infty$ and terminating at point $\gamma + i\infty$ for some $\gamma \in (-\infty, \infty)$.

We need a set of conditions to ensure the convergence of the integral in (2). To explain this, we need to introduce some notation (following p. 2 of [KS]). First denote

$$a_1^* := \sum_{j=1}^m \beta_j - \sum_{i=n+1}^p \alpha_i,$$

$$a_2^* := \sum_{i=1}^n \alpha_i - \sum_{j=m+1}^q \beta_j.$$

The following two parameters play the most important role:

$$a^* := a_2^* + a_1^* = \sum_{i=1}^n \alpha_i - \sum_{i=n+1}^p \alpha_i + \sum_{j=1}^m \beta_j - \sum_{j=m+1}^q \beta_j;$$

$$\Delta := a_2^* - a_1^* = \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i.$$

Similar to a^* , we define

$$\xi := \sum_{i=1}^n a_i - \sum_{i=n+1}^p a_i + \sum_{j=1}^m b_j - \sum_{j=m+1}^q b_j.$$

Additionally, set

$$c^* := m + n - \frac{p+q}{2}.$$

In the critical cases, we need to use the following two parameters:

$$\delta := \prod_{i=1}^p \alpha_i^{-\alpha_i} \prod_{j=1}^q \beta_j^{\beta_j};$$

$$\mu := \sum_{j=1}^q b_j - \sum_{i=1}^p a_i + \frac{p-q}{2}$$

The well-posedness of the Fox H -function is given by Theorems 1.1 and 1.2 of [KS], which are summarized in the following figure 2:

2 Examples

2.1 Example FoxH-2_9_11.wls

File content

```
(* (2.9.11) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {{1,1},{1,1}}
```




Figure 1: Diagram for the parameterization of the Fox H -function.




Figure 2: Well-posedness of the Fox H -function.

```

},
(* Lower List *) {
  (* Lower Front List *) {{1,1}},
  (* Lower Rear List *) {{0,0}}
}
}

```

Fox H-function

$$H_{2,2}^{1,0} \left(\cdot \left| \begin{array}{c} (1,1), (1,1) \\ (1,1), (0,0) \end{array} \right. \right)$$

$$H_{2,2}^{1,0} \left(\cdot \left| \begin{array}{c|c} & (1,1), (1,1) \\ \hline (1,1) & (0,0) \end{array} \right. \right)$$

Summary

$$\begin{aligned}
a^* &= -1 \\
\Delta &= -1 \\
\delta &= \text{Indeterminate} \\
\mu &= -1 \\
a_1^* &= -1 \\
a_2^* &= 0 \\
\xi &= -1 \\
c^* &= -1
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 & -11 \end{array} \right)$$

2.2 Example FoxH-2_9_12.wls

File content

```

(* (2.9.12) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {

```

```

(* Upper Front List *) {{1/2,1},{1/2,1}},
(* Upper Rear List *) {}
},
(* Lower List *) {
(* Lower Front List *) {{0,1}},
(* Lower Rear List *) {{-1/2,1}}
}
}

```

Fox H-function

$$H_{2,2}^{1,2} \left(. \left| \begin{array}{c} (\frac{1}{2}, 1), (\frac{1}{2}, 1) \\ (0, 1), (-\frac{1}{2}, 1) \end{array} \right. \right)$$

$$H_{2,2}^{1,2} \left(. \left| \frac{(\frac{1}{2}, 1), (\frac{1}{2}, 1)}{(0, 1)} \right| \frac{}{(-\frac{1}{2}, 1)} \right)$$

Summary

$$\begin{aligned}
a^* &= 2 \\
\Delta &= 0 \\
\delta &= \text{ComplexInfinity} \\
\mu &= -\frac{3}{2} \\
a_1^* &= 1 \\
a_2^* &= 1 \\
\xi &= \frac{3}{2} \\
c^* &= 1
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} \frac{1}{2} & \frac{3}{2} & \frac{5}{2} & \frac{7}{2} & \frac{9}{2} & \frac{11}{2} & \frac{13}{2} & \frac{15}{2} & \frac{17}{2} & \frac{19}{2} & \frac{21}{2} \\ \frac{1}{2} & \frac{3}{2} & \frac{5}{2} & \frac{7}{2} & \frac{9}{2} & \frac{11}{2} & \frac{13}{2} & \frac{15}{2} & \frac{17}{2} & \frac{19}{2} & \frac{21}{2} \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{array} \right)$$

2.3 Example FoxH-2_9_13.wls

File content

```
(* (2.9.13) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{1/2,1},{1,1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{1/2,1}},
    (* Lower Rear List *) {{0,1}}
  }
}
```

Fox H-function

$$H_{2,2}^{1,2} \left(. \left| \begin{array}{c} (\frac{1}{2}, 1), (1, 1) \\ (\frac{1}{2}, 1), (0, 1) \end{array} \right. \right)$$

$$H_{2,2}^{1,2} \left(. \left| \begin{array}{c} (\frac{1}{2}, 1), (1, 1) \\ (\frac{1}{2}, 1) \end{array} \right| \begin{array}{c} \\ (0, 1) \end{array} \right)$$

Summary

$$\begin{aligned} a^* &= 2 \\ \Delta &= 0 \\ \delta &= \text{ComplexInfinity} \\ \mu &= -1 \\ a_1^* &= 1 \\ a_2^* &= 1 \\ \xi &= 2 \\ c^* &= 1 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} \frac{1}{2} & \frac{3}{2} & \frac{5}{2} & \frac{7}{2} & \frac{9}{2} & \frac{11}{2} & \frac{13}{2} & \frac{15}{2} & \frac{17}{2} & \frac{19}{2} & \frac{21}{2} \\ 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} -\frac{1}{2} & -\frac{3}{2} & -\frac{5}{2} & -\frac{7}{2} & -\frac{9}{2} & -\frac{11}{2} & -\frac{13}{2} & -\frac{15}{2} & -\frac{17}{2} & -\frac{19}{2} & -\frac{21}{2} \end{array} \right)$$

2.4 Example FoxH-2_9_4.wls

File content

```
(* (2.9.4) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{b, β}},
    (* Lower Rear List *) {}
  }
}
```

Fox H-function

$$H_{0,1}^{1,0} \left(\cdot \left| \begin{array}{c} \\ (b, \beta) \end{array} \right. \right)$$

$$H_{0,1}^{1,0} \left(\cdot \left| \begin{array}{c} \hline (b, \beta) \end{array} \right. \right)$$

Summary

$$\begin{aligned} a^* &= \beta \\ \Delta &= \beta \\ \delta &= \text{Indeterminate} \\ \mu &= b - \frac{1}{2} \\ a_1^* &= \beta \\ a_2^* &= 0 \\ \xi &= b \\ c^* &= \frac{1}{2} \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(-\frac{b}{\beta} \quad -\frac{b+1}{\beta} \quad -\frac{b+2}{\beta} \quad -\frac{b+3}{\beta} \quad -\frac{b+4}{\beta} \quad -\frac{b+5}{\beta} \quad -\frac{b+6}{\beta} \quad -\frac{b+7}{\beta} \quad -\frac{b+8}{\beta} \quad -\frac{b+9}{\beta} \quad -\frac{b+10}{\beta} \right)$$

2.5 Example FoxH-2_9_5.wls

File content

```
(* (2.9.5) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{1-a,1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{0, 1}},
    (* Lower Rear List *) {}
  }
}
```

Fox H-function

$$H_{1,1}^{1,1} \left(\cdot \left| \begin{array}{c} (1-a, 1) \\ (0, 1) \end{array} \right. \right)$$

$$H_{1,1}^{1,1} \left(\cdot \left| \frac{(1-a, 1)}{(0, 1)} \right| \right)$$

Summary

$$\begin{aligned} a^* &= 2 \\ \Delta &= 0 \\ \delta &= \text{Indeterminate} \\ \mu &= a - 1 \\ a_1^* &= 1 \\ a_2^* &= 1 \\ \xi &= 1 - a \\ c^* &= 1 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} a & a+1 & a+2 & a+3 & a+4 & a+5 & a+6 & a+7 & a+8 & a+9 & a+10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{array} \right)$$

2.6 Example FoxH-2_9_6.wls

File content

```
(* (2.9.6) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {{α+β+1,1}}
  },
  (* Lower List *) {
    (* Lower Front List *) {{α, 1}},
    (* Lower Rear List *) {}
  }
}
```

Fox H-function

$$H_{1,1}^{1,0} \left(\begin{array}{c} \cdot \\ \left| \begin{array}{c} (\alpha + \beta + 1, 1) \\ (\alpha, 1) \end{array} \right. \end{array} \right)$$

$$H_{1,1}^{1,0} \left(\begin{array}{c} \cdot \\ \left| \frac{\cdot}{(\alpha, 1)} \right| \begin{array}{c} (\alpha + \beta + 1, 1) \\ \hline \end{array} \end{array} \right)$$

Summary

$$\begin{aligned} a^* &= 0 \\ \Delta &= 0 \\ \delta &= \text{Indeterminate} \\ \mu &= -\beta - 1 \\ a_1^* &= 0 \\ a_2^* &= 0 \\ \xi &= -\beta - 1 \\ c^* &= 0 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} -\alpha & -\alpha-1 & -\alpha-2 & -\alpha-3 & -\alpha-4 & -\alpha-5 & -\alpha-6 & -\alpha-7 & -\alpha-8 & -\alpha-9 & -\alpha-10 \end{array} \right)$$

2.7 Example FoxH32-21.wls

File content

```
{
(* Upper List *) {
  (* Upper Front List *) {{1,  $\alpha^{-1}$ }},
  (* Upper Rear List *) {{Ceiling[ $\beta$ ],  $\beta$ }}
},
(* Lower List *) {
  (* Lower Front List *) {{1/2,  $\alpha/2$ }, {1, 1}},
  (* Lower Rear List *) {{1,  $\alpha/2$ }}
}
}
```

Fox H-function

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c} (1, \frac{1}{\alpha}), (\lceil \beta \rceil, \beta) \\ (\frac{1}{2}, \frac{\alpha}{2}), (1, 1), (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c|c} (1, \frac{1}{\alpha}) & (\lceil \beta \rceil, \beta) \\ \hline (\frac{1}{2}, \frac{\alpha}{2}), (1, 1) & (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

Summary

$$\begin{aligned}
a^* &= \frac{1}{\alpha} - \beta + 1 \\
\Delta &= \alpha - \frac{1}{\alpha} - \beta + 1 \\
\delta &= 2^{-\alpha} \left(\frac{1}{\alpha} \right)^{-1/\alpha} \left(2^{\alpha/2} \alpha^{\alpha/2} + \alpha^\alpha \right) \beta^{-\beta} \\
\mu &= 1 - \lceil \beta \rceil \\
a_1^* &= \frac{1}{2} (\alpha - 2\beta + 2) \\
a_2^* &= \frac{1}{\alpha} - \frac{\alpha}{2} \\
\xi &= \frac{3}{2} - \lceil \beta \rceil \\
c^* &= \frac{1}{2}
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \begin{pmatrix} 0 & \alpha & 2\alpha & 3\alpha & 4\alpha & 5\alpha & 6\alpha & 7\alpha & 8\alpha & 9\alpha & 10\alpha \end{pmatrix}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} -\frac{1}{\alpha} & -\frac{3}{\alpha} & -\frac{5}{\alpha} & -\frac{7}{\alpha} & -\frac{9}{\alpha} & -\frac{11}{\alpha} & -\frac{13}{\alpha} & -\frac{15}{\alpha} & -\frac{17}{\alpha} & -\frac{19}{\alpha} & -\frac{21}{\alpha} \\ -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 & -11 \end{pmatrix}$$

2.8 Example FoxH32-21-Y.wls

File content

```

{
  (* Upper List *) {
    (* Upper Front List *) {{1, 1}},
    (* Upper Rear List *) {{β + γ, β}}
  },
  (* Lower List *) {
    (* Lower Front List *) {{d/2, α/2}, {1, 1}},
    (* Lower Rear List *) {{1, α/2}}
  }
}

```

Fox H-function

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c} (1, 1), (\beta + \gamma, \beta) \\ (\frac{d}{2}, \frac{\alpha}{2}), (1, 1), (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

$$H_{2,3}^{2,1} \left(\cdot \left| \frac{(1, 1)}{(\frac{d}{2}, \frac{\alpha}{2}), (1, 1)} \right| \frac{(\beta + \gamma, \beta)}{(1, \frac{\alpha}{2})} \right)$$

Summary

$$\begin{aligned} a^* &= 2 - \beta \\ \Delta &= \alpha - \beta \\ \delta &= 2^{-\alpha} \left(2^{\alpha/2} \alpha^{\alpha/2} + \alpha^\alpha \right) \beta^{-\beta} \\ \mu &= \frac{1}{2} (-2\beta - 2\gamma + d + 1) \\ a_1^* &= \frac{1}{2} (\alpha - 2\beta + 2) \\ a_2^* &= 1 - \frac{\alpha}{2} \\ \xi &= \frac{1}{2} (d - 2(\beta + \gamma - 1)) \\ c^* &= \frac{1}{2} \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} -\frac{d}{\alpha} & -\frac{d+2}{\alpha} & -\frac{d+4}{\alpha} & -\frac{d+6}{\alpha} & -\frac{d+8}{\alpha} & -\frac{d+10}{\alpha} & -\frac{d+12}{\alpha} & -\frac{d+14}{\alpha} & -\frac{d+16}{\alpha} & -\frac{d+18}{\alpha} & -\frac{d+20}{\alpha} \\ -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 & -11 \end{array} \right)$$

2.9 Example FoxH32-21-Z.wls

File content

```
{
  (* Upper List *) {
    (* Upper Front List *) {{1, 1}},
```

```

(* Upper Rear List *) {{Ceiling[β], β}}
},
(* Lower List *) {
(* Lower Front List *) {{d/2, α/2}, {1, 1}},
(* Lower Rear List *) {{1, α/2}}
}
}

```

Fox H-function

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c} (1,1), (\lceil \beta \rceil, \beta) \\ (\frac{d}{2}, \frac{\alpha}{2}), (1,1), (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c|c} (1,1) & (\lceil \beta \rceil, \beta) \\ \hline (\frac{d}{2}, \frac{\alpha}{2}), (1,1) & (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

Summary

$$\begin{aligned}
a^* &= 2 - \beta \\
\Delta &= \alpha - \beta \\
\delta &= 2^{-\alpha} \left(2^{\alpha/2} \alpha^{\alpha/2} + \alpha^\alpha \right) \beta^{-\beta} \\
\mu &= \frac{1}{2} (-2\lceil \beta \rceil + d + 1) \\
a_1^* &= \frac{1}{2} (\alpha - 2\beta + 2) \\
a_2^* &= 1 - \frac{\alpha}{2} \\
\xi &= \frac{1}{2} (-2\lceil \beta \rceil + d + 2) \\
c^* &= \frac{1}{2}
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} -\frac{d}{\alpha} & -\frac{d+2}{\alpha} & -\frac{d+4}{\alpha} & -\frac{d+6}{\alpha} & -\frac{d+8}{\alpha} & -\frac{d+10}{\alpha} & -\frac{d+12}{\alpha} & -\frac{d+14}{\alpha} & -\frac{d+16}{\alpha} & -\frac{d+18}{\alpha} & -\frac{d+20}{\alpha} \\ -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 & -11 \end{array} \right)$$

2.10 Example FoxH32-21-Z-Star.wls

File content

```
{
  (* Upper List *) {
    (* Upper Front List *) {{1, 1}},
    (* Upper Rear List *)  {{1, β}}
  },
  (* Lower List *) {
    (* Lower Front List *) {{d/2, α/2}, {1, 1}},
    (* Lower Rear List *)  {{1, α/2}}
  }
}
```

Fox H-function

$$H_{2,3}^{2,1} \left(\cdot \left| \begin{array}{c} (1, 1), (1, \beta) \\ (\frac{d}{2}, \frac{\alpha}{2}), (1, 1), (1, \frac{\alpha}{2}) \end{array} \right. \right)$$

$$H_{2,3}^{2,1} \left(\cdot \left| \frac{(1, 1)}{(\frac{d}{2}, \frac{\alpha}{2}), (1, 1)} \right| \frac{(1, \beta)}{(1, \frac{\alpha}{2})} \right)$$

Summary

$$\begin{aligned} a^* &= 2 - \beta \\ \Delta &= \alpha - \beta \\ \delta &= 2^{-\alpha} \left(2^{\alpha/2} \alpha^{\alpha/2} + \alpha^\alpha \right) \beta^{-\beta} \\ \mu &= \frac{d-1}{2} \\ a_1^* &= \frac{1}{2}(\alpha - 2\beta + 2) \\ a_2^* &= 1 - \frac{\alpha}{2} \\ \xi &= \frac{d}{2} \\ c^* &= \frac{1}{2} \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} -\frac{d}{\alpha} & -\frac{d+2}{\alpha} & -\frac{d+4}{\alpha} & -\frac{d+6}{\alpha} & -\frac{d+8}{\alpha} & -\frac{d+10}{\alpha} & -\frac{d+12}{\alpha} & -\frac{d+14}{\alpha} & -\frac{d+16}{\alpha} & -\frac{d+18}{\alpha} & -\frac{d+20}{\alpha} \\ -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 & -11 \end{pmatrix}$$

2.11 Example FoxH-Bessel-J_2_9_18.wls

File content

```
(* (2.9.18) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{(a+η)/2, 1}},
    (* Lower Rear List *) {{(a-η)/1, 1}}
  }
}
```

Fox H-function

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c} \\ \left(\frac{a+\eta}{2}, 1 \right), (a-\eta, 1) \end{array} \right. \right)$$

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c|c} & \\ \hline \left(\frac{a+\eta}{2}, 1 \right) & (a-\eta, 1) \end{array} \right. \right)$$

Summary

$$\begin{aligned} a^* &= 0 \\ \Delta &= 2 \\ \delta &= \text{ComplexInfinity} \\ \mu &= \frac{1}{2}(3a - \eta - 2) \\ a_1^* &= 1 \\ a_2^* &= -1 \\ \xi &= \frac{1}{2}(3\eta - a) \\ c^* &= 0 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\{\}, \{\}, \{\}, \{\}, \{\}, \{\}, \{\}, \{\}, \{\}, \{\}, \{\}\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} \frac{1}{2}(-a - \eta) \\ \frac{1}{2}(-a - \eta - 2) \\ \frac{1}{2}(-a - \eta - 4) \\ \frac{1}{2}(-a - \eta - 6) \\ \frac{1}{2}(-a - \eta - 8) \\ -\frac{a}{2} - \frac{\eta}{2} - 5 \\ -\frac{a}{2} - \frac{\eta}{2} - 6 \\ -\frac{a}{2} - \frac{\eta}{2} - 7 \\ -\frac{a}{2} - \frac{\eta}{2} - 8 \\ -\frac{a}{2} - \frac{\eta}{2} - 9 \\ \frac{1}{2}(-a - \eta - 20) \end{pmatrix}$$

2.12 Example FoxH-Bessel-K_2_9_19.wls

File content

```
(* (2.9.19) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{(a-η)/2, 1},{(a+η)/2, 1}},
    (* Lower Rear List *) {}
  }
}
```

Fox H-function

$$H_{0,2}^{2,0} \left(\cdot \left| \begin{pmatrix} \frac{a-\eta}{2}, 1 \end{pmatrix}, \begin{pmatrix} \frac{a+\eta}{2}, 1 \end{pmatrix} \right. \right)$$

$$H_{0,2}^{2,0} \left(\cdot \left| \frac{\quad}{\left(\frac{a-\eta}{2}, 1\right), \left(\frac{a+\eta}{2}, 1\right)} \right| \right)$$

Summary

$$\begin{aligned} a^* &= 2 \\ \Delta &= 2 \\ \delta &= \text{Indeterminate} \\ \mu &= a - 1 \\ a_1^* &= 2 \\ a_2^* &= 0 \\ \xi &= a \\ c^* &= 1 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} \frac{\eta-a}{2} & \frac{1}{2}(-a+\eta-2) & \frac{1}{2}(-a+\eta-4) & \frac{1}{2}(-a+\eta-6) & \frac{1}{2}(-a+\eta-8) & \frac{1}{2}(-a+\eta-10) & \frac{1}{2}(-a+\eta-12) \\ \frac{1}{2}(-a-\eta) & \frac{1}{2}(-a-\eta-2) & \frac{1}{2}(-a-\eta-4) & \frac{1}{2}(-a-\eta-6) & \frac{1}{2}(-a-\eta-8) & -\frac{a}{2}-\frac{\eta}{2}-5 & -\frac{a}{2}-\frac{\eta}{2}-7 \end{pmatrix}$$

2.13 Example FoxH-Bessel-Y_2_9_20.wls

File content

```
(* (2.9.20) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {{(a-η-1)/2, 1}}
  },
  (* Lower List *) {
    (* Lower Front List *) {{(a-η)/2, 1},{(a+η)/2, 1}},
    (* Lower Rear List *) {{(a-η-1)/2, 1}}
  }
}
```

Fox H-function

$$H_{1,3}^{2,0} \left(\cdot \left| \begin{array}{c} (\frac{1}{2}(a-\eta-1), 1) \\ (\frac{a-\eta}{2}, 1), (\frac{a+\eta}{2}, 1), (\frac{1}{2}(a-\eta-1), 1) \end{array} \right. \right)$$

$$H_{1,3}^{2,0} \left(\cdot \left| \frac{\phantom{(\frac{1}{2}(a-\eta-1), 1)}}{(\frac{a-\eta}{2}, 1), (\frac{a+\eta}{2}, 1)} \right| \begin{array}{c} (\frac{1}{2}(a-\eta-1), 1) \\ (\frac{1}{2}(a-\eta-1), 1) \end{array} \right)$$

Summary

$$\begin{aligned} a^* &= 0 \\ \Delta &= 2 \\ \delta &= \text{ComplexInfinity} \\ \mu &= a - 1 \\ a_1^* &= 1 \\ a_2^* &= -1 \\ \xi &= \eta + 1 \\ c^* &= 0 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccc} \frac{\eta-a}{2} & \frac{1}{2}(-a+\eta-2) & \frac{1}{2}(-a+\eta-4) & \frac{1}{2}(-a+\eta-6) & \frac{1}{2}(-a+\eta-8) & \frac{1}{2}(-a+\eta-10) & \frac{1}{2}(-a+\eta-12) \\ \frac{1}{2}(-a-\eta) & \frac{1}{2}(-a-\eta-2) & \frac{1}{2}(-a-\eta-4) & \frac{1}{2}(-a-\eta-6) & \frac{1}{2}(-a-\eta-8) & -\frac{a}{2}-\frac{\eta}{2}-5 & -\frac{a}{2}-\frac{\eta}{2}-7 \end{array} \right)$$

2.14 Example FoxH-Cos.wls

File content

```
(* (2.9.8) and (2.9.10) of Kilbas & Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{0, 1}},
    (* Lower Rear List *) {{1/2, 1}}
  }
}
```

Fox H-function

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c} \\ (0,1), (\frac{1}{2},1) \end{array} \right. \right)$$

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c|c} & \\ (0,1) & (\frac{1}{2},1) \end{array} \right. \right)$$

Summary

$$\begin{aligned} a^* &= 0 \\ \Delta &= 2 \\ \delta &= \text{ComplexInfinity} \\ \mu &= -\frac{1}{2} \\ a_1^* &= 1 \\ a_2^* &= -1 \\ \xi &= -\frac{1}{2} \\ c^* &= 0 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{array} \right)$$

2.15 Example FoxH-H.G_2_9_14.wls

File content

```
(* (2.9.14) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{1-a,1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{0,1}},
    (* Lower Rear List *) {{1-c,1}}
  }
}
```

Fox H-function

$$H_{1,2}^{1,1} \left(\cdot \left| \begin{array}{c} (1-a, 1) \\ (0, 1), (1-c, 1) \end{array} \right. \right)$$

$$H_{1,2}^{1,1} \left(\cdot \left| \begin{array}{c|c} (1-a, 1) & \\ \hline (0, 1) & (1-c, 1) \end{array} \right. \right)$$

Summary

$$\begin{aligned} a^* &= 1 \\ \Delta &= 1 \\ \delta &= \text{ComplexInfinity} \\ \mu &= a - c - \frac{1}{2} \\ a_1^* &= 1 \\ a_2^* &= 0 \\ \xi &= c - a \\ c^* &= \frac{1}{2} \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} a & a+1 & a+2 & a+3 & a+4 & a+5 & a+6 & a+7 & a+8 & a+9 & a+10 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccccc} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{array} \right)$$

2.16 Example FoxH-H.G_2_9_15.wls

File content

```
(* (2.9.15) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{1-a,1},{1-b,1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{0,1}},
    (* Lower Rear List *) {{1-c,1}}
  }
}
```

Fox H-function

$$H_{2,2}^{1,2} \left(\begin{matrix} \cdot \\ (1-a, 1), (1-b, 1) \\ (0, 1), (1-c, 1) \end{matrix} \right)$$

$$H_{2,2}^{1,2} \left(\begin{matrix} \cdot \\ \frac{(1-a, 1), (1-b, 1)}{(0, 1)} \\ (1-c, 1) \end{matrix} \right)$$

Summary

$$\begin{aligned} a^* &= 2 \\ \Delta &= 0 \\ \delta &= \text{ComplexInfinity} \\ \mu &= a + b - c - 1 \\ a_1^* &= 1 \\ a_2^* &= 1 \\ \xi &= -a - b + c + 1 \\ c^* &= 1 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \begin{pmatrix} a & a+1 & a+2 & a+3 & a+4 & a+5 & a+6 & a+7 & a+8 & a+9 & a+10 \\ b & b+1 & b+2 & b+3 & b+4 & b+5 & b+6 & b+7 & b+8 & b+9 & b+10 \end{pmatrix}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{pmatrix}$$

2.17 Example FoxH-Lommel_2_9_22.wls

File content

```
(* (2.9.22) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{(1+μ)/2,1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
```

```

(* Lower Front List *) {{(1+μ)/2,1},{η/2,1},{-η/2,1}},
(* Lower Rear List *) {}
}
}

```

Fox H-function

$$H_{1,3}^{3,1} \left(\cdot \left| \begin{array}{c} \left(\frac{\mu+1}{2}, 1 \right) \\ \left(\frac{\mu+1}{2}, 1 \right), \left(\frac{\eta}{2}, 1 \right), \left(-\frac{\eta}{2}, 1 \right) \end{array} \right. \right)$$

$$H_{1,3}^{3,1} \left(\cdot \left| \frac{\left(\frac{\mu+1}{2}, 1 \right)}{\left(\frac{\mu+1}{2}, 1 \right), \left(\frac{\eta}{2}, 1 \right), \left(-\frac{\eta}{2}, 1 \right)} \right| \right)$$

Summary

$$\begin{aligned}
a^* &= 4 \\
\Delta &= 2 \\
\delta &= \text{Indeterminate} \\
\mu &= -1 \\
a_1^* &= 3 \\
a_2^* &= 1 \\
\xi &= \mu + 1 \\
c^* &= 2
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\frac{1-\mu}{2} \quad \frac{3-\mu}{2} \quad \frac{5-\mu}{2} \quad \frac{7-\mu}{2} \quad \frac{9-\mu}{2} \quad \frac{11-\mu}{2} \quad \frac{13-\mu}{2} \quad \frac{15-\mu}{2} \quad \frac{17-\mu}{2} \quad \frac{19-\mu}{2} \quad \frac{21-\mu}{2} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(\begin{array}{cccccccccc} \frac{1}{2}(-\mu-1) & \frac{1}{2}(-\mu-3) & \frac{1}{2}(-\mu-5) & \frac{1}{2}(-\mu-7) & \frac{1}{2}(-\mu-9) & \frac{1}{2}(-\mu-11) & \frac{1}{2}(-\mu-13) & \frac{1}{2}(-\mu-15) & \frac{1}{2}(-\mu-17) & \frac{1}{2}(-\mu-19) \\ -\frac{\eta}{2} & -\frac{\eta}{2}-1 & -\frac{\eta}{2}-2 & -\frac{\eta}{2}-3 & -\frac{\eta}{2}-4 & -\frac{\eta}{2}-5 & -\frac{\eta}{2}-6 & -\frac{\eta}{2}-7 & -\frac{\eta}{2}-8 & -\frac{\eta}{2}-9 \\ \frac{\eta}{2} & \frac{\eta-2}{2} & \frac{\eta-4}{2} & \frac{\eta-6}{2} & \frac{\eta-8}{2} & \frac{\eta}{2}-5 & \frac{\eta}{2}-6 & \frac{\eta}{2}-7 & \frac{\eta}{2}-8 & \frac{\eta}{2}-9 \end{array} \right)$$

2.18 Example FoxH-Mittag-Leffler.wls

File content

```
(* (2.9.27) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {{0, 1}},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{0, 1}},
    (* Lower Rear List *) {{1 - μ, ρ}}
  }
}
```

Fox H-function

$$H_{1,2}^{1,1} \left(\cdot \left| \begin{array}{c} (0, 1) \\ (0, 1), (1 - \mu, \rho) \end{array} \right. \right)$$

$$H_{1,2}^{1,1} \left(\cdot \left| \frac{(0, 1)}{(0, 1) \mid (1 - \mu, \rho)} \right. \right)$$

Summary

$$\begin{aligned} a^* &= 2 - \rho \\ \Delta &= \rho \\ \delta &= \text{ComplexInfinity} \\ \mu &= \frac{1}{2} - \mu \\ a_1^* &= 1 \\ a_2^* &= 1 - \rho \\ \xi &= \mu - 1 \\ c^* &= \frac{1}{2} \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \left(\begin{array}{cccccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \end{array} \right)$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} 0 & -1 & -2 & -3 & -4 & -5 & -6 & -7 & -8 & -9 & -10 \end{pmatrix}$$

2.19 Example FoxH-Sin.wls

File content

```
(* (2.9.7) and (2.9.9) of Kilbas & Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {}
  },
  (* Lower List *) {
    (* Lower Front List *) {{1/2, 1}},
    (* Lower Rear List *) {{0, 1}}
  }
}
```

Fox H-function

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c} \\ (\frac{1}{2}, 1), (0, 1) \end{array} \right. \right)$$

$$H_{0,2}^{1,0} \left(\cdot \left| \begin{array}{c|c} & \\ \hline (\frac{1}{2}, 1) & (0, 1) \end{array} \right. \right)$$

Summary

$$\begin{aligned} a^* &= 0 \\ \Delta &= 2 \\ \delta &= \text{ComplexInfinity} \\ \mu &= -\frac{1}{2} \\ a_1^* &= 1 \\ a_2^* &= -1 \\ \xi &= \frac{1}{2} \\ c^* &= 0 \end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \left(-\frac{1}{2} \quad -\frac{3}{2} \quad -\frac{5}{2} \quad -\frac{7}{2} \quad -\frac{9}{2} \quad -\frac{11}{2} \quad -\frac{13}{2} \quad -\frac{15}{2} \quad -\frac{17}{2} \quad -\frac{19}{2} \quad -\frac{21}{2} \right)$$

2.20 Example FoxH-Whittaker_2_9_21.wls

File content

```
(* (2.9.21) of Kilbas and Saigo 04 *)
{
  (* Upper List *) {
    (* Upper Front List *) {},
    (* Upper Rear List *) {{a-λ+1,1}}
  },
  (* Lower List *) {
    (* Lower Front List *) {{a+μ+1/2,1},{a-μ+1/2,1}},
    (* Lower Rear List *) {}
  }
}
```

Fox H-function

$$H_{1,2}^{2,0} \left(. \left| \begin{array}{c} (a - \lambda + 1, 1) \\ (a + \mu + \frac{1}{2}, 1), (a - \mu + \frac{1}{2}, 1) \end{array} \right. \right)$$

$$H_{1,2}^{2,0} \left(. \left| \frac{}{(a + \mu + \frac{1}{2}, 1), (a - \mu + \frac{1}{2}, 1)} \right| (a - \lambda + 1, 1) \right)$$

Summary

$$\begin{aligned}
a^* &= 1 \\
\Delta &= 1 \\
\delta &= \text{Indeterminate} \\
\mu &= a + \lambda - \frac{1}{2} \\
a_1^* &= 1 \\
a_2^* &= 0 \\
\xi &= a + \lambda \\
c^* &= \frac{1}{2}
\end{aligned}$$

Poles 1. First ten poles from upper front list

$$a_{i,k} = \{\}$$

2. First ten poles from lower front list

$$b_{j,\ell} = \begin{pmatrix} -a - \mu - \frac{1}{2} & -a - \mu - \frac{3}{2} & -a - \mu - \frac{5}{2} & -a - \mu - \frac{7}{2} & -a - \mu - \frac{9}{2} & -a - \mu - \frac{11}{2} & -a - \mu - \frac{13}{2} & -a - \mu - \frac{15}{2} \\ -a + \mu - \frac{1}{2} & -a + \mu - \frac{3}{2} & -a + \mu - \frac{5}{2} & -a + \mu - \frac{7}{2} & -a + \mu - \frac{9}{2} & -a + \mu - \frac{11}{2} & -a + \mu - \frac{13}{2} & -a + \mu - \frac{15}{2} \end{pmatrix}$$

References

- [Fox61] Charles Fox. The G and H functions as symmetrical Fourier kernels. *Trans. Amer. Math. Soc.*, 98:395–429, 1961.
- [KS] Anatoly A. Kilbas and Megumi Saigo. H -transforms, volume 9 of *Analytical Methods and Special Functions*. Chapman & Hall/CRC, Boca Raton, FL. Theory and applications.