

Probing Strangeness Canonical Ensemble with K^- , $\phi(1020)$ and Ξ^- Production in Au+Au Collisions at $\sqrt{s_{\text{NN}}} = 3 \text{ GeV}$

M. S. Abdallah,⁵ B. E. Aboona,⁵⁵ J. Adam,⁶ L. Adamczyk,² J. R. Adams,³⁹ J. K. Adkins,³⁰ G. Agakishiev,²⁸ I. Aggarwal,⁴¹ M. M. Aggarwal,⁴¹ Z. Ahammed,⁶⁰ I. Alekseev,^{3,35} D. M. Anderson,⁵⁵ A. Aparin,²⁸ E. C. Aschenauer,⁶ M. U. Ashraf,¹¹ F. G. Atetalla,²⁹ A. Attri,⁴¹ G. S. Averichev,²⁸ V. Bairathi,⁵³ W. Baker,¹⁰ J. G. Ball Cap,²⁰ K. Barish,¹⁰ A. Behera,⁵² R. Bellwied,²⁰ P. Bhagat,²⁷ A. Bhasin,²⁷ J. Bielcik,¹⁴ J. Bielcikova,³⁸ I. G. Bordyuzhin,³ J. D. Brandenburg,⁶ A. V. Brandin,³⁵ I. Bunzarov,²⁸ J. Butterworth,⁴⁵ X. Z. Cai,⁵⁰ H. Caines,⁶³ M. Calderón de la Barca Sánchez,⁸ D. Cebra,⁸ I. Chakaberia,^{31,6} P. Chaloupka,¹⁴ B. K. Chan,⁹ F.-H. Chang,³⁷ Z. Chang,⁶ N. Chankova-Bunzarova,²⁸ A. Chatterjee,¹¹ S. Chattopadhyay,⁶⁰ D. Chen,¹⁰ J. Chen,⁴⁹ J. H. Chen,¹⁸ X. Chen,⁴⁸ Z. Chen,⁴⁹ J. Cheng,⁵⁷ M. Chevalier,¹⁰ S. Choudhury,¹⁸ W. Christie,⁶ X. Chu,⁶ H. J. Crawford,⁷ M. Csanád,¹⁶ M. Daugherty,¹ T. G. Dedovich,²⁸ I. M. Deppner,¹⁹ A. A. Derevschikov,⁴³ A. Dhamija,⁴¹ L. Di Carlo,⁶² L. Didenko,⁶ P. Dixit,²² X. Dong,³¹ J. L. Drachenberg,¹ E. Duckworth,²⁹ J. C. Dunlop,⁶ N. Elsey,⁶² J. Engelage,⁷ G. Eppley,⁴⁵ S. Esumi,⁵⁸ O. Evdokimov,¹² A. Ewigleben,³² O. Eyser,⁶ R. Fatemi,³⁰ F. M. Fawzi,⁵ S. Fazio,⁶ P. Federic,³⁸ J. Fedorisin,²⁸ C. J. Feng,³⁷ Y. Feng,⁴⁴ P. Filip,²⁸ E. Finch,⁵¹ Y. Fisyrak,⁶ A. Francisco,⁶³ C. Fu,¹¹ L. Fulek,² C. A. Gagliardi,⁵⁵ T. Galatyuk,¹⁵ F. Geurts,⁴⁵ N. Ghimire,⁵⁴ A. Gibson,⁵⁹ K. Gopal,²³ X. Gou,⁴⁹ D. Grosnick,⁵⁹ A. Gupta,²⁷ W. Guryn,⁶ A. I. Hamad,²⁹ A. Hamed,⁵ Y. Han,⁴⁵ S. Harabasz,¹⁵ M. D. Harasty,⁸ J. W. Harris,⁶³ H. Harrison,³⁰ S. He,¹¹ W. He,¹⁸ X. H. He,²⁶ Y. He,⁴⁹ S. Heppelmann,⁸ S. Heppelmann,⁴² N. Herrmann,¹⁹ E. Hoffman,²⁰ L. Holub,¹⁴ Y. Hu,¹⁸ H. Huang,³⁷ H. Z. Huang,⁹ S. L. Huang,⁵² T. Huang,³⁷ X. Huang,⁵⁷ Y. Huang,⁵⁷ T. J. Humanic,³⁹ G. Igo,^{9,*} D. Isenhowe,¹ W. W. Jacobs,²⁵ C. Jena,²³ A. Jentsch,⁶ Y. Ji,³¹ J. Jia,^{6,52} K. Jiang,⁴⁸ X. Ju,⁴⁸ E. G. Judd,⁷ S. Kabana,⁵³ M. L. Kabir,¹⁰ S. Kagamaster,³² D. Kalinkin,^{25,6} K. Kang,⁵⁷ D. Kapukchyan,¹⁰ K. Kauder,⁶ H. W. Ke,⁶ D. Keane,²⁹ A. Kechechyan,²⁸ M. Kelsey,⁶² Y. V. Khyzhniak,³⁵ D. P. Kikoła,⁶¹ C. Kim,¹⁰ B. Kimelman,⁸ D. Kincses,¹⁶ I. Kisel,¹⁷ A. Kiselev,⁶ A. G. Knospe,³² H. S. Ko,³¹ L. Kochenda,³⁵ L. K. Kosarzewski,¹⁴ L. Kramarik,¹⁴ P. Kravtsov,³⁵ L. Kumar,⁴¹ S. Kumar,²⁶ R. Kunnawalkam Elayavalli,⁶³ J. H. Kwasizur,²⁵ R. Lacey,⁵² S. Lan,¹¹ J. M. Landgraf,⁶ J. Lauret,⁶ A. Lebedev,⁶ R. Lednicky,^{28,38} J. H. Lee,⁶ Y. H. Leung,³¹ C. Li,⁴⁹ C. Li,⁴⁸ W. Li,⁴⁵ X. Li,⁴⁸ Y. Li,⁵⁷ X. Liang,¹⁰ Y. Liang,²⁹ R. Licenik,³⁸ T. Lin,⁴⁹ Y. Lin,¹¹ M. A. Lisa,³⁹ F. Liu,¹¹ H. Liu,²⁵ H. Liu,¹¹ P. Liu,⁵² T. Liu,⁶³ X. Liu,³⁹ Y. Liu,⁵⁵ Z. Liu,⁴⁸ T. Ljubicic,⁶ W. J. Llope,⁶² R. S. Longacre,⁶ E. Loyd,¹⁰ N. S. Lukow,⁵⁴ X. F. Luo,¹¹ L. Ma,¹⁸ R. Ma,⁶ Y. G. Ma,¹⁸ N. Magdy,¹² D. Mallick,³⁶ S. Margetis,²⁹ C. Markert,⁵⁶ H. S. Matis,³¹ J. A. Mazer,⁴⁶ N. G. Minaev,⁴³ S. Mioduszewski,⁵⁵ B. Mohanty,³⁶ M. M. Mondal,⁵² I. Mooney,⁶² D. A. Morozov,⁴³ A. Mukherjee,¹⁶ M. Nagy,¹⁶ J. D. Nam,⁵⁴ Md. Nasim,²² K. Nayak,¹¹ D. Neff,⁹ J. M. Nelson,⁷ D. B. Nemes,⁶³ M. Nie,⁴⁹ G. Nigmatkulov,³⁵ T. Niida,⁵⁸ R. Nishitani,⁵⁸ L. V. Nogach,⁴³ T. Nonaka,⁵⁸ A. S. Nunes,⁶ G. Odyniec,³¹ A. Ogawa,⁶ S. Oh,³¹ V. A. Okorokov,³⁵ B. S. Page,⁶ R. Pak,⁶ J. Pan,⁵⁵ A. Pandav,³⁶ A. K. Pandey,⁵⁸ Y. Panebratsev,²⁸ P. Parfenov,³⁵ B. Pawlik,⁴⁰ D. Pawlowska,⁶¹ H. Pei,¹¹ C. Perkins,⁷ L. Pinsky,²⁰ R. L. Pintér,¹⁶ J. Pluta,⁶¹ B. R. Pokhrel,⁵⁴ G. Pomatkin,³⁸ J. Porter,³¹ M. Posik,⁵⁴ V. Prozorova,¹⁴ N. K. Pruthi,⁴¹ M. Przybycien,² J. Putschke,⁶² H. Qiu,²⁶ A. Quintero,⁵⁴ C. Racz,¹⁰ S. K. Radhakrishnan,²⁹ N. Raha,⁶² R. L. Ray,⁵⁶ R. Reed,³² H. G. Ritter,³¹ M. Robotkova,³⁸ O. V. Rogachevskiy,²⁸ J. L. Romero,⁸ D. Roy,⁴⁶ L. Ruan,⁶ J. Rusnak,³⁸ N. R. Sahoo,⁴⁹ H. Sako,⁵⁸ S. Salur,⁴⁶ J. Sandweiss,^{63,*} S. Sato,⁵⁸ W. B. Schmidke,⁶ N. Schmitz,³³ B. R. Schweid,⁵² F. Seck,¹⁵ J. Seger,¹³ M. Sergeeva,⁹ R. Seto,¹⁰ P. Seyboth,³³ N. Shah,²⁴ E. Shahaliev,²⁸ P. V. Shanmuganathan,⁶ M. Shao,⁴⁸ T. Shao,¹⁸ A. I. Sheikh,²⁹ D. Shen,⁵⁰ S. S. Shi,¹¹ Y. Shi,⁴⁹ Q. Y. Shou,¹⁸ E. P. Sichtermann,³¹ R. Sikora,² M. Simko,³⁸ J. Singh,⁴¹ S. Singha,²⁶ M. J. Skoby,⁴⁴ N. Smirnov,⁶³ Y. Söhnngen,¹⁹ W. Solyst,²⁵ P. Sorensen,⁶ H. M. Spinka,^{4,*} B. Srivastava,⁴⁴ T. D. S. Stanislaus,⁵⁹ M. Stefaniak,⁶¹ D. J. Stewart,⁶³ M. Strikhanov,³⁵ B. Stringfellow,⁴⁴ A. A. P. Suaide,⁴⁷ M. Sumera,³⁸ B. Summa,⁴² X. M. Sun,¹¹ X. Sun,¹² Y. Sun,⁴⁸ Y. Sun,²¹ B. Surrow,⁵⁴ D. N. Svirida,³ Z. W. Sweger,⁸ P. Szymanski,⁶¹ A. H. Tang,⁶ Z. Tang,⁴⁸ A. Taranenko,³⁵ T. Tarnowsky,³⁴ J. H. Thomas,³¹ A. R. Timmins,²⁰ D. Tlusty,¹³ T. Todoroki,⁵⁸ M. Tokarev,²⁸ C. A. Tomkiel,³² S. Trentalange,⁹ R. E. Tribble,⁵⁵ P. Tribedy,⁶ S. K. Tripathy,¹⁶ T. Truhlar,¹⁴ B. A. Trzeciak,¹⁴ O. D. Tsai,⁹ Z. Tu,⁶ T. Ullrich,⁶ D. G. Underwood,^{4,59} I. Upsal,⁴⁵ G. Van Buren,⁶ J. Vanek,³⁸ A. N. Vasiliev,⁴³ I. Vassiliev,¹⁷ V. Verkest,⁶² F. Videbæk,⁶ S. Vokal,²⁸ S. A. Voloshin,⁶² F. Wang,⁴⁴ G. Wang,⁹ J. S. Wang,²¹ P. Wang,⁴⁸ Y. Wang,¹¹ Y. Wang,⁵⁷ Z. Wang,⁴⁹ J. C. Webb,⁶ P. C. Weidenkaff,¹⁹ L. Wen,⁹ G. D. Westfall,³⁴ H. Wieman,³¹ S. W. Wissink,²⁵ J. Wu,²⁶ Y. Wu,¹⁰ B. Xi,⁵⁰ Z. G. Xiao,⁵⁷ G. Xie,³¹ W. Xie,⁴⁴ H. Xu,²¹ N. Xu,³¹ Q. H. Xu,⁴⁹ Y. Xu,⁴⁹ Z. Xu,⁶ Z. Xu,⁹ C. Yang,⁴⁹ Q. Yang,⁴⁹ S. Yang,⁴⁵ Y. Yang,³⁷ Z. Ye,⁴⁵ Z. Ye,¹² L. Yi,⁴⁹ K. Yip,⁶ Y. Yu,⁴⁹ H. Zbroszczyk,⁶¹ W. Zha,⁴⁸ C. Zhang,⁵² D. Zhang,¹¹ J. Zhang,⁴⁹ S. Zhang,¹² S. Zhang,¹⁸ X. P. Zhang,⁵⁷ Y. Zhang,²⁶ Y. Zhang,⁴⁸ Y. Zhang,¹¹ Z. J. Zhang,³⁷ Z. Zhang,⁶ Z. Zhang,¹² J. Zhao,⁴⁴ C. Zhou,¹⁸ Y. Zhou,¹¹ X. Zhu,⁵⁷ M. Zurek,⁴ and M. Zyzak¹⁷

(STAR Collaboration)

- ¹Abilene Christian University, Abilene, Texas 79699
- ²AGH University of Science and Technology, FPACS, Cracow 30-059, Poland
- ³Alikhanov Institute for Theoretical and Experimental Physics NRC "Kurchatov Institute", Moscow 117218, Russia
- ⁴Argonne National Laboratory, Argonne, Illinois 60439
- ⁵American University of Cairo, New Cairo 11835, New Cairo, Egypt
- ⁶Brookhaven National Laboratory, Upton, New York 11973
- ⁷University of California, Berkeley, California 94720
- ⁸University of California, Davis, California 95616
- ⁹University of California, Los Angeles, California 90095
- ¹⁰University of California, Riverside, California 92521
- ¹¹Central China Normal University, Wuhan, Hubei 430079
- ¹²University of Illinois at Chicago, Chicago, Illinois 60607
- ¹³Creighton University, Omaha, Nebraska 68178
- ¹⁴Czech Technical University in Prague, FNSPE, Prague 115 19, Czech Republic
- ¹⁵Technische Universität Darmstadt, Darmstadt 64289, Germany
- ¹⁶ELTE Eötvös Loránd University, Budapest, Hungary H-1117
- ¹⁷Frankfurt Institute for Advanced Studies FIAS, Frankfurt 60438, Germany
- ¹⁸Fudan University, Shanghai, 200433
- ¹⁹University of Heidelberg, Heidelberg 69120, Germany
- ²⁰University of Houston, Houston, Texas 77204
- ²¹Huzhou University, Huzhou, Zhejiang 313000
- ²²Indian Institute of Science Education and Research (IISER), Berhampur 760010, India
- ²³Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India
- ²⁴Indian Institute Technology, Patna, Bihar 801106, India
- ²⁵Indiana University, Bloomington, Indiana 47408
- ²⁶Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou, Gansu 730000
- ²⁷University of Jammu, Jammu 180001, India
- ²⁸Joint Institute for Nuclear Research, Dubna 141 980, Russia
- ²⁹Kent State University, Kent, Ohio 44242
- ³⁰University of Kentucky, Lexington, Kentucky 40506-0055
- ³¹Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ³²Lehigh University, Bethlehem, Pennsylvania 18015
- ³³Max-Planck-Institut für Physik, Munich 80805, Germany
- ³⁴Michigan State University, East Lansing, Michigan 48824
- ³⁵National Research Nuclear University MEPhI, Moscow 115409, Russia
- ³⁶National Institute of Science Education and Research, HBNI, Jatni 752050, India
- ³⁷National Cheng Kung University, Tainan 70101
- ³⁸Nuclear Physics Institute of the CAS, Rez 250 68, Czech Republic
- ³⁹Ohio State University, Columbus, Ohio 43210
- ⁴⁰Institute of Nuclear Physics PAN, Cracow 31-342, Poland
- ⁴¹Panjab University, Chandigarh 160014, India
- ⁴²Pennsylvania State University, University Park, Pennsylvania 16802
- ⁴³NRC "Kurchatov Institute", Institute of High Energy Physics, Protvino 142281, Russia
- ⁴⁴Purdue University, West Lafayette, Indiana 47907
- ⁴⁵Rice University, Houston, Texas 77251
- ⁴⁶Rutgers University, Piscataway, New Jersey 08854
- ⁴⁷Universidade de São Paulo, São Paulo, Brazil 05314-970
- ⁴⁸University of Science and Technology of China, Hefei, Anhui 230026
- ⁴⁹Shandong University, Qingdao, Shandong 266237
- ⁵⁰Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800
- ⁵¹Southern Connecticut State University, New Haven, Connecticut 06515
- ⁵²State University of New York, Stony Brook, New York 11794
- ⁵³Instituto de Alta Investigación, Universidad de Tarapacá, Arica 1000000, Chile
- ⁵⁴Temple University, Philadelphia, Pennsylvania 19122
- ⁵⁵Texas A&M University, College Station, Texas 77843
- ⁵⁶University of Texas, Austin, Texas 78712
- ⁵⁷Tsinghua University, Beijing 100084
- ⁵⁸University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
- ⁵⁹Valparaiso University, Valparaiso, Indiana 46383
- ⁶⁰Variable Energy Cyclotron Centre, Kolkata 700064, India
- ⁶¹Warsaw University of Technology, Warsaw 00-661, Poland
- ⁶²Wayne State University, Detroit, Michigan 48201

⁶³Yale University, New Haven, Connecticut 06520

(Dated: September 12, 2021)

We report the first multi-differential measurements of strange hadrons of K^- , ϕ and Ξ^- yields as well as the ratios of ϕ/K^- and ϕ/Ξ^- in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV with the STAR experiment fixed target configuration at RHIC. The ϕ mesons and Ξ^- hyperons are measured through hadronic decay channels, $\phi \rightarrow K^+K^-$ and $\Xi^- \rightarrow \Lambda\pi^-$. Collision centrality and rapidity dependence of the transverse spectra for these strange hadrons are presented. The 4π yields and ratios are compared to thermal model and hadronic transport model predictions. At the collision energy, thermal model with grand canonical ensemble (GCE) under-predicts the ϕ/K^- ratio while the result of canonical ensemble (CE) calculations reproduce well the ratios of ϕ/K^- , with the correlation length $r_c \sim 2.7$ fm, and ϕ/Ξ^- , $r_c \sim 4.2$ fm, for the 0-10% central collisions. Hadronic transport models including high mass resonance decays could also describe the ratios. While thermal calculations with GCE work well for strangeness production in high energy collisions, the change to CE at 3 GeV implies a rather different medium property at high baryon density.

Relativistic heavy ion physics is aiming at the detailed investigation of phase structures of strongly interacting matter, governed by quantum chromodynamics (QCD), under extreme conditions of high temperature and density [1–3]. Particle production has been studied to investigate properties of the produced QCD matter in heavy-ion collisions. The strange quark mass is comparable to the QCD renormalization scale ($\Lambda_{QCD} \sim 200$ MeV) [4, 5], therefore strange quark dynamics plays an important role in understanding the Equation-of-State (EoS) of QCD matter particularly in the high density region [6–11].

Statistical thermal models have often been used to characterize thermal properties of the produced media [12–18]. In these models, grand canonical ensemble (GCE) and canonical ensemble (CE) statistical descriptions can be applied to conserve electric charge, baryon number, and strangeness number in order to compute the final state particle yields. Both GCE and CE models are able to describe various particle yields including strange particles produced in heavy-ion collisions at RHIC and the LHC at center-of-mass energy ($\sqrt{s_{NN}}$) greater than 7.7 GeV. It has been argued that at lower energies, strangeness number needs to be conserved locally on an event-by-event basis described by the CE, which leads to a reduction in the yields of hadrons with non-zero strangeness number (“Canonical Suppression”) [13, 19, 20], but not for the $\phi(1020)$ meson with zero net strangeness number ($S=0$). The ϕ/K^- ratio is expected to increase with decreasing collision energy in models using the CE treatment for strangeness, opposite to the trend in the GCE treatment. The canonical suppression power for Ξ^- ($S=2$) is even larger than for K^- ($S=1$). The ϕ/K^- and ϕ/Ξ^- ratios offer a unique test to scrutinize thermodynamic properties of strange quarks in the hot and dense QCD environment.

In heavy-ion collisions, the near/sub-threshold production of multi-strange hadrons can be achieved from the multiple collisions of nucleons, produced particles, and short-lived resonances [21]. The particle production in heavy-ion collisions below its free nucleon-nucleon (NN)

threshold ($\sqrt{s_{NN}} \sim 2.89$ GeV for ϕ and ~ 3.25 GeV for Ξ^-) is expected to be sensitive to the stiffness of the nuclear EoS at high density [22], as it is for single-strange hadrons [9, 23]. The near/sub-threshold production further provides the possibility to observe exotic states of QCD matter [24] and signatures of “soft deconfinement” [25].

Previous measurements show that the ϕ/K^- ratio in heavy-ion collisions stays remarkably flat (~ 0.15) at collision energies $\sqrt{s_{NN}} > 5$ GeV [26–28]. Recent measurements of the ϕ/K^- ratio in heavy-ion collisions at collision energies below the ϕ free NN-threshold show a hint of relative enhancement compared to those from high energies [29–32], indicative of the applicability of the CE description for strangeness production at these energies. In this Letter, we report high precision measurement of ϕ/K^- and ϕ/Ξ^- ratios in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV from the STAR experiment.

The dataset used in this analysis was collected under the fixed target (FXT) setup [33] in the 2018 RHIC run. A single beam was provided by RHIC with total energy equal to 3.85 GeV/nucleon and incident on a gold target of thickness 0.25 mm, corresponding to a 1% interaction probability. The target is installed inside the vacuum pipe, 2 cm below the center of the beam axis, and located 200 cm to the west of the center of the STAR detector. The main detectors used are the Time Projection Chamber (TPC) [34], the Time of Flight (TOF) detector [35], and the Beam-Beam Counter (BBC) [36]. The trigger is provided by the signal in the east BBC detector and at least five hits in the TOF detector. Tracking and particle identification (PID) are done using the TPC and TOF. Both the TPC and TOF detectors have full azimuthal coverage within a pseudorapidity range of $0 < \eta < 1.88$ for the TPC and $0 < \eta < 1.5$ for the TOF in FXT mode [34, 35]. Events are selected with the offline reconstructed collision vertex within 1.5 cm of the target center along the beam direction. Approximately 2.6×10^8 minimum bias (MB) triggered events passed the selection criteria and are used in this analysis.

The centrality class is selected using measured charged particle multiplicity within the TPC acceptance. A Monte Carlo Glauber model, used in conjunction with

* Deceased

a negative binomial distribution to model particle production in hadronic collisions, is optimized in order to best match the data and determine the centrality class. Due to the trigger inefficiency in the low multiplicity region (corresponding to the most peripheral collisions), we only report the results from the 0–60% centrality in this paper.

ϕ mesons are reconstructed via the decay channel $\phi \rightarrow K^+K^-$ with a branching ratio (BR) of 49.2%, while the Ξ^- hyperons decay via $\Xi^- \rightarrow \Lambda\pi^- \rightarrow p\pi^-\pi^-$ with a BR of 63.8% [37]. Ξ^- reconstruction is performed using the KFPARTICLE package based on the Kalman Filter method [38, 39]. The charged tracks are reconstructed with the TPC in a 0.5 T uniform magnetic field, and are required to consist of at least 20 TPC hits (out of a maximum of 45) and have a ratio between the number of hit points and the maximum possible number of hit points larger than 0.52 to ensure good tracking and avoid track splitting. The charged tracks are identified via a combination of the ionization energy loss measurement with the TPC and the time-of-flight measurement with the TOF. A minimum p_T cut of 0.2 GeV/c is required in the analysis. Since the K^-/π^- ratio is much smaller than the K^+/π^+ ratio at low energies, to reduce the contamination from π^- tracks, a strict PID criterion for K^- requiring both TPC and TOF is implemented [40, 41].

Figure 1 (a) shows the invariant mass distribution of K^+K^- pairs in the transverse momentum (p_T) region of 0.4–1.6 GeV/c for 0–60% central collisions. The combinatorial background is estimated with the mixed-event (ME) technique in which K^+ and K^- from different events of similar characteristics (centrality, event plane angle) are paired. The mixed-event spectra are normalized to the same-event (SE) distributions in the mass range of 1.04–1.08 GeV/c². After the subtraction of the combinatorial background, the remainder distribution, shown as red solid circles, is fitted with a Breit-Wigner function for the signal plus a linear function which represents the remaining correlated background (< 1%) from a partial reconstruction of strange hadrons. The ϕ meson raw yields are extracted from the Breit-Wigner function fit within the corresponding 3σ mass window. The extracted ϕ signal shape is consistent with its intrinsic properties convoluted with the detector smearing effect due to finite momentum resolution (< 3% for single track). Figure 1 (b) shows the invariant mass distribution of $\Lambda(p\pi^-)\pi^-$ in the p_T region of 0.5–2.0 GeV/c for 0–40% central collisions. The combinatorial background is estimated with the rotating daughter (Rot) method, in which a daughter track of Ξ^- is rotated by a random angle between 150 to 210 degrees in the transverse plane. The rotated spectra are normalized to the same-event distributions in the mass ranges of 1.30–1.31 and 1.34–1.35 GeV/c². After the combinatorial background is subtracted, the $\Lambda\pi^-$ invariant mass distribution is fitted with a Gaussian for the signal plus a linear function for the remaining correlated background. The Ξ^- raw yields are obtained via histogram bin counting from the invari-




FIG. 1. Invariant mass distributions of K^+K^- (a) and $\Lambda\pi^-$ (b) in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. Black open circles represent the same-event unlike-sign distribution. The grey shaded histogram represents the normalized mixed-event (rotating daughters for Ξ^-) unlike-sign distribution that is used to estimate the combinatorial background. The red solid circles depict the ϕ meson (a) and Ξ^- (b) signals obtained by subtracting the combinatorial background from the same-event distribution. Reconstructed ϕ (c) and Ξ^- (d) acceptance, p_T vs. rapidity in the center-of-mass frame (y_{cm}) in the same collisions. The dotted line indicates the target rapidity location. The red curve represents the TPC and TOF acceptance edge.

ant mass distributions with all background subtracted within mass windows of 3σ . The reconstructed ϕ and Ξ^- acceptances (p_T vs. y_{cm}) in the collision center-of-mass frame are shown in Fig. 1 (c) and (d), respectively. The target is located at $y_{cm} = -1.05$, using the convention where the beam travels in the positive direction. The red curve represents the TPC and TOF acceptance edge.

Particle raw yields are calculated in each centrality and p_T bin within each rapidity slice. The raw yields are corrected for the TPC acceptance and tracking efficiency, the particle identification efficiency, and the TOF matching and PID efficiency. The final average reconstruction (including acceptance etc.) efficiency is ~ 0.30 , 0.04, and 0.02 for K^- , ϕ and Ξ^- , respectively. As a cross-check, we conducted the measurement of Ξ^- lifetime from the same data and the result is 164.2 ± 6.6 ps, consistent with the PDG value, 163.9 ± 1.5 ps. The corrected p_T spectra in symmetric rapidity bins $(-0.2, 0)$ vs. $(0, 0.2)$ are also consistent.

The systematic uncertainty of the raw yield extraction




FIG. 2. K^- (a), ϕ meson (b) and Ξ^- (c) invariant yields as a function of $m_T - m_0$ for various rapidity regions in 0–10% central Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. Statistical and systematic uncertainties are added quadratically here for plotting. Solid and dashed black lines depict m_T exponential function fits to the measured data points with scaling factors in each rapidity windows.

is estimated by changing the histogram fitting method to bin counting method or by changing the fitting ranges. The maximum difference between these scenarios and the default one is considered as one standard deviation. The contribution varies by p_T , rapidity, and centrality and the overall contribution is less than 5% for the invariant yield. The systematic uncertainty in the TPC acceptance and efficiency correction ε_{TPC} is estimated by varying the cuts on track selection criteria and topological variables (for Ξ^- only). The contribution to the total yield is 4–5% for K^- , 13–16% for ϕ and 6–10% for Ξ^- . This leads to a 10–13% (12–18%) uncertainty in the measured ϕ/K^- (ϕ/Ξ^-) ratio. The uncertainty of the PID efficiency correction is estimated by varying the PID selection cuts and the contribution is less than 3% to the total yield. For the p_T integrated yield, the uncertainty due to the extrapolation to the full p_T range is estimated by choosing several fitting functions [42], and the maximum difference between these scenarios and the default one (m_T -exponential) is quoted as one standard deviation. This contribution is 5–7% for K^- , 14–17% for ϕ and 13–15% for Ξ^- , respectively.

Figure 2 shows the acceptance \times efficiency corrected K^- , ϕ and Ξ^- invariant yields as a function of $m_T - m_0$ ($m_T = \sqrt{m_0^2 + p_T^2}$) for various rapidity ranges in 0–10% central Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. Dashed and solid lines depict fits to the spectra with the m_T -exponential function in order to extrapolate to the unmeasured p_T ranges. Please see the Supplemental Material




FIG. 3. Rapidity density distributions of K^- (squares), ϕ meson (circles) and Ξ^- (diamonds) p_T -integrated yields dN/dy in 0–10% (a), 10–40% (b) and 40–60% central (c) Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. Solid lines depict Gaussian function fits to the data points.

for the p_T spectra in other rapidity and centrality bins. The p_T integrated rapidity distributions dN/dy are displayed in Fig. 3 for Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV for three different centralities. Solid curves depict Gaussian function fits to the data points with the centroid parameter fixed to zero. They are used to extrapolate to the unmeasured rapidity region for calculating total multiplicities.

The ϕ/K^- and ϕ/Ξ^- ratios are presented in Fig. 4 as a function of collision energy $\sqrt{s_{NN}}$, including the midrapidity data in central Au+Au or Pb+Pb data from the AGS, SPS and RHIC BES at higher energies and 4π acceptance data from SIS at lower energies. The black solid circles show our measurements in the 0–10% centrality bin in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. The measured ϕ , K^- and Ξ^- yields in 4π and the ϕ/K^- , ϕ/Ξ^- ratios in different centrality bins are listed in Tab. I. The ϕ/K^- and ϕ/Ξ^- ratios measured at 3 GeV are slightly higher than, or comparable to, the values at high energies for $\sqrt{s_{NN}} \geq 5$ GeV [26–28, 43, 49–53] despite the collision energy being very close to the ϕ threshold and below the Ξ^- threshold in NN collisions.

Various curves in Fig. 4 represent the predictions of ϕ/K^- and ϕ/Ξ^- ratios from several model calculations in central A+A collisions. Statistical model calculations, based on the Grand Canonical Ensemble and Canonical Ensemble for strangeness with several different choices of strangeness correlation length (r_c), were calculated using the THERMUS package [44] with energy dependent freeze-out parameters (T_{ch} , μ_B) taken from [12]. We noted that the ϕ/K^- and ϕ/Ξ^- ratios from GCE depend on μ_S . The same relation, $\mu_S = \mu_B/4$ from the STAR BES-I data [28], was used in this GCE calculation. With unprecedented precision, our data exclude the GCE calculation, which indicates the event-by-event strangeness conservation is crucial [54] in such collisions. In the canonical approach, the correlation length, r_c , defines a region of the particle production phase space inside which the production of the strangeness is canonically

TABLE I. ϕ , K^- , Ξ^- integrated yields and ϕ/K^- and ϕ/Ξ^- ratios for given centrality classes in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV. The first error given corresponds to the statistical one, the second to the systematic error.

Centrality	ϕ (10^{-3})	K^- (10^{-2})	ϕ/K^-	Ξ^- (10^{-3})	ϕ/Ξ^-
0–10%	$20.1 \pm 1.4 \pm 3.8$	$8.70 \pm 0.02 \pm 0.53$	$0.231 \pm 0.016 \pm 0.042$	$13.9 \pm 0.8 \pm 2.4$	$1.45 \pm 0.13 \pm 0.34$
10–40%	$8.5 \pm 0.4 \pm 1.7$	$3.39 \pm 0.01 \pm 0.20$	$0.249 \pm 0.011 \pm 0.046$	$3.61 \pm 0.32 \pm 0.59$	$2.34 \pm 0.23 \pm 0.65$
40–60%	$2.6 \pm 0.2 \pm 0.5$	$0.79 \pm 0.01 \pm 0.06$	$0.327 \pm 0.029 \pm 0.069$	—	—




FIG. 4. ϕ/K^- (a) and ϕ/Ξ^- (b) ratio as a function of collision energy, $\sqrt{s_{NN}}$. The solid black circles show the measurements presented here in 0–10% centrality bin, while empty markers in black are used for data from various other energies and/or collision systems [26–32, 43]. The vertical grey bands on the data points represent the systematic uncertainties. The grey solid line represents a THERMUS calculation based on the Grand Canonical Ensemble (GCE) while the dotted lines depict calculations based on the Canonical Ensemble (CE) with different values of the strangeness correlation radius (r_c) [12, 44]. The green dashed line, green shaded band and the solid red line show transport model calculations from the public versions UrQMD¹ [45, 46], modified UrQMD² [47] and SMASH [48], respectively.

conserved. Both the ϕ/K^- and ϕ/Ξ^- data from our measurement favor the CE thermodynamics for strangeness with a small strangeness correlation length ($r_c \sim 2.7$ fm for ϕ/K^- and $r_c \sim 4.2$ fm for ϕ/Ξ^-). It is worthwhile to point out that the CE calculations with the same r_c parameter cannot describe our ϕ/K^- and ϕ/Ξ^- data simultaneously (as also observed in lighter systems and at lower collision energy [31]). A global thermal model fit with all the particle yields at 3 GeV will help to precisely determine these thermal parameters in the future.

Previous measurements from smaller collision systems

(Ar+KCl and Al+Al collisions) show comparable or higher ϕ/K^- and/or ϕ/Ξ^- ratios at energies below 3 GeV [29–31, 43]. The measurement in $p+p$ collisions at 2.7 GeV shows a much larger ϕ/K^- ratio (1.04 ± 0.23) [55], while the measured ratio at 17.3 GeV (0.11 ± 0.01) is comparable to that in central Au+Au/Pb+Pb collisions at similar energies. The ϕ/Ξ^- ratio in $p+p$ collisions at 17.3 GeV, 5.09 ± 0.36 , is also significantly larger than that in central Au+Au/Pb+Pb collisions [56–58]. In our measurement at 3 GeV, there is no obvious difference in the ϕ/K^- ratio between the 0–10% and 10–40% central bins, while the result in the most peripheral 40–60% central bin shows a hint of a larger value, as shown in Tab. I. Similarly, the ϕ/Ξ^- ratio in mid-central collisions seems to be larger than that in central collisions. Overall, these observations are qualitatively consistent with the expectation that a smaller canonical volume in the smaller system leads to a higher observed ϕ/K^- and/or ϕ/Ξ^- ratio.

Hadronic transport models are widely used in the high baryon density region to study the properties of the produced dense matter [45–48, 59, 60]. In the modified version of the Ultra-relativistic Quantum Molecular Dynamics (UrQMD) model [47], new decay channels from high mass baryon resonances to ϕ and Ξ^- are deployed. The relevant decay branching fraction was determined by fitting the experimental data from $p+p$ collisions [55]. From the comparison shown in Fig. 4, the modified UrQMD² calculation for central ($b < 5$ fm) Au+Au collisions agrees with the data points at low $\sqrt{s_{NN}}$, including our new measurement for ϕ/K^- . However calculations from the public UrQMD¹ model underestimate our measurements for both ϕ/K^- and ϕ/Ξ^- . The SMASH (Simulating Many Accelerated Strongly-interacting Hadrons) model attempts to incorporate the newest available experimental data from both elementary hadronic cross sections and dilepton invariant mass spectra to constrain the resonance branching ratios [48]. The ϕ/K^- ratio is reasonably reproduced using SMASH in the smaller system and $\sqrt{s_{NN}}$ below 3 GeV, despite the overestimation of each individual (ϕ , K^-) transverse mass spectrum measured, e.g. in Au+Au 0–40% system by HADES [32, 48]. The predicted ϕ/K^- ratio from the same model is about 2.5σ higher than central Au+Au 0–10% collisions at 3 GeV. This indicates that some important in-medium mechanism for strangeness production and propagation may be missing for the large system in SMASH.

In summary, we report the systematic measurements of K^- , $\phi(1020)$ and Ξ^- production yields and the ϕ/K^- ,

ϕ/Ξ^- ratios in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV with the STAR experiment at RHIC. The measured ϕ/K^- ratio is significantly larger than the statistical model prediction based on Grand Canonical Ensemble in the 0–10% central collisions. Both the results of ϕ/K^- ($r_c \sim 2.7$ fm) and ϕ/Ξ^- ($r_c \sim 4.2$ fm) ratios favor the Canonical Ensemble model for strangeness production in such collisions. Transport models, including the resonance decays, could reasonably describe our measured ϕ/K^- ratio at 3 GeV and the increasing trend of ϕ/Ξ^- at lower energies. Note that the measurement of collective flow from the 3 GeV Au+Au collisions demonstrated a new EoS dominated by baryonic interactions [61]. The new results from this paper suggest a significant change in the strangeness production at $\sqrt{s_{NN}} = 3$ GeV compared to higher collision energies, providing new insights towards the understanding of the QCD medium properties at high baryon density [6].

We would like to thank K. Redlich and J. Steinheimer for fruitful discussions. We thank the RHIC Operations Group and RCF at BNL, the NERSC Center at LBNL, and the Open Science Grid consortium for providing re-

sources and support. This work was supported in part by the Office of Nuclear Physics within the U.S. DOE Office of Science, the U.S. National Science Foundation, the Ministry of Education and Science of the Russian Federation, National Natural Science Foundation of China, Chinese Academy of Science, the Ministry of Science and Technology of China and the Chinese Ministry of Education, the Higher Education Sprout Project by Ministry of Education at NCKU, the National Research Foundation of Korea, Czech Science Foundation and Ministry of Education, Youth and Sports of the Czech Republic, Hungarian National Research, Development and Innovation Office, New National Excellency Programme of the Hungarian Ministry of Human Capacities, Department of Atomic Energy and Department of Science and Technology of the Government of India, the National Science Centre of Poland, the Ministry of Science, Education and Sports of the Republic of Croatia, RosAtom of Russia and German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), Helmholtz Association, Ministry of Education, Culture, Sports, Science, and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS).

-
- [1] Y. Akiba *et al.*, (2015), arXiv:1502.02730 [nucl-ex].
 - [2] W. Busza, K. Rajagopal, and W. van der Schee, *Ann. Rev. Nucl. Part. Sci.* **68**, 339 (2018).
 - [3] J. Adams *et al.* (STAR), *Nucl. Phys. A* **757**, 102 (2005).
 - [4] J. Rafelski and B. Muller, *Phys. Rev. Lett.* **48**, 1066 (1982), [Erratum: *Phys. Rev. Lett.* **56**, 2334 (1986)].
 - [5] P. Koch, B. Muller, and J. Rafelski, *Phys. Rept.* **142**, 167 (1986).
 - [6] C. M. Ko, *EPJ Web Conf.* **171**, 03002 (2018).
 - [7] P. Danielewicz, R. Lacey, and W. G. Lynch, *Science* **298**, 1592 (2002).
 - [8] T. Galatyuk, *JPS Conf. Proc.* **32**, 010079 (2020).
 - [9] J. Aichelin and C. M. Ko, *Phys. Rev. Lett.* **55**, 2661 (1985).
 - [10] J. Adamczewski-Musch *et al.*, *Phys. Lett. B* **793**, 457 (2019).
 - [11] W. Cassing, E. Bratkovskaya, and A. Sibirtsev, *Nucl. Phys. A* **691**, 753 (2001).
 - [12] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, *Nature* **561**, 321 (2018).
 - [13] J. Rafelski and M. Danos, *Physics Letters B* **97**, 279 (1980).
 - [14] J. Cleymans and H. Satz, *Z. Phys. C* **57**, 135 (1993).
 - [15] F. Becattini, M. Gazdzicki, and J. Sollfrank, *Eur. Phys. J. C* **5**, 143 (1998).
 - [16] W. Florkowski, W. Broniowski, and M. Michalec, *Acta Phys. Polon. B* **33**, 761 (2002).
 - [17] J. Cleymans, A. Frster, H. Oeschler, K. Redlich, and F. Uhlig, *Phys. Lett. B* **603**, 146 (2004).
 - [18] M. Petráň and J. Rafelski, *Phys. Rev. C* **82**, 011901 (2010).
 - [19] J. Rafelski and J. Letessier, *Journal of Physics G: Nuclear and Particle Physics* **28**, 1819 (2002).
 - [20] K. Redlich and A. Tounsi, *Eur. Phys. J. C* **24**, 589 (2002).
 - [21] G. Zeeb, M. Reiter, and M. Bleicher, *Phys. Lett. B* **586**, 297 (2004).
 - [22] G.-C. Yong, Z.-G. Xiao, Y. Gao, and Z.-W. Lin, (2021), arXiv:2105.10284 [nucl-th].
 - [23] C. Fuchs, *Prog. Part. Nucl. Phys.* **56**, 1 (2006).
 - [24] L. McLerran and R. D. Pisarski, *Nucl. Phys. A* **796**, 83 (2007).
 - [25] K. Fukushima, T. Kojo, and W. Weise, *Phys. Rev. D* **102**, 096017 (2020).
 - [26] B. B. Back *et al.* (E917 Collaboration), *Phys. Rev. C* **69**, 054901 (2004).
 - [27] C. Alt *et al.* (NA49 Collaboration), *Phys. Rev. C* **78**, 044907 (2008).
 - [28] J. Adam *et al.* (STAR Collaboration), *Phys. Rev. C* **102**, 034909 (2020).
 - [29] P. Gasik *et al.*, *Eur. Phys. J. A* **52**, 177 (2016).
 - [30] K. Piasecki *et al.* (FOPI Collaboration), *Phys. Rev. C* **91**, 054904 (2015).
 - [31] G. Agakishiev *et al.* (HADES Collaboration), *Phys. Rev. C* **80**, 025209 (2009).
 - [32] J. Adamczewski-Musch *et al.* (HADES Collaboration), *Phys. Lett. B* **778**, 403 (2018).
 - [33] K. C. Meehan, *J. Phys. Conf. Ser.* **742**, 012022 (2016).
 - [34] M. Anderson *et al.*, *Nucl. Instrum. Meth. A* **499**, 659 (2003).
 - [35] W. J. Llope (for STAR), *Nucl. Instrum. Meth. A* **661**, S110 (2012).
 - [36] C. A. Whitten (STAR), *AIP Conf. Proc.* **980**, 390 (2008).
 - [37] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
 - [38] I. Kisel (CBM), *J. Phys. Conf. Ser.* **1070**, 012015 (2018).
 - [39] J. Adam *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **126**, 162301 (2021).
 - [40] Y. Xu *et al.*, *Nucl. Instrum. Meth. A* **614**, 28 (2010).

- [41] M. Shao *et al.*, Nucl. Instrum. Meth. **A558**, 419 (2006).
 [42] B. I. Abelev *et al.* (STAR Collaboration), Phys. Rev. C **79**, 034909 (2009).
 [43] G. Agakishiev *et al.* (HADES Collaboration), Phys. Rev. Lett. **103**, 132301 (2009).
 [44] S. Wheaton, J. Cleymans, and M. Hauer, Comput. Phys. Commun. **180**, 84 (2009).
 [45] M. Bleicher *et al.*, J. Phys. G **25**, 1859 (1999).
 [46] S. Bass *et al.*, Prog. Part. Nucl. Phys. **41**, 255 (1998).
 [47] J. Steinheimer and M. Bleicher, J. Phys. G: Nucl. Part. Phys. **43**, 015104 (2015).
 [48] V. Steinberg *et al.*, Phys. Rev. C **99**, 064908 (2019).
 [49] C. Alt *et al.* (NA49 Collaboration), Phys. Rev. C **77**, 024903 (2008).
 [50] S. V. Afanasiev *et al.* (NA49 Collaboration), Phys. Rev. C **66**, 054902 (2002).
 [51] Phys. Rev. C **78**, 034918 (2008).
 [52] B. Abelev *et al.* (ALICE Collaboration), Phys. Rev. C **91**, 024609 (2015).
 [53] B. I. Abelev *et al.* (STAR Collaboration), Phys. Rev. C **79**, 064903 (2009).
 [54] P. Braun-Munzinger, K. Redlich, and J. Stachel, QuarkGluon Plasma 3 , 491599 (2004).
 [55] Y. Maeda *et al.*, Phys. Rev. C **77**, 015204 (2008).
 [56] A. Aduszkiewicz *et al.* (NA61/SHINE), Eur. Phys. J. C **77**, 671 (2017), arXiv:1705.02467 [nucl-ex].
 [57] A. Aduszkiewicz *et al.* (NA61/SHINE), Eur. Phys. J. C **80**, 199 (2020), arXiv:1908.04601 [nucl-ex].
 [58] A. Aduszkiewicz *et al.* (NA61/SHINE), Eur. Phys. J. C **80**, 833 (2020), arXiv:2006.02062 [nucl-ex].
 [59] C. Hartnack, H. Oeschler, Y. Leifels, E. L. Bratkovskaya, and J. Aichelin, Phys. Rept. **510**, 119 (2012).
 [60] T. Song, L. Tolos, J. Wirth, J. Aichelin, and E. Bratkovskaya, Phys. Rev. C **103**, 044901 (2021).
 [61] M. Abdallah *et al.* (STAR Collaboration), (2021), arXiv:2108.00908 [nucl-ex].