

# Algebraic Topology is Inevitable

A friendly introduction to the applications of AT in other  
mathematics

Guanyu Li

Olivetti, March 2023

# I'm not a topologist!

## Topologist's Morning Routine

Cup of Coffee



Pants



Shirt



# What is Algebraic Topology

# What is Algebraic Topology

## Question

Is what we refer to algebraic topology clear by definition?

# What is Algebraic Topology

## Question

Is what we refer to algebraic topology clear by definition?

## Almost certainly wrong definition

Algebraic topology is the study of (maybe generalised) homology groups, cohomology groups, and homotopy groups of spaces, and the connections among these algebraic objects.

# Examples

## Examples

- ① We have the definition of the fundamental groups, it is all the (based) loops up to homotopy equivalences

## Examples

- ① We have the definition of the fundamental groups, it is all the (based) loops up to homotopy equivalences

$$\pi_1(X, x_0) = [(S^1, s_0), (X, x_0)]_*.$$



## Examples

- 1 We have the definition of the fundamental groups, it is all the (based) loops up to homotopy equivalences

$$\pi_1(X, x_0) = [(S^1, s_0), (X, x_0)]_*.$$

Could we generalise it?

## Examples

- 1 We have the definition of the fundamental groups, it is all the (based) loops up to homotopy equivalences

$$\pi_1(X, x_0) = [(S^1, s_0), (X, x_0)]_*.$$

Could we generalise it?

$$\pi_n(X, x_0) = [(S^n, s_0), (X, x_0)]_*.$$

## Examples

- ① We have the definition of the fundamental groups, it is all the (based) loops up to homotopy equivalences

$$\pi_1(X, x_0) = [(S^1, s_0), (X, x_0)]_*.$$

Could we generalise it?

$$\pi_n(X, x_0) = [(S^n, s_0), (X, x_0)]_*.$$

- ② Hurewicz: Suppose  $X$  is a connected topological space. Then the map

$$\pi_1(X) \rightarrow H_1(X), [\gamma] \mapsto (\sigma : \Delta^1 \rightarrow X, t \mapsto \gamma(t))$$

is a surjective group homomorphism, whose kernel is exactly the commutator subgroup  $[\pi_1(X), \pi_1(X)]$ . Hence

$$H_1 = (\pi_1)_{\text{ab}}.$$

## Examples

- ③ By analogy,  $H_n$  is the abelianisation of  $\pi_n$ , and since  $\pi_n$  is abelian for  $n \geq 2$ ,  $H_n = \pi_n$  for  $n \geq 2$ .

## Examples

- ③ By analogy,  $H_n$  is the abelianisation of  $\pi_n$ , and since  $\pi_n$  is abelian for  $n \geq 2$ ,  $H_n = \pi_n$  for  $n \geq 2$ .

NO! Counterexamples will be showing up later.

## Examples

- ③ By analogy,  $H_n$  is the abelianisation of  $\pi_n$ , and since  $\pi_n$  is abelian for  $n \geq 2$ ,  $H_n = \pi_n$  for  $n \geq 2$ .  
NO! Counterexamples will be showing up later.
- ④ In fact,  $\pi_n$ 's are quite powerful. Whitehead: Suppose  $f : X \rightarrow Y$  is a map between “good” topological spaces, if the induced maps

$$f_* : \pi_n(X, x_0) \rightarrow \pi_n(Y, f(x_0))$$

are isomorphisms for all  $x_0 \in X$  and all  $n \geq 0$ , then  $f$  is a homotopy equivalence.

## Examples

- ③ By analogy,  $H_n$  is the abelianisation of  $\pi_n$ , and since  $\pi_n$  is abelian for  $n \geq 2$ ,  $H_n = \pi_n$  for  $n \geq 2$ .  
NO! Counterexamples will be showing up later.
- ④ In fact,  $\pi_n$ 's are quite powerful. Whitehead: Suppose  $f : X \rightarrow Y$  is a map between “good” topological spaces, if the induced maps

$$f_* : \pi_n(X, x_0) \rightarrow \pi_n(Y, f(x_0))$$

are isomorphisms for all  $x_0 \in X$  and all  $n \geq 0$ , then  $f$  is a homotopy equivalence.

We turned topological information into algebraic information!

## Examples

- ⑤ The homotopy groups are powerful, however it is hard to compute. Even the question

$$\pi_i(S^n) = ?$$

is not fully answered. This question is still somewhat one of the central problems in AT.



## Examples

## Examples

What do we mean by the dimension of a space?

## Examples

What do we mean by the dimension of a space?

- ① One might say  $\dim \mathbb{R}^n$  is the number of elements in a basis, which is  $n$ . However this is by linear algebra, not by topology.

## Examples

What do we mean by the dimension of a space?

- 1 One might say  $\dim \mathbb{R}^n$  is the number of elements in a basis, which is  $n$ . However this is by linear algebra, not by topology.
- 2 One might say in geometry, we *define* a space locally looking like  $\mathbb{R}^n$  of dimension  $n$ . Then there is another question arise immediately: How do we know

$$\mathbb{R}^0, \mathbb{R}^1, \dots, \mathbb{R}^n, \dots$$

are different?

## Examples

- ③ This is a simple consequence of AT.

## Examples

- ③ This is a simple consequence of AT.
  - ① Supposedly  $\mathbb{R}^m \cong \mathbb{R}^n$  as topological spaces, then

$$\mathbb{R}^m - \{x\} \cong \mathbb{R}^n - \{x'\}.$$

## Examples

③ This is a simple consequence of AT.

① Supposedly  $\mathbb{R}^m \cong \mathbb{R}^n$  as topological spaces, then

$$\mathbb{R}^m - \{x\} \cong \mathbb{R}^n - \{x'\}.$$

② Both sides deformation retracts to spheres, so we have  $S^{m-1} \cong S^{n-1}$ .

## Examples

③ This is a simple consequence of AT.

① Supposedly  $\mathbb{R}^m \cong \mathbb{R}^n$  as topological spaces, then

$$\mathbb{R}^m - \{x\} \cong \mathbb{R}^n - \{x'\}.$$

② Both sides deformation retracts to spheres, so we have  $S^{m-1} \cong S^{n-1}$ .

③ By homology comparison,  $m = n$ .



## Examples

Imagine that one is tying one's shoes, using such a way.



In reality one knows that this works - it is different from being untied.

## Examples

How can we know that mathematically it is different from doing nothing?

## Examples


How can we know that mathematically it is different from doing nothing?



We need to know that this is different from a circle.


## Examples

After doing a little thing, we get this beautiful picture:



## Examples


After doing a little thing, we get this beautiful picture:



However

## Examples

After doing a little thing, we get this beautiful picture:



However if you have some knowledge of topology: No matter how we tangle the (closed) shoelace, it is still a circle itself!

## Examples

The difference is coming from how it is embedded into the ambient space, in this case,  $\mathbb{R}^3$ . Not only on what the spaces are.

## Examples

The difference is coming from how it is embedded into the ambient space, in this case,  $\mathbb{R}^3$ . Not only on what the spaces are.

## Definition

- A knot is an embedding  $S^1 \hookrightarrow \mathbb{R}^3$ .




## Examples

The difference is coming from how it is embedded into the ambient space, in this case,  $\mathbb{R}^3$ . Not only on what the spaces are.

## Definition

- A knot is an embedding  $S^1 \hookrightarrow \mathbb{R}^3$ .



- We call such a knot trefoil and we should have that trefoil is different from an untwisted circle.

## Examples

However (I can hear a lot of complaints now)

## Examples

However (I can hear a lot of complaints now) memorizing a map is hard! We want to see more concrete things!

## Examples

However (I can hear a lot of complaints now) memorizing a map is hard! We want to see more concrete things!

Instead of remembering the map, we consider the **complement** of the embedded knot in  $\mathbb{R}^3$ .

## Examples

However (I can hear a lot of complaints now) memorizing a map is hard! We want to see more concrete things!

Instead of remembering the map, we consider the **complement** of the embedded knot in  $\mathbb{R}^3$ . If we can know  $\mathbb{R}^3 - S^1$  is different from  $\mathbb{R}^3 - K$  ( $K$  is the trefoil), then we are done.

## Examples

However (I can hear a lot of complaints now) memorizing a map is hard! We want to see more concrete things!

Instead of remembering the map, we consider the **complement** of the embedded knot in  $\mathbb{R}^3$ . If we can know  $\mathbb{R}^3 - S^1$  is different from  $\mathbb{R}^3 - K$  ( $K$  is the trefoil), then we are done.

But how?

## Examples

However (I can hear a lot of complaints now) memorizing a map is hard! We want to see more concrete things!

Instead of remembering the map, we consider the **complement** of the embedded knot in  $\mathbb{R}^3$ . If we can know  $\mathbb{R}^3 - S^1$  is different from  $\mathbb{R}^3 - K$  ( $K$  is the trefoil), then we are done.

But how? We have algebraic topology!

## Examples

①  $\pi_1(\mathbb{R}^3 - S^1) = \mathbb{Z},$



## Examples

①  $\pi_1(\mathbb{R}^3 - S^1) = \mathbb{Z},$

②

$$\begin{aligned}\pi_1(\mathbb{R}^3 - K) &= \langle x_1, x_2, x_3 \mid x_1 x_2 x_1^{-1} x_3^{-1}, x_2 x_3 x_2^{-1} x_1^{-1}, x_3 x_1 x_3^{-1} x_2^{-1} \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 x_1^{-1} x_2^{-1} = x_1, x_1 x_2 x_1^{-1} x_1 (x_1 x_2 x_1^{-1})^{-1} = x_2 \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 = x_1 x_2 x_1 \rangle \\ &\cong \langle x, y \mid x^3 = y^2 \rangle\end{aligned}$$

where the last isomorphism is by the substitution  
 $y = x_2 x_1, x = x_1 y$  so  $x_1 = x y^{-1}$  and  $x_2 = y^2 x^{-1}$ .

## Examples

①  $\pi_1(\mathbb{R}^3 - S^1) = \mathbb{Z},$

②

$$\begin{aligned}\pi_1(\mathbb{R}^3 - K) &= \langle x_1, x_2, x_3 \mid x_1 x_2 x_1^{-1} x_3^{-1}, x_2 x_3 x_2^{-1} x_1^{-1}, x_3 x_1 x_3^{-1} x_2^{-1} \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 x_1^{-1} x_2^{-1} = x_1, x_1 x_2 x_1^{-1} x_1 (x_1 x_2 x_1^{-1})^{-1} = x_2 \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 = x_1 x_2 x_1 \rangle \\ &\cong \langle x, y \mid x^3 = y^2 \rangle\end{aligned}$$

where the last isomorphism is by the substitution

$y = x_2 x_1, x = x_1 y$  so  $x_1 = x y^{-1}$  and  $x_2 = y^2 x^{-1}$ .

If these two groups are different, then mathematically we are done!

## Examples

①  $\pi_1(\mathbb{R}^3 - S^1) = \mathbb{Z},$

②

$$\begin{aligned}\pi_1(\mathbb{R}^3 - K) &= \langle x_1, x_2, x_3 \mid x_1 x_2 x_1^{-1} x_3^{-1}, x_2 x_3 x_2^{-1} x_1^{-1}, x_3 x_1 x_3^{-1} x_2^{-1} \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 x_1^{-1} x_2^{-1} = x_1, x_1 x_2 x_1^{-1} x_1 (x_1 x_2 x_1^{-1})^{-1} = x_2 \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 = x_1 x_2 x_1 \rangle \\ &\cong \langle x, y \mid x^3 = y^2 \rangle\end{aligned}$$

where the last isomorphism is by the substitution

$$y = x_2 x_1, x = x_1 y \text{ so } x_1 = x y^{-1} \text{ and } x_2 = y^2 x^{-1}.$$

If these two groups are different, then mathematically we are done!  
But again, why are they different?

## Examples

①  $\pi_1(\mathbb{R}^3 - S^1) = \mathbb{Z},$

②

$$\begin{aligned}\pi_1(\mathbb{R}^3 - K) &= \langle x_1, x_2, x_3 \mid x_1 x_2 x_1^{-1} x_3^{-1}, x_2 x_3 x_2^{-1} x_1^{-1}, x_3 x_1 x_3^{-1} x_2^{-1} \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 x_1^{-1} x_2^{-1} = x_1, x_1 x_2 x_1^{-1} x_1 (x_1 x_2 x_1^{-1})^{-1} = x_2 \rangle \\ &= \langle x_1, x_2 \mid x_2 x_1 x_2 = x_1 x_2 x_1 \rangle \\ &\cong \langle x, y \mid x^3 = y^2 \rangle\end{aligned}$$

where the last isomorphism is by the substitution

$y = x_2 x_1, x = x_1 y$  so  $x_1 = xy^{-1}$  and  $x_2 = y^2 x^{-1}$ .

If these two groups are different, then mathematically we are done!

But again, why are they different?

Is it possible to see  $\langle x, y \mid x^3 = y^2 \rangle$  is nonabelian, for instance  $xy \neq yx$ ?

# Applications in Combinatorics

## Examples

- A *finite presentation* of a group  $G$  is a quotient

$$G \cong F/N,$$

where  $F = \langle x_1, \dots, x_m \rangle$  is the free group generated by  $x_1, \dots, x_m$ , and  $N$  is the smallest normal subgroup generated by elements  $R_1, \dots, R_n$  which are called relations. Clearly an element  $w \in F$  is in the normal subgroup  $N$  if and only if it can be written in the form

$$w = \prod_{i=1}^N T_i R_{j_i}^{e_i} T_i^{-1} \quad (1)$$

where  $T_i \in F$  for all  $i$  and  $e_i = \pm 1$ .

## Examples

- The *Word Problem* is to determine when two elements of  $F$  (or two 'words') represent the same element of  $G$ .  
Equivalently, given an element  $w \in F$ , when does it lie in  $N$  so that it has a form of 1. In general, this is always a hard problem in combinatorics. **Magnus** solved this question for  $G = \langle x_1, \dots, x_m \mid R_1, \dots, R_n \rangle$  when  $n = 1$ . We call such groups *1-relator groups*.

## Examples

- The *Word Problem* is to determine when two elements of  $F$  (or two 'words') represent the same element of  $G$ .  
Equivalently, given an element  $w \in F$ , when does it lie in  $N$  so that it has a form of 1. In general, this is always a hard problem in combinatorics. **Magnus** solved this question for  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$  when  $n = 1$ . We call such groups *1-relator groups*.

## Warning

Word problem is generally very difficult -



## Examples

- The *Word Problem* is to determine when two elements of  $F$  (or two 'words') represent the same element of  $G$ .  
Equivalently, given an element  $w \in F$ , when does it lie in  $N$  so that it has a form of 1. In general, this is always a hard problem in combinatorics. **Magnus** solved this question for  $G = \langle x_1, \dots, x_m \mid R_1, \dots, R_n \rangle$  when  $n = 1$ . We call such groups *1-relator groups*.

## Warning

Word problem is generally very difficult - for instance, it is not easy to see  $xy \neq yx$  in  $\langle x, y \mid x^3 = y^2 \rangle$ .

## Examples

- There is also complementary problem, asking the uniqueness of the expression of relations, i.e., determining all relations of  $R_1, \dots, R_n$  of the form

$$\prod_{i=1}^N T_i R_{t_i}^{e_i} T_i^{-1} = 1, \quad (2)$$

## Examples

- There is also complementary problem, asking the uniqueness of the expression of relations, i.e., determining all relations of  $R_1, \dots, R_n$  of the form

$$\prod_{i=1}^N T_i R_{t_i}^{e_i} T_i^{-1} = 1, \quad (2)$$

such as  $R_2 R_1 R_2^{-1} R_2 R_1^{-1} R_2^{-1} = 1$ .

## Examples

- There is also complementary problem, asking the uniqueness of the expression of relations, i.e., determining all relations of  $R_1, \dots, R_n$  of the form

$$\prod_{i=1}^N T_i R_{t_i}^{e_i} T_i^{-1} = 1, \quad (2)$$

such as  $R_2 R_1 R_2^{-1} R_2 R_1^{-1} R_2^{-1} = 1$ .

Also in the case when  $n = 1$ , the problem is solved.

## Theorem (Magnus)

## Theorem (Simple Identity Theorem, Lyndon)

Let  $F$  be the free group on generators  $x_1, \dots, x_m$  and  $G = F/N$  where  $N$  is the smallest normal subgroup containing the relation  $R$ . Then

$$\prod_{i=1}^N T_i R^{e_i} T_i^{-1} = 1$$

implies that the indices  $1, \dots, N$  can be grouped into pairs  $(i, j)$  such that  $t_i = t_j$ ,  $e_i = -e_j$ , and for certain integers  $c_i$ ,  $T_i \equiv T_j Q^{e_i} \pmod{N}$ .

## Theorem (Magnus)

## Theorem (Simple Identity Theorem, Lyndon)

Let  $F$  be the free group on generators  $x_1, \dots, x_m$  and  $G = F/N$  where  $N$  is the smallest normal subgroup containing the relation  $R$ . Then

$$\prod_{i=1}^N T_i R^{e_i} T_i^{-1} = 1$$

implies that the indices  $1, \dots, N$  can be grouped into pairs  $(i, j)$  such that  $t_i = t_j$ ,  $e_i = -e_j$ , and for certain integers  $c_i$ ,  $T_i \equiv T_j Q^{e_i} \pmod{N}$ .

But what does it mean?

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ ,



## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ , and for each generator  $x_i$ , glue a segment on the point, so we get  $\bigvee_{i=1}^m S^1$ .

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ , and for each generator  $x_i$ , glue a segment on the point, so we get  $\bigvee_{i=1}^m S^1$ .
- 2 For each relation  $R_j$ , glue a disk  $D_j$  along the word given by the relation.

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ , and for each generator  $x_i$ , glue a segment on the point, so we get  $\bigvee_{i=1}^m S^1$ .
- 2 For each relation  $R_j$ , glue a disk  $D_j$  along the word given by the relation.
- 3 For instance, if  $G = \langle a, b | aba^{-1}b^{-1} \rangle$ , we get a torus.

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ , and for each generator  $x_i$ , glue a segment on the point, so we get  $\bigvee_{i=1}^m S^1$ .
- 2 For each relation  $R_j$ , glue a disk  $D_j$  along the word given by the relation.
- 3 For instance, if  $G = \langle a, b | aba^{-1}b^{-1} \rangle$ , we get a torus.
- 4 The point is, by van Kampen, we can *read* that the fundamental group of the Cayley complex is exactly  $G$ .

## Examples

Let's consider the construction which is called Cayley complex (associated to the group  $G = \langle x_1, \dots, x_m | R_1, \dots, R_n \rangle$ ):

- 1 Pick one point  $p$ , and for each generator  $x_i$ , glue a segment on the point, so we get  $\bigvee_{i=1}^m S^1$ .
- 2 For each relation  $R_j$ , glue a disk  $D_j$  along the word given by the relation.
- 3 For instance, if  $G = \langle a, b | aba^{-1}b^{-1} \rangle$ , we get a torus.
- 4 The point is, by van Kampen, we can *read* that the fundamental group of the Cayley complex is exactly  $G$ .

Then we can we say more if we have the simple identity theorem?

## Theorem

Given a group of finite presentation  $G = \langle x_1, \dots, x_m | R \rangle$ , if  $R$  is not a proper power (i.e. if  $R = Q^q$  then  $q = 1$ ), its Cayley complex is *aspherical*, namely the universal covering space is contractible.

This theorem is actually equivalent to the simple identity theorem!

## Theorem

Given a group of finite presentation  $G = \langle x_1, \dots, x_m \mid R \rangle$ , if  $R$  is not a proper power (i.e. if  $R = Q^q$  then  $q = 1$ ), its Cayley complex is *aspherical*, namely the universal covering space is contractible.

This theorem is actually equivalent to the simple identity theorem! We again pause for a moment to see what can we derive from this theorem.

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms.



## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

# Applications in Algebra

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

We probably need some other tools to do this. Define

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

We probably need some other tools to do this. Define

- For any group  $G$ , there is a (good) topological space  $BG$  s.t.
  - ①  $\pi_1(BG) = G$ , and

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

We probably need some other tools to do this. Define

- For any group  $G$ , there is a (good) topological space  $BG$  s.t.
  - ①  $\pi_1(BG) = G$ , and
  - ② the universal covering space is contractible.

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

We probably need some other tools to do this. Define

- For any group  $G$ , there is a (good) topological space  $BG$  s.t.
  - ①  $\pi_1(BG) = G$ , and
  - ② the universal covering space is contractible.

We call such a space **the** classifying space of  $G$ .

## Examples

One of the key problem in algebra is the classification of (discrete) groups, up to isomorphisms. It's a very difficult problem, similar to the problem of the classification of topological spaces.

We probably need some other tools to do this. Define

- For any group  $G$ , there is a (good) topological space  $BG$  s.t.
  - ①  $\pi_1(BG) = G$ , and
  - ② the universal covering space is contractible.

We call such a space **the** classifying space of  $G$ .

- For a (discrete) group  $G$ , its *group homology* is defined as

$$H_n(G; \mathbb{Z}) := H_n(BG; \mathbb{Z}).$$

## Examples

- Then another question pops out: Do we have other choices of classifying space? Do other choices give different group homologies?

## Examples

- Then another question pops out: Do we have other choices of classifying space? Do other choices give different group homologies?
- There is a canonical way of constructing such a space, getting such a sequence

$$G \hookrightarrow EG \rightarrow BG.$$



## Examples

- Then another question pops out: Do we have other choices of classifying space? Do other choices give different group homologies?
- There is a canonical way of constructing such a space, getting such a sequence

$$G \hookrightarrow EG \rightarrow BG.$$

- For any two choices, they are homotopy equivalent, so they give the same group homology.

## Examples

- Then another question pops out: Do we have other choices of classifying space? Do other choices give different group homologies?
- There is a canonical way of constructing such a space, getting such a sequence

$$G \hookrightarrow EG \rightarrow BG.$$

- For any two choices, they are homotopy equivalent, so they give the same group homology.

However,

## Examples

- Then another question pops out: Do we have other choices of classifying space? Do other choices give different group homologies?
- There is a canonical way of constructing such a space, getting such a sequence

$$G \hookrightarrow EG \rightarrow BG.$$

- For any two choices, they are homotopy equivalent, so they give the same group homology.

However, do you feel good?

## Examples

- ① Given a group  $G$ , let

$$F_n := \mathbb{Z}[G] \otimes_{\mathbb{Z}} \mathbb{Z}[G] \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} \mathbb{Z}[G]$$

where there are  $n + 1$  terms in total. Let  $G$  acting on  $F_n$  freely by

$$g \cdot (g_0 \otimes g_1 \otimes \cdots \otimes g_n) := (g \cdot g_0) \otimes g_1 \otimes \cdots \otimes g_n$$

and  $F_n$  is a free  $\mathbb{Z}[G]$ -module with a basis of  $\{1 \otimes g_1 \otimes \cdots \otimes g_n\}_{g_i \in G}$ .

## Examples

- ② One could define

$$d_i^{[n]} : F_n \rightarrow F_{n-1}, \quad 0 \leq i \leq n$$

linearly extended by

$$d_i^{[n]}(1 \otimes g_1 \otimes \cdots \otimes g_n) := \begin{cases} g_1 \cdot (1 \otimes g_2 \otimes \cdots \otimes g_n) \\ (1 \otimes g_1 \otimes \cdots \otimes g_{i-1} \otimes g_i g_{i+1} \otimes \cdots \otimes g_n) \\ 1 \otimes g_1 \otimes \cdots \otimes g_{n-1} \end{cases}$$

where  $\{d_i^{[n]}\}_{0 \leq i \leq n}$  is a  $\mathbb{Z}[G]$ -module homomorphism satisfying  $d_i^{[n]} d_j^{[n]} = d_{j-1}^{[n]} d_i^{[n]}$ , so  $\partial_n := \sum_{i=0}^n (-1)^i d_i^{[n]}$  is the boundary map of the chain complex

$$0 \leftarrow F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} \cdots \xleftarrow{\partial_n} F_n \xleftarrow{\partial_{n+1}} \cdots$$

## Examples

- ③  $\epsilon : F_0 = \mathbb{Z}[G] \rightarrow \mathbb{Z}, \sum_{i=1}^N n_i g_i \mapsto \sum_{i=1}^N n_i$  gives an augmentation

$$0 \leftarrow \mathbb{Z} \xleftarrow{\epsilon} F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} \dots \xleftarrow{\partial_n} F_n \xleftarrow{\partial_{n+1}} \dots$$

## Examples

- ③  $\epsilon : F_0 = \mathbb{Z}[G] \rightarrow \mathbb{Z}, \sum_{i=1}^N n_i g_i \mapsto \sum_{i=1}^N n_i$  gives an augmentation

$$0 \leftarrow \mathbb{Z} \xleftarrow{\epsilon} F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} \cdots \xleftarrow{\partial_n} F_n \xleftarrow{\partial_{n+1}} \cdots$$

- ④ The  $n$ -th group homology  $H_n(G; \mathbb{Z})$  is defined as the  $n$ -th homology group of the chain complex

$$0 \leftarrow F_0 \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_1} F_1 \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_2} \cdots \xleftarrow{\partial_n} F_n \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_{n+1}} \cdots$$

## Examples

- ③  $\epsilon : F_0 = \mathbb{Z}[G] \rightarrow \mathbb{Z}, \sum_{i=1}^N n_i g_i \mapsto \sum_{i=1}^N n_i$  gives an augmentation

$$0 \leftarrow \mathbb{Z} \xleftarrow{\epsilon} F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} \dots \xleftarrow{\partial_n} F_n \xleftarrow{\partial_{n+1}} \dots$$

- ④ The  $n$ -th group homology  $H_n(G; \mathbb{Z})$  is defined as the  $n$ -th homology group of the chain complex

$$0 \leftarrow F_0 \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_1} F_1 \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_2} \dots \xleftarrow{\partial_n} F_n \otimes_{\mathbb{Z}[G]} \mathbb{Z} \xleftarrow{\partial_{n+1}} \dots$$

## Theorem

Two definitions of group homology  $H_*(G; \mathbb{Z})$  agree.



## Examples

Computation time!

## Examples

Computation time!

- $B\mathbb{Z} = S^1$ , and  $H_i(S^1) = \mathbb{Z} \oplus \mathbb{Z}$ , so

$$H_n(\mathbb{Z}; \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } n = 0, 1, \\ 0 & \text{otherwise.} \end{cases}$$

## Examples

Computation time!

- $B\mathbb{Z} = S^1$ , and  $H_i(S^1) = \mathbb{Z} \oplus \mathbb{Z}$ , so

$$H_n(\mathbb{Z}; \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } n = 0, 1, \\ 0 & \text{otherwise.} \end{cases}$$

- How about  $H_n(G; \mathbb{Z})$  when  $G = \langle x, y \mid x^3 = y^2 \rangle$ ? If the group homology could tell us that they are different, how nice it could be!

## Examples

Computation time!

- $B\mathbb{Z} = S^1$ , and  $H_i(S^1) = \mathbb{Z} \oplus \mathbb{Z}$ , so

$$H_n(\mathbb{Z}; \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } n = 0, 1, \\ 0 & \text{otherwise.} \end{cases}$$

- How about  $H_n(G; \mathbb{Z})$  when  $G = \langle x, y \mid x^3 = y^2 \rangle$ ? If the group homology could tell us that they are different, how nice it could be!
- Simple identity theorem says its Cayley complex has the property that the universal covering space is contractible, so **the Cayley complex is already (a choice of)  $BG$ !**

## Examples

Computation time continued!

## Examples

Computation time continued!

- The cellular complex of the Cayley complex is easy to compute:

$$\mathbb{Z} \xleftarrow{0} \mathbb{Z}x \oplus \mathbb{Z}y \xleftarrow{1 \mapsto 3x-2y} \mathbb{Z}.$$

## Examples

Computation time continued!

- The cellular complex of the Cayley complex is easy to compute:

$$\mathbb{Z} \xleftarrow{0} \mathbb{Z}x \oplus \mathbb{Z}y \xleftarrow{1 \mapsto 3x-2y} \mathbb{Z}.$$

So the homology groups are

$$H_0 = \mathbb{Z}, \quad H_1 = \mathbb{Z} \quad H_i = 0 \quad i > 1.$$

## Examples

Computation time continued!

- The cellular complex of the Cayley complex is easy to compute:

$$\mathbb{Z} \xleftarrow{0} \mathbb{Z}x \oplus \mathbb{Z}y \xleftarrow{1 \mapsto 3x-2y} \mathbb{Z}.$$

So the homology groups are

$$H_0 = \mathbb{Z}, \quad H_1 = \mathbb{Z} \quad H_i = 0 \quad i > 1.$$

- Oh no! This is the same as the homology  $H_*(\mathbb{Z}; \mathbb{Z})$ .



## Examples

Computation time continued!

- The cellular complex of the Cayley complex is easy to compute:

$$\mathbb{Z} \xleftarrow{0} \mathbb{Z}x \oplus \mathbb{Z}y \xleftarrow{1 \mapsto 3x-2y} \mathbb{Z}.$$

So the homology groups are

$$H_0 = \mathbb{Z}, \quad H_1 = \mathbb{Z} \quad H_i = 0 \quad i > 1.$$

- Oh no! This is the same as the homology  $H_*(\mathbb{Z}; \mathbb{Z})$ . We cannot differ these two groups via group homology.

## Examples

Computation time continued!

- The cellular complex of the Cayley complex is easy to compute:

$$\mathbb{Z} \xleftarrow{0} \mathbb{Z}x \oplus \mathbb{Z}y \xleftarrow{1 \mapsto 3x - 2y} \mathbb{Z}.$$

So the homology groups are

$$H_0 = \mathbb{Z}, \quad H_1 = \mathbb{Z} \quad H_i = 0 \quad i > 1.$$

- Oh no! This is the same as the homology  $H_*(\mathbb{Z}; \mathbb{Z})$ . We cannot differ these two groups via group homology.

## Remark

The difficulty relies on the presentation of the group / the choice of  $BG$ .

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?
- And,

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?
- And, can algebraic be applied to other mathematics?

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?
- And, can algebraic be applied to other mathematics?

But before answering these questions

# Why is Algebraic Topology Inevitable



# Why is Algebraic Topology Inevitable

## Definition

# Why is Algebraic Topology Inevitable

## Definition

Algebraic topology is more of a way of understanding mathematics. It is a machinery, eating objects we care about, spitting out algebraic information such as numbers, polynomials, matrices, groups, modules etc. to classify the objects.

# Why is Algebraic Topology Inevitable

## Definition

Algebraic topology is more of a way of understanding mathematics. It is a machinery, eating objects we care about, spitting out algebraic information such as numbers, polynomials, matrices, groups, modules etc. to classify the objects.

In the famous textbook written by a (retired) professor here at Cornell, AT is divided into several parts, homologies, cohomologies, homotopies, etc. But algebraic topology is far more influencing than these studies. It gives birth to a lot of subjects like category theory, homological algebra, homotopical algebra, and so on, rooted unexpectedly in almost all mathematics.

# Applications in Geometry

## Examples

In the examples before, we see that different choices of things (the name is resolutions) give us the same homology. The general setting of this phenomenon happens is called *homological algebra*, born by algebraic topology. With AT itself, there are many big names related:

## Examples

In the examples before, we see that different choices of things (the name is resolutions) give us the same homology. The general setting of this phenomenon happens is called *homological algebra*, born by algebraic topology. With AT itself, there are many big names related:

- Chern classes and other characteristic classes.

## Examples

In the examples before, we see that different choices of things (the name is resolutions) give us the same homology. The general setting of this phenomenon happens is called *homological algebra*, born by algebraic topology. With AT itself, there are many big names related:

- Chern classes and other characteristic classes.
- Kodaira vanishing and so Kodaira embedding.

## Examples

In the examples before, we see that different choices of things (the name is resolutions) give us the same homology. The general setting of this phenomenon happens is called *homological algebra*, born by algebraic topology. With AT itself, there are many big names related:

- Chern classes and other characteristic classes.
- Kodaira vanishing and so Kodaira embedding.
- Hodge decomposition.



## Examples

In the examples before, we see that different choices of things (the name is resolutions) give us the same homology. The general setting of this phenomenon happens is called *homological algebra*, born by algebraic topology. With AT itself, there are many big names related:

- Chern classes and other characteristic classes.
- Kodaira vanishing and so Kodaira embedding.
- Hodge decomposition.
- ...

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

# Some answers

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

## Answers

- There is a topological reasons, where van Kampen implies knots embedded in  $\mathbb{R}^3$  has the same fundamental group of knots embedded in  $S^3$ , and the latter one is aspherical and its homology is determined by Alexander duality.

# Some answers

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

## Answers

- There is a topological reasons, where van Kampen implies knots embedded in  $\mathbb{R}^3$  has the same fundamental group of knots embedded in  $S^3$ , and the latter one is aspherical and its homology is determined by Alexander duality.
- Yes of course!

# Some answers

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

## Answers

- There is a topological reasons, where van Kampen implies knots embedded in  $\mathbb{R}^3$  has the same fundamental group of knots embedded in  $S^3$ , and the latter one is aspherical and its homology is determined by Alexander duality.
- Yes of course! But IDK the combinatorics to do it  $> - <$

# Some answers

## Questions

- Why do these two groups have the same group homology?
- Can we distinguish these two group?
- Is there other ways of see the trefoil is different from unknot?

## Answers

- There is a topological reasons, where van Kampen implies knots embedded in  $\mathbb{R}^3$  has the same fundamental group of knots embedded in  $S^3$ , and the latter one is aspherical and its homology is determined by Alexander duality.
- Yes of course! But IDK the combinatorics to do it  $> - <$
- Yes!

# Thank you!