

What's in a Name? Etymology and History of Antibacterials and Antifungals

Chad Beisel,^{1,●} Monica Donnelley,^{1,●} Pegah Capaul,¹ Tanya Savenka,^{2,●} Ashly Nham,^{1,●} Matthew Gosen,² James Lewis II,³ Michael Barbachyn,⁴ Angel N. Desai,² and George R. Thompson III^{2,5,●}

¹Department of Pharmacy, University of California–Davis Medical Center, Sacramento, California, USA, ²Division of Infectious Diseases, Department of Internal Medicine, University of California–Davis Medical Center, Sacramento, California, USA, ³Department of Pharmacy, University of Oregon Health Sciences University, Portland, Oregon, USA, ⁴Barbachyn Consulting, LLC, Kalamazoo, Michigan, USA, and ⁵Department of Medical Microbiology and Immunology, University of California–Davis, Davis, California, USA

The development of antimicrobials represents some of the most storied and transformative scientific milestones of the modern era. This review provides an overview of their historical development and etymologic roots, highlighting their enduring contributions to contemporary medicine and public health.

Keywords. antibacterials; antifungals; antimicrobials; etymology; history.

日期：20260907

報告者：R3 施朝貴



PDF 下載

Historical Foundations: From Antiquity to the Antibiotic Era



Antiquity: Natural Remedies



Antiquity and Natural Biocides

Early attempts at wound antisepsis date back millennia:

- **Honey:** Prized by Sumerians, Egyptians, and Greeks; biocidal effects stem from low moisture, low pH, and H_2O_2 .
- **Moldy Bread:** Ancient Egyptian manuscripts record applying moldy bread to wounds.



1909: Magic Bullet



Paul Ehrlich and "The Magic Bullet"

Dr. Paul Ehrlich formulated the concept of "*magic bullet*" (*Magische Kugel*)—targeting pathogens without injuring host tissues. Culminated in **arsphenamine (Salvarsan)** for syphilis.



1930s-40s: Antibiotic Era



Sulfonamides, Penicillin, and ICU

Discovery of **sulfonamides (1930s)** and **penicillin (1940s)** launched the antibiotic era:

- Enabled reliable treatment of fatal infections.
- Directly enabled surgery, immunosuppression, and **intensive care medicine**.



1941+: The Arms Race



Selman Waksman and The "Arms Race"

Waksman introduced the term "**antibiotic**" in 1941 for naturally produced microbial substances. Initiated the continuous clinical "**arms race**" against resistance.

Principles of Antimicrobial Naming: USAN and WHO INN

Balancing structural heritage, global consistency, and medication safety



Prefix (2 Syllables)

+



Stem / Suffix (Class Target)

=



Approved Generic Name



Traditional vs. Modern Governance

- **Traditional Naming:** Early agents were named organically from discovery sources or chemical structures.
- **Modern Standardization:** Overseen by the **USAN Council** in coordination with WHO **INN Programme**.



Suffix and Prefix Dynamics

- **Class Suffix (Stem):** Preserved across new drugs in the same class to denote shared mechanism.
- **Distinct Prefix:** Proposed by developers, typically **2 syllables** to ensure distinct phonetic separation.



Regulatory Rules & Excluded Letters

Strict criteria enforced by USAN and WHO INN:


- **No outcome promises:** Cannot promise efficacy (e.g., no "Cureall").
- **No medical terminology:** Avoid anatomical or disease references.
- **Excluded Letters (H, K, J, W, Y):** Avoided because they are absent in certain Roman alphabets.
- **Cross-Lingual Audits:** Avoid negative foreign meanings (e.g., "nova-" = "no go" in Spanish).



Scope of this Review: Focuses on currently available **antibacterials and antifungals** (antiviral and antimycobacterial agents are excluded).

SECTION

Antibacterial Agents



The Penicillin Saga: Discovery, Wartime Secrecy & Etymology

From Fleming's paintbrush mold to corn steep liquor and secret codes

Discovery & Paintbrush Mold

- **Alexander Fleming (1928):** Observed staphylococcal inhibition surrounding *Penicillium notatum* (*P. rubens*).
- **Etymology:** Latin *penicillus* = "**Paintbrush**", describing conidiophore morphology under microscope. Suffix **-in** = active chemical constituent.
- **Nobel Prize (1945):** Shared by Fleming, Howard Florey, and Ernst Chain for clinical development.

WWII Production Miracle

- **Corn Steep Liquor & Beer Fermentation:** Peoria, IL byproduct + deep-tank beer fermentation multiplied yield for Allied D-Day troops (1944).

Penicillin G: Sequential Lettering

- **Naming Origin:** In the UK, natural variants were numbered in Roman numerals (Penicillin I, II). In the US, letters were used sequentially (A, B, C... G).
- **Penicillin G (Benzylpenicillin):** 7th US sequential variant (Penicillin II in UK). Acid-labile → IV/IM administration only.

Penicillin V: Cold War Secret Code

- **The Secret:** Synthesized in Austria in 1950s; "V" stands for German "**Vertraulich**" (**Confidential/Secret**), not letter 22!
- **PenVK:** "K" denotes **potassium salt** (*Kalium*) for solubility. Acid-stable → oral administration.

Penicillin Class

Etymology and chemical derivations of major penicillin analogs (Beisel et al., Table 1)

Penicillin Analog	Sub-Class / Group	Etymological & Chemical Origin	Pharmacological & Clinical Significance
Penicillin	Natural Product	Latin <i>penicillus</i> = paintbrush mold. Suffix -in = chemical constituent.	Original natural product class sharing core β -lactam ring nucleus.
Dicloxacillin	Isoxazoly	di- = 2 chlorine atoms ; cloxacillin derived from oxacillin.	Semisynthetic penicillinase-resistant oral agent; blocks staphylococcal cleavage.
Nafcillin	Antistaphylococcal	naf- = naphthalene ring derivative + -cillin .	Semisynthetic parenteral antistaphylococcal agent (MSSA).
Oxacillin	Isoxazoly Prototype	oxa- refers to oxygen atom in the isoxazoly ring .	Semisynthetic penicillinase-resistant prototype.
Flucloxacillin	Isoxazoly	Fluorinated & chlorinated derivative of oxacillin (flu- + cloxacillin).	Semisynthetic isoxazoly penicillin widely used for resistant staphylococci.
		amox- = amino + methoxy /hydroxy-	Hydroxyl modification nearly doubles oral

β-Lactamase Inhibitors

Evolution and etymology of "-bactams" (Beisel et al., Table 2)

Discovery & Structural Evolution

- **Natural Prototype:** Clavulanate isolated from *S. clavuligerus* culture filtrate.
- **Structural Shift:** Evolved from β-lactam sulfones to cyclic boronic acids & non-β-lactam DBOs(Diazabicyclooctane (雙氮雜雙環辛烷) ◦).

Nomenclature & Stem Rules

- **Universal Suffix** -bactam : USAN/WHO contraction for β-lactamase inhibitors.
- **Prefix Origins:** Derived from chemical groups (*Tazo-*), early codes (*Avi-*), or brand names (*Va-*).

Inhibitor Name	Category / Stem	Etymological & Chemical Origin
Clavulanic acid	Natural Product	Isolated from <i>Streptomyces clavuligerus</i> culture filtrate. Clavulanate (salt form) has little standalone antimicrobial activity but significantly reduces MIC when combined with β-lactams.
Sulbactam	Sulfone	Halogenated derivative of penicillanic acid (penicillanic acid sulfone). Sulb- derived from sulfone ; -bactam contraction for β-lactamase inhibitor.
Tazobactam	Sulfone Analogue	Analogue of sulbactam. Tazo- prefix is derived from the triazolymethyl group.

Cephalosporins: Discovery, History & Structural Evolution

From Sardinian sewage outfalls to broad-spectrum β -lactam therapy

The Sardinian Sewage Discovery (1948)

Italian microbiologist **Giuseppe Brotzu** investigated sewage outfalls(污水出口) in Sardinia after noticing locals swimming there rarely developed typhoid fever. Isolated *Acremonium chrysogenum* (formerly *Cephalosporium acremonium*), producing **Cephalosporin C**.

The 7-ACA Nucleus & Cephalothin

Chemical cleavage in 1962 produced **7-aminocephalosporanic acid (7-ACA)**, the core template for all semisynthetic derivatives. **Cephalothin** acquired its name from the added thiophene (-th-) group.

Generational Advancement

- **1st Gen:** Active vs Gram-positive cocci (Cefazolin, Cephalexin).
- **2nd Gen / Cephamecins:** Expanded Gram-negative & anaerobic activity (Cefoxitin, Cefotetan).
- **3rd Gen:** High porin penetration & CNS penetration (Ceftriaxone, Ceftazidime).

4th/5th Gen & Trojan Horse Innovation

- **Cefepime (4th Gen):** Zwitterion; C-7 methoxyimino + C-3 pyrrolidine.
- **Ceftaroline (5th Gen):** Thiazole side chain targeting MRSA PBP2a.
- **Cefiderocol:** Siderophore cephalosporin exploiting active Fe^{3+} transport like a Trojan Horse.

Cephalosporin

Etymology and chemical derivations of major cephalosporin analogs (Beisel et al., Table 3)

Agent Name	Sub-Class / Generation	Etymological & Chemical Origin
Cefazolin	1st Generation	Cef- class prefix + -azol- referring to the thiadiazole ring + -in antimicrobial suffix. Derived from 7-ACA.
Cephalexin	1st Generation	Ceph- class prefix + -alexin from Greek <i>alexo</i> (" to ward off ") and an antiquated immunological term for complement.
Cefadroxil	1st Generation (Oral)	Semisynthetic oral agent; -adroxil suffix refers to a hydroxylated side chain.
Cefuroxime	2nd Generation	-uroxime combines oxy (oxygen atom in β -lactam ring) + -ime indicating an oxime group .
Cefotetan	2nd Gen (Cephamycin)	Cephamycin subset; -tetan derives from its 7-methyl-tetrazol-5-yl side chain.
Cefaclor	2nd Generation	Named after the class prefix + -clor referring to chlorine at C-3 position of cephem nucleus.
	2nd Gen	Derived from cephamycin C (<i>Streptomyces lactamdurans</i>): -ox- = 7-α-methoxyl group + -

Carbapenems & Monobactams

From fragile Thienamycin to DHP-1 stable lifesavers and monocyclic β -lactams (Beisel et al., Table 4)

Carbapenem Discovery & Imipenem Milestone

- **1960s-70s Search:** Penicillin resistance spurred the isolation of *Thienamycin* from *S. cattleya* (**thiena-** = sulfur side chain).
- **1985 Imipenem Milestone:** Fragile thienamycin stabilized into Imipenem, paired with Cilastatin to defeat renal DHP-1 degradation.

Bedside Evolution: DHP-1 Stability & OPAT

- **1990s Meropenem:** Engineered C1-methyl modification to resist DHP-1 metabolism independently without needing cilastatin.
- **2001 Ertapenem:** Extended serum albumin binding, unlocking *once-daily dosing*

Monobactam Concept & Natural Origin

- **1981 Monocyclic Concept:** Sykes et al. defined *Monobactam* for monocyclic 2-azetidinone compounds lacking fused bicyclic rings.
- **Natural Isolation:** Isolated *Sulfazecin* & *Isosulfazecin* from *Pseudomonas*, though natural forms had weak antimicrobial activity.

Aztreonam Synthetic Landmark & Clinical Role

- **Squibb Breakthrough:** Fully synthetic Aztreonam prepared using *Threonine* (**-tre-**) as amino acid starting material.
- **MBL Stability:** Gram-negative specific agent; resists Class B metallo- β -lactamases & minimal penicillin cross-allergy.

Agent
Name

Category / Class

Etymological & Structural Origin

Aminoglycosides

Historical discovery, the Nobel controversy, and the "-mycin" vs "-micin" rule (Beisel et al., Table 5)

The "-mycin" vs. "-micin" Genus Rule

- **Etymology:** Portmanteau of *amino sugars* + *glycosidic bonds*.
- **-mycin** (with 'y'): Genus ***Streptomyces*** (Streptomycin, Tobramycin).
- **-micin** (with 'i'): Genus ***Micromonospora*** (Gentamicin, Netilmicin).

Streptomycin & The 1952 Nobel Controversy

- **First Aminoglycoside (1944):** Streptomycin isolated from *Streptomyces griseus* in Waksman's laboratory.
- **Nobel Controversy:** 1952 Nobel awarded solely to lab director **Selman Waksman**. Graduate student **Albert Schatz**, who actually isolated it, was recognized as co-discoverer only after legal settlement and review.

Agent Name	Category / Class	Etymological & Microbial Origin
Streptomycin	Streptomyces (-mycin)	strepto- refers to twisting/coiled nature of <i>Streptomyces griseus</i> ; -mycin denotes <i>Streptomyces</i> genus.
Neomycin	Streptomyces (-mycin)	neo- from Greek for " new " (new antibiotic discovered from <i>Streptomyces fradiae</i> distinguishing it from earlier agents) + -mycin .
Kanamycin	Streptomyces (-mycin)	kana- originates from isolated organism <i>Streptomyces kanamyceticus</i> + -mycin .

Macrolides

Macrocyclic lactones, R.B. Woodward, and Table 6 etymology (Beisel et al., Table 6)



Molecular Architecture & R.B. Woodward

- **Etymology & Core:** Coined by organic chemist **R.B. Woodward** (1950s); Greek *makros* (large) + *-lide* (lactone). Large 12 to 16-atom lactone ring + sugar molecules.
- **Pikromycin (1950s Prototype):** First recognized macrolide from *Streptomyces venezuelae*; named from Greek *pikros* = **bitter taste**.



Azalide Derivation & Clinical Indications

- **Azalide Derivation:** **aza-** denotes **nitrogen atom replacement** inserted into lactone ring + **-lide** (lactone).
- **Clinical Role:** Essential for respiratory infections, *Helicobacter pylori*, mycobacterial, and other infections.

Agent Name	Category / Stem	Etymological & Structural Origin
Erythromycin	Natural Prototype (-mycin)	Isolated from soil bacterium <i>Streptomyces erythreus</i> (now <i>Saccharopolyspora erythraea</i>) screened from Philippine soil. erythro- derived from Greek <i>erythros</i> = " red " (color of producing bacterium) + -mycin (<i>Streptomyces</i> origin).
Azithromycin	15-Membered Azalide	Semisynthetic derivative of erythromycin A with nitrogen inserted into lactone ring to create a 15-membered azalide. aza- prefix indicates nitrogen atom added + -thromycin suffix indicating similarity to erythromycin.
Semisynthetic derivative of erythromycin A with nitrogen inserted into lactone ring to create a 15-membered azalide.		

Quinolones

From Nalidixic acid to C6-fluorine and C7-piperazine evolution (Beisel et al., Table 7)

1960s Serendipity & C6-Fluorine Milestone

- **Accidental By-product:** *Nalidixic acid* isolated by George Leshner (Sterling Drugs) during chloroquine synthesis.
- **C6-Fluorine Revolution:** *Flumequine* added C6-fluorine (Fluoro-), launching 2nd-gen agents. -oxacin stem reflects exocyclic ketone/4-oxo oxygen.

C7-Piperazine Ring & Spectrum Expansion

- **C7-Piperazine Ring:** Added in 2nd-gen agents (Enoxacin, Ciprofloxacin) to cover *Pseudomonas* species.
- **3rd/4th Gen Advancement:** Ofloxacin expanded to Gram-positives (*S. aureus*); later agents optimized PK, anaerobes & Gram-positive potency.

Agent Name	Category / Generation	Etymological & Structural Origin
Ciprofloxacin	2nd Gen (Cyclopropyl)	Semisynthetic derivative of norfloxacin with an added cyclopropyl group. cipro- refers to the cyclopropyl group + -floxacin stem.
Ofloxacin	2nd Gen (Racemic Oxazine)	Synthetic racemic mixture. o- likely derived from the oxazine ring in drug structure + -floxacin stem.
Levofloxacin	3rd Gen (Pure L-Isomer)	Active L-isomer of ofloxacin, increasing potency, reducing required dose, and lowering toxicity vs racemic ofloxacin. Levo- refers to L-isomer form + -floxacin stem.

Tetracyclines & Glycylcyclines

4-Ring architecture, soil actinomycetes, and glycylamido derivatives (Beisel et al., Table 8)

📖 4-Ring Origins & Soil Actinomycetes

- **Structure:** Named for 4 (tetra-) fused rings (-cycl-).
- **Chlortetracycline (Aureomycin, 1948):** Benjamin Duggar from *S. aureofaciens* (Latin *aureus* = **golden mold**).
- **Oxytetracycline (Terramycin, 1950):** Isolated from *S. rimosus* in Terre Haute, IN (Latin *terra* = **earth/soil**).

🧬 PK Refinement & Glycylcycline Innovation

- **2nd Gen (Doxo / Mino):** Enhanced oral bioavailability & extended half-lives via chemical modifications.
- **Glycylcyclines (Tigecycline, etc.):** Added C-9 **glycylamido moiety** to defeat efflux pumps & ribosomal protection in MDR pathogen.

Agent Name	Category / Stem	Etymological & Structural Origin
Tetracycline	Class Prototype	Derived from catalytic hydrogenation of chlortetracycline; named based on chemical structure and loss of C-7 chlorine.
Doxycycline	2nd Gen (C6-Deoxy)	Semisynthetic from oxytetracycline. doxy- is a contraction of " deoxy- ", reflecting absence of hydroxyl group at C-6 position.
Minocycline	2nd Gen (C7-Dimethylamino)	Refers to its additional dimethylamino group at C-7 position, distinguishing it from prior tetracyclines.

Sulfonamides & Antifolate Agents

From Prontosil red azo dye to nucleic acid antimetabolites and folate synergy (Beisel et al., Table 9)



Domagk's Prontosil & WWII Sulfa Miracle

- **Gerhard Domagk (1932):** Discovered *Prontosil* (red azo dye); cleaved *in vivo* into active **sulfanilamide** ($-\text{SO}_2\text{NH}_2$).
- **WWII Field Medicine:** *Sulfathiazole* used extensively for Allied wound infections, saving countless soldiers.



Hitchings' Antifolate Synergy & Nobel Prize

- **George Hitchings (1988 Nobel Prize):** Nucleic acid antimetabolite research yielding Trimethoprim & Pyrimethamine.
- **Trimethoprim Etymology:** tri- (3 methoxy) + meth- + -o- (oxy) + -prim (pyrimidine). Formulated as **TMP-SMX (1968)**.

Agent Name	Category / Stem	Etymological & Structural Origin
Sulfanilamide	Azo Dye Metabolite	Active metabolite of Prontosil. Combination of <i>sulfanil(ic acid)</i> + <i>-amide</i> .
Sulfadiazine	Pyrimidine Derivative	Semisynthetic from sulfanilamide; -diazine refers to the pyrimidine ring attached to sulfa group.
Sulfamethoxazole	Isoxazole Derivative	Optimized for half-life. Conjugation of sulfa- + -methoxazole (from methoxy-isoxazole ring).

Glycopeptides & Lipoglycopeptides

From Borneo jungles to Mt. Ararat and lipoglycopeptide evolution (Beisel et al., Table 10)

Vancomycin & Lipoglycopeptide Evolution

- **Borneo Jungle (1953):** Edmund Kornfeld (Eli Lilly) isolated *Vancomycin* from *A. orientalis* soil sample; named from "**Vanquish**" (defeating MRSA).
- **2nd Gen Lipoglycopeptides:** *Telavancin* (Theravance + -vancin), *Dalbavancin*, *Oritavancin* with extended PK & lipophilic anchors.

Daptomycin & Membrane Depolarization

- **Mount Ararat, Turkey (1980s):** Isolated from *Streptomyces roseosporus* soil actinomycete.
- **Etymology & MOA:** dap- = **decanoyl** (fatty acid tail) + **amino peptide** core; membrane depolarization mechanism distinct from cell wall inhibition.

Agent Name	Category / Class	Etymological & Structural Origin
Vancomycin	Natural Glycopeptide	Isolated from <i>Amycolatopsis orientalis</i> (Borneo soil). Name derived from " vanquish " + -mycin (<i>Streptomyces</i> origin).
Teicoplanin	Natural Glycopeptide Complex	From <i>Actinoplanes teichomyceticus</i> (Indian soil). teicho- (Greek for wall , teichoic acid-like binding) + -planin .
Telavancin	Dual-MOA	Semisynthetic vancomycin derivative with lipophilic/hydrophilic moieties. Named after

Oxazolidinones & Lincosamides

From DuPont toxicity to Upjohn breakthroughs and Lincoln, Nebraska soil (Beisel et al., Tables 11 & 12)

🔥 Oxazolidinones: DuPont to Upjohn & Tedizolid Naming

- **Upjohn Breakthrough:** Steven Brickner overcame DuPont myelosuppression; **Linezolid** (1993) combines morpho line- + oxa -zolid inone.
- **Tedizolid Naming Journey:** TR-700 assigned *Tetrezolid* (tetrazole ring) → rejected by USAN → *Torezolid* → rejected → finalized as **Tedizolid**.

🌱 Lincosamides: Lincoln, Nebraska & Chlorination

- **Lincoln, Nebraska Soil (1960s): Lincomycin** isolated from *S. lincolnensis*; -amide refers to substituted prolinamide group.
- **Clindamycin Synergy:** Chlorinated derivative (cl- = **chlorinated**), dramatically improving oral bioavailability & anaerobic potency.

Agent Name	Class / Group	Etymological & Chemical Origin
Linezolid	Oxazolidinone Prototype	Fully synthetic compound. line- taken from morpholine structural component + -zolid from oxazolidinone class.
Tedizolid	2nd Gen Oxazolidinone	Fully synthetic compound. tedi- prefix finalized after <i>tetrezolid</i> (tetrazole ring) and <i>torezolid</i> were rejected by USAN.
Lincomycin	Lincosamide Prototype	Natural fermentation product named after <i>Streptomyces lincolnensis</i> , discovered from soil in Lincoln, Nebraska. -amide refers to substituted prolinamide group.

Rifamycins & Polymyxins

From 1950s French mob films to slimy soil bacilli and cationic polypeptides (Beisel et al., Tables 13 & 14)

Rifamycins: Lepetit Labs & "Rififi" Heist Film

- **Milan Discovery (1957):** Isolated by Piero Sensi's team from *Streptomyces mediterranei* (*A. rifamycinica*). Rifamycin B isolated in pure form.
- **French Mob Movie Naming:** Nicknamed after 1955 French crime film Rififi (slang for fight/scuffle) → **Rifamycin!**

Polymyxins: Slimy Bacilli & Cationic Polypeptides

- **Soil Origins (1947/1949):** Isolated from *Paenibacillus polymyxa* (Poly B) & *Bacillus colistinus* (Colistin / Poly E).
- **Etymology:** poly- = **polypeptide**; myx- = **slimy exudates** (exopolysaccharide). Chosen for lower toxicity.

Pleuromutilins: Mushroom Origin & C14 Modifications

- **NY Botanical Garden (1951):** Screened mushrooms *Pleurotus mutilus* & *P. passeckerianus* at Columbia Univ.
- **C-14 Side Chain & Stem:** Modified C-14 side chain for potency; universal stem -mutilin (Tiamulin, Retapamulin, Lefamulin).

Agent Name	Class / Group	Etymological & Structural Origin
Rifampin / Rifampicin	Rifamycin Prototype	Product of <i>A. rifamycinica</i> ; rif- from French film <i>Rififi</i> ; named after <i>N</i> -amino- <i>N</i> -methylpiperazine derivative → rif-amp-icin .
Rifabutin	Isobutyl Derivative	Derivative of rifamycin S; -butin adapted from " -butyl " referring to the isobutyl side chain.

Miscellaneous Antibiotics: Spanish Mountain Soil, Dental Miracles & Gut Targets

Fosfomycin, Metronidazole, and Fidaxomicin

Fosfomycin (Spanish Mountain Soil)

- **Mount Montgó, Spain (1969):** Merck & CEPA isolated *Streptomyces fradiae* near Alicante, Spain.
- **Etymology:** Originally *phosphonomycin* (epoxide + phosphonic acid structure); renamed **Fosfomycin** per INN standards.

Metronidazole (The 1962 Dental Miracle)

- **Rhône-Poulenc Antiparasitic:** Derived from *azomycin* for *Trichomonas vaginalis*.
- **Bedside Surprise:** 1962 patient treated for trichomoniasis had trench mouth (gingivitis) resolve simultaneously → premier antianaerobic agent!

Fidaxomicin (Flaw Becomes a Feature)

- **Hamden, CT Soil (1970s):** Abbott isolated *Tiacumicin B* (*tiacum-* from *D. aurantiacum*); abandoned due to 0% oral absorption.
- **2000s Revival:** 0% systemic absorption became the ultimate feature for gut-targeted *C. difficile* therapy! Retained *-micin* (actinomycete).

Miscellaneous Antibiotics: WWII Synthesis, Venezuelan Soil & Bacterial Antagonism

Nitrofurantoin, Chloramphenicol, and Mupirocin

Nitrofurantoin (WWII Synthetic Search)

- **Eaton Labs (1940s / 1953 FDA):** Synthesized during post-WWII search for novel antibacterials.
- **Etymology & Structure:** Nitrofuran ring + hydantoin side chain → **Nitro-furan-toin.**

Chloramphenicol (Venezuelan Soil Landmark)


- **Caracas Soil (1947):** Yale botanist Paul Burkholder isolated *S. venezuelae*; Mildred Rebstock (Parke-Davis) synthesized it (first natural antibiotic mass-produced!).
- **Name Breakdown:** chlor- (2 Cl) + -am- (amide) + -phen- (phenyl) + -icol (glycol).

Mupirocin (Bacterial Antagonism)

- **1890s Discovery to 1960s Beecham:** Freudenreich noticed *P. fluorescens* antagonism. Beecham isolated *pseudomonic acid A* for topical use.
- **Renaming Rationale:** Renamed *Mupirocin* (m- monic acid + -piro- *Pseudomonas* + -cin) to avoid implying activity was solely vs *Pseudomonas*.

SECTION

Antifungal Agents



Polyenes: Discovery, Chemical Nature & Formulation Evolution

Macrocyclic lactones with conjugated double bonds ("poly-")

Nystatin: Early Limitations & Toxicity

- **Structure & Origin:** Macrolide with conjugated double bonds ("poly-"). Discovered by NY State Dept of Health from *S. noursei*.
- **Oral & Systemic Failure:** Inadequate oral absorption → predicted inefficacious for systemic invasive fungal infections.
- **Formulation Toxicity:** Oral suspensions & IV attempts failed due to vein sclerosing and severe infusion reactions.

Amphotericin B (1953): Orinoco Soil & Lipid Rescue

- **Orinoco Screening (Jan 1953):** *Streptomyces nodosus* (M-4575) from Venezuela soil → Amphotericin A & B.
- **Tetraene Chromophore:** AmB demonstrated superior antifungal activity over AmA → primary clinical focus.
- **Deoxycholate Toxicity:** Poor oral absorption required IV sodium deoxycholate → high nephrotoxicity & infusion reaction risk.
- **Lipid Rescue (FDA 1995/1997):** Lipid Complex (ABLC, Nov 1995) & Liposomal (L-AmB, Aug 1997) approved to spare nephrons.

Polyene Agent	Microbial Source / Geographic Origin	Etymological & Chemical Naming	Clinical & Formulation Significance
Nystatin	<i>Streptomyces noursei</i>	Originally " <i>fungicidin</i> ". Renamed ny- (New York) + -sta- (State) + -tin (<i>Streptomyces</i>).	Broad spectrum; systemic toxicity limits use to topical & oral mucosal candidiasis.

Azoles: Chemical Evolution, Ring Architecture & Etymology

From topical imidazoles to mold-active triazoles & low-toxicity tetrazoles



Azole Class Evolution: Overcoming Polyenes

- **Rationale:** Developed after polyenes to reduce infusion reactions & dose-dependent nephrotoxicity of Amphotericin B.
- **Benzimidazole Foundation & Topical Era:** Pioneered by benzimidazole; first marketed topicals were **Clotrimazole** (1969), **Miconazole** (1969), and **Econazole** (1974).
- **Triazoles vs Molds:** Later triazoles (Voriconazole, Posaconazole, Isavuconazole) expanded coverage to molds (*Aspergillus*).



Nitrogen Ring Architecture & Suffix System

- **Class Suffix "-azole":** Refers to imidazole or triazole (3 nitrogens ["**tri-**"] in 5-membered ring) structural backbone.
- **Stem "-conazole":** Originated from Miconazole naming convention and perpetuated across the class.
- **Novel Tetrazoles (Oteseconazole):** Engineered with 4 nitrogens ("**tetra-**") + 1 carbon atom to decrease host toxicity.

Azole Agent	Ring Sub-Class	Etymological & Chemical Naming Rationale	Clinical / Chemical Significance
Clotrimazole	1st Gen Imidazole	clo- = chlorophenyl group + -im- = imidazole moiety.	First marketed topical product (1969).
Miconazole	1st Gen Imidazole	mico- = mycology (Greek <i>mykes</i> [fungus]) + -azole =	Established the standardized -

Glucan Synthase Inhibitors: Echinocandins & Triterpenoids

Targeting 1,3-β-D-glucan synthesis in the fungal cell wall

Echinocandins: Spiny Molds to Cell Wall Blockers

- **1970s Fermentation Discovery:** Echinocandin B isolated from *Aspergillus nidulans* var. *echinulatus*. Greek "echin-" = **spiny/prickly** (echinate conidiophores).
- **Overcoming Toxicity:** Early compounds caused severe hemolysis. Semisynthetic derivatives produced **Cilofungin** (first clinical trial) and **Pneumocandins** (*Zalerion arboricola*).
- **1990s Optimization:** Chemical tuning improved PK & solubility, yielding **Micafungin** (precursors from *Coleophoma empetria*) with potent anti-*Candida* activity.

Triterpenoids: Enfumafungin to Ibrexafungerp

- **Enfumafungin Landmark (2000):** Natural triterpene isolated from endophytic fungus *Hormonema carpetanum* via HTS.
- **Enfumafungin Breakdown:** en- (endophytic) + -fuma- (*A. **fuma**gatus*) + -fungin.
- **Ibrexafungerp Evolution:** Poor *in vivo* stability of enfumafungin led to semisynthetic **Ibrexafungerp** (first-in-class triterpenoid).
- **Suffix "-erp":** fung- = echinocandin-like spectrum; -erp = unique stem for **triterpenoids**.

Agent	Microbial Source / Derivative Origin	Etymological & Chemical Naming	Clinical & Structural Significance
Caspofungin	Semisynthetic from Pneumocandin B0 (<i>Zalerion arboricola</i>)	cas- = dual activity vs. <i>Candida</i> & <i>Aspergillus</i> spp.; -fungin = echinocandin stem.	First FDA-approved echinocandin (2001).

Other Antifungals: Repurposed Drugs, Allylamines & Chemical Diversity

Pyrimidines, allylamines, oxaboroles, hydroxypyridones, and fatty acids

Repurposed Pyrimidines & Penicillium Derivatives

- **Flucytosine Repurposing (1957 → 1960s):** Fluorinated cytosine synthesized in 1957 as an anticancer antimetabolite. Weak antineoplastic efficacy led to 1960s repurposing for systemic mycoses (cryptococcal meningitis).
- **Griseofulvin Color Origin:** Isolated from *Penicillium griseofulvum*. Latin *griseus* (**gray**) + *fulvus* (**tawny**) describing the gray-tawny pigment of the producing mold.

Allylamines & Diverse Chemical Classes

- **Allylamine Serendipity (Sandoz, Vienna):** Discovered in 1970s–1980s via serendipity + medicinal chemistry. Named for **allyl group** ($\text{CH}_2=\text{CH}-\text{CH}_2-$) bound to an **amine** (nitrogen).
- **Synthetic Diversity:** Encompasses Naphthalene cores (**Naftifine**, **Terbinafine**), Benzylamines (**Butenafine**), Oxaboroles (**Tavaborole**), Hydroxypyridones (**Ciclopirox**), and Castor oil derivatives (**Undecylenic acid**).

Antifungal Agent	Chemical Sub-Class	Etymological & Chemical Naming Rationale	Clinical / Biological Significance
Flucytosine	Fluorinated Pyrimidine	Fluorinated analogue of cytosine . Designed as anticancer antimetabolite (1957), repurposed as antifungal.	Synergistic oral agent for cryptococcal meningitis.
Griseofulvin	Penicillium Derivative	Derived directly from <i>Penicillium griseofulvum</i> . Latin <i>griseus</i> (gray) + <i>fulvus</i> (tawny).	Early oral agent for dermatophytosis & tinea capitis.

Epilogue: The Power of Antimicrobial Nomenclature

Connecting 80 years of drug discovery history with modern ICU antimicrobial stewardship

CORE THESIS

"Antimicrobial nomenclature is an evolutionary roadmap. Etymological roots transform drug names from arbitrary labels into immediate clinical insights on structure, spectrum, and pharmacokinetics."

Reference: Beisel C, et al. *What's in a Name? Etymology and History of Antibacterials and Antifungals*. Open Forum Infect Dis, 2026.