

Ciências / Ciência da computação / Introduction to the Theory of Computation (3rd Edition)

Exercício 16

Capítulo 2, Página 156

...



Introduction to the Theory of Computation

ISBN: 9781133187790

[Índice](#)**Solução** Certificado Solução fornecida há 2 anos**Passo 1**

1 de 2

Union and concatenation are trivial. Assume that we are given languages L_1 and L_2 generated by context-free grammars G_1 and G_2 respectively. Then grammar which includes **all rules of both** grammar G_1 and G_2 and **one additional rule** $S \rightarrow S_1 \mid S_2$, where S is the new starting variable of grammar G , and S_1 and S_2 are old starting variables of grammars G_1 and G_2 , recognizes exactly **union** $L_1 \cup L_2$. Concatenation is obtained analogously, by adding rule $S \rightarrow S_1 S_2$.

Star is also not difficult, but last $\text{\textit{Exercise}}$ shows th

$$S_0 \rightarrow S_0 S_0 \mid S \mid \varepsilon$$

to the new grammar, where S is the old starting variable. Note the analogy with adding a new initial state to NFA which recognizes star of language.

Resultado

2 de 2

We modify the context-free grammars generating given languages to capture their union, concatenation and star.

Avaliar esta solução[< Exercício 15](#)[Exercício 17 >](#)

[Privacidade](#) [Termos de serviço](#)

[Português \(BR\)](#) ▾