


THE DIAGONAL OF THE MULTIPLIHEDRA AND THE TENSOR PRODUCT OF A_∞ -MORPHISMS

GUILLAUME LAPLANTE-ANFOSSI AND THIBAUT MAZUIR

ABSTRACT. We define a cellular approximation for the diagonal of the Forcey–Loday realizations of the multiplihedra, and endow them with a compatible topological cellular operadic bimodule structure over the Loday realizations of the associahedra. This provides us with a model for topological and algebraic A_∞ -morphisms, as well as a universal and explicit formula for their tensor product. We study the monoidal properties of this newly defined tensor product and conclude by outlining several applications, notably in algebraic and symplectic topology.



Date: July 19, 2022.

2010 Mathematics Subject Classification. Primary 52B11, 18M70.

Key words and phrases. Multiplihedra, approximation of the diagonal, associahedra, operads, tensor product, A -infinity algebras, A -infinity morphisms, A -infinity categories.

The first author was supported by the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 754362 , by the Natural Sciences and Engineering Research Council of Canada (NSERC) and by the ANR-20-CE40-0016 (Higher Algebra, Geometry and Topology). The second author was supported by the ANR-21-CE40-0002 (New challenges in Symplectic and Contact Topology).

CONTENTS

Introduction	2
1. Realizations of the multiplihedra	4
2. Diagonal of the multiplihedra	11
3. Cellular formula for the diagonal of the multiplihedra	15
4. Tensor product of A_∞ -morphisms and A_∞ -functors	20
5. Further applications	31
References	37

INTRODUCTION

The n -dimensional associahedron, a polytope whose faces are in bijection with planar trees with $n + 2$ leaves, was first introduced as a topological cell complex by J. Stasheff to describe algebras whose product is associative up to homotopy [Sta63]. The problem of giving polytopal realizations of these CW-complexes has a rich history [CZ12], and the algebras that they encode, called A_∞ -algebras, have been extensively studied in various branches of mathematics. They were used in algebraic topology for the study of iterated loop spaces [May72, BV73] or the study of homotopy theory of differential graded associative algebras [LH03, Val20] ; in symplectic topology to define Fukaya categories of symplectic manifolds [Sei08, FOOO09a, FOOO09b], through the interpretation of the associahedra as moduli spaces of disks with marked boundary points; and more recently, in mathematical physics, mirror symmetry, Galois cohomology or non-commutative probability.

The n -dimensional multiplihedron is a polytope whose faces are in bijection with 2-colored planar trees with $n + 1$ leaves. It was first introduced as a topological cell complex by J. Stasheff to describe morphisms between A_∞ -algebras [Sta70]. It was only recently realized as a convex polytope in the work of S. Forcey [For08], followed by the work of S. Forcey and S. Devadoss [DF08], F. Ardila and J. Doker [AD13], and F. Chapoton and V. Pilaud [CP22]. The multiplihedra were studied in algebraic topology [BV73], as well as in symplectic topology [MW10, MWW18] and Morse theory [Maz21a, Maz21b], as they can be respectively realized as moduli spaces of quilted disks with marked boundary points and as moduli spaces of 2-colored metric trees.

In this paper, we define and study a cellular approximation of the diagonal of the multiplihedra. The need for such an approximation comes from the fact that the standard thin diagonal $\Delta_P : P \rightarrow P \times P, x \mapsto (x, x)$ of a polytope P is not cellular in general, i.e. its image is not a union of faces of $P \times P$. A cellular approximation of the diagonal is a cellular map $\Delta_P^{\text{cell}} : P \rightarrow P \times P$ which is homotopic to Δ_P and which agrees with Δ_P on the vertices of P .

The Alexander–Whitney map [EML53] and the Serre diagonal [Ser51] respectively define cellular approximations for the diagonal of the simplices and for the diagonal of the cubes, yielding the cup product in singular cohomology and the cup product in cubical cohomology. A cellular approximation for the diagonal of the associahedra was constructed in [MTTV21] and yields a universal formula for the tensor product of two A_∞ -algebras. See also [SU04, MS06]. By the term *universal*, we mean that the same formula applies uniformly to any pair of A_∞ -algebras. In a similar fashion, the cellular approximation of the diagonal of the

multiplihedra will be used to define a universal tensor product of A_∞ -morphisms in this paper. Our main results can be summarized as follows.

- (1) We define a cellular approximation of the diagonal on Forcey–Loday realizations of the multiplihedra (Definition 2.12).
- (2) We endow them with a compatible operadic bimodule structure over the Loday realizations of the associahedra (Theorem 1).
- (3) We compute explicitly the associated combinatorial formula for the cellular image of the diagonal (Theorem 2).
- (4) We apply the cellular chains functor to the diagonal in order to define a universal tensor product of A_∞ -morphisms (Proposition 4.18), and we study its properties (Section 4.4).

To achieve these goals, we use the theory of cellular approximations of diagonals developed by the first author in [Lap22], which is based on the theory of fiber polytopes of [BS92] and the method introduced in [MTTV21]. We prove that the Forcey–Loday realizations of the multiplihedra [For08] can be obtained from the Ardila–Doker realization of the multiplihedra [AD13] by projection (Proposition 1.16). These last realizations are generalized permutahedra, in the sense of A. Postnikov [Pos09], which allows us to apply the results of [Lap22] directly, both to define a cellular approximation of the diagonal and to describe its cellular image combinatorially.

The tensor product of A_∞ -morphisms defined by this diagonal does not however define a symmetric monoidal structure on the category $\infty\text{-}A_\infty\text{-alg}$ of A_∞ -algebras and their A_∞ -morphisms, since it is not strictly compatible with the composition. This is not a defect of our construction: in Proposition 4.25, we prove that there is no tensor product of A_∞ -morphisms which is strictly compatible with the composition of A_∞ -morphisms. This proposition should be compared to a similar result by M. Markl and S. Shnider, saying that there is no strictly associative tensor product of A_∞ -algebras [MS06, Theorem 13]. The preceding two properties are in fact always satisfied up to homotopy (see Proposition 4.26), which points towards the idea that the category $\infty\text{-}A_\infty\text{-alg}$ should possess some kind of *homotopy* symmetric monoidal structure. An analogous phenomenon was already observed for the category of homotopy representations of an algebraic group [AACD11, Pol20].

Our results can be readily applied to different fields. The operadic bimodule structure of Point (2) above was used in the work of the second author, in order to realize A_∞ -algebras and A_∞ -morphisms in Morse theory [Maz21a, Maz21b]. The algebraic tensor product in Point (4) has applications in Heegaard Floer homology and could be used to relate the Fukaya categories of products of symplectic manifolds via Lagrangian correspondences, see Section 5.3. We also expect future applications of our work to the computation of the homology of fibered spaces, using the construction of the convolution A_∞ -algebra associated to an A_∞ -coalgebra and an A_∞ -algebra in Proposition 5.4. This last construction can also be related to the deformation theory of ∞ -morphisms developed in [RNW19b, RNW19a], see Section 5.2.3. Moreover, our geometric methods shed a new light on a result of M. Markl and S. Shnider [MS06], pointing towards possible links with discrete and continuous Morse theory (Remark 5.3).

Finally, the results of this paper can be straightforwardly extended to the "multioperalihedra", a family of polytopes which is to the operahedra of [Lap22] what the multiplihedra are to the associahedra. They belong at the same time to the families of graph-multiplihedra [DF08] and of nestomultiplihedra [AD13]. Together with the results of [Lap22, Section 4],

one would obtain a tensor product of ∞ -morphisms between homotopy operads, defined by explicit formulæ.

Layout. We introduce the Forcey–Loday and the Ardila–Doker realizations of the multiplihedra in Section 1. We define a cellular approximation of their diagonal and endow the Forcey–Loday multiplihedra with an operadic bimodule structure over the Loday associahedra in Section 2. We compute explicitly the associated combinatorial formula for the image of our diagonal in Section 3. We define a tensor product of A_∞ -algebras and of A_∞ -morphisms and study its properties in Section 4. We finally sketch future applications of our work in Section 5.

Conventions. We use the conventions and notations of [Zie95] for convex polytopes and the ones of [LV12] for operads. The word operad will always mean non-symmetric operad [LV12, Section 5.2.8] in this paper. We denote by $[n] := \{1, \dots, n\}$ and by $\{e_i\}_{i \in [n]}$ the standard basis of \mathbb{R}^n . The abbreviation "dg" will stand for the words "differential graded".

Acknowledgements. We would like to thank Bruno Vallette for numerous discussions and for his careful reading of our paper, as well as Alexandru Oancea and Eric Hoffbeck for their comments on earlier versions. We are also indebted to Lino Amorim and Robert Lipshitz, for explaining to us their work and for their detailed insights on possible applications of our results in symplectic topology. We finally express our gratitude to Sushmita Venugopalan, for taking the time to discuss potential connections between our work and results on toric varieties, and to Daniel Robert-Nicoud, for discussing his work with us and suggesting new directions of research.

1. REALIZATIONS OF THE MULTIPLIHEDRA

Drawing from the work of Forcey in [For08], we define the weighted Forcey–Loday realizations of the multiplihedra and describe their geometric properties in Proposition 1.10. We then show how they can be recovered from the Ardila–Doker realizations of the multiplihedra, which are in particular generalized permutahedra.

1.1. 2-colored trees and multiplihedra.

1.1.1. *2-colored trees.* We consider in this section *planar rooted trees*, which we simply abbreviate as *trees*. The term *edge* refers to both internal and external edges. The external edges will sometimes be called leaves.

Definition 1.1 (Cut). *A cut of a tree is a subset of edges or vertices which contains precisely one edge or vertex in each non-self crossing path from an incoming edge to the root.*

A cut divides a tree into an upper part that we color in blue and a lower part that we color in red. The edges and vertices of the cut are represented by drawing a black line over them, as pictured in Figure 1.

Definition 1.2 (2-colored tree). *A 2-colored tree is a tree together with a cut. We call 2-colored maximal tree a 2-colored binary tree whose cut is made of edges only.*

We denote by CT_n (resp. CMT_n) the set of 2-colored trees (resp. 2-colored maximal trees) with n leaves, for $n \geq 1$.

Definition 1.3 (Face order). *The face order $s \subset t$ on 2-colored trees is defined as follows: a 2-colored tree s is less than a 2-colored tree t if t can be obtained from s by a sequence of contractions of monochrome edges or moves of the cut from a family of edges to an adjacent vertex.*





FIGURE 1. Two 2-colored trees, related by the face order.

Definition 1.4 (Tamari-type order). *The Tamari-type order $s < t$ on 2-colored maximal trees is generated by the following three covering relations:*



where each t_i , $1 \leq i \leq 4$, is a binary tree of the appropriate color.

We add a minimum element \emptyset_n to the poset of 2-colored trees (CT_n, \subset) .

Proposition 1.5. *The posets (CT_n, \subset) and $(CMT_n, <)$ are lattices.*

Proof. The poset of 2-colored trees was proven in [For08] to be isomorphic to the face lattice of a polytope, the multiplihedron; see Point (3) of Proposition 1.10. The Hasse diagram of the poset of 2-colored maximal trees was proven to be isomorphic to the oriented 1-skeleton of the multiplihedron, and also to be the Hasse diagram of a lattice in [CP22, Proposition 117]. \square

REMARK 1.6. F. Chapoton and V. Pilaud introduced in [CP22] the shuffle of two generalized permutahedra (see Section 1.3 for definition and examples). The fact that the poset $(CMT_n, <)$ is a lattice follows from the fact that the multiplihedron arises as the shuffle of the associahedron and the interval, which both have the lattice property, and that the shuffle operation preserves the lattice property in this case, see [CP22, Corollary 95].

1.1.2. Grafting of trees. We will denote the operation of grafting a planar tree v at the i^{th} -leaf of a 2-colored tree u by $u \circ_i v$. We will also denote the grafting of a level of 2-colored trees v_1, \dots, v_k on the k leaves of a planar tree by $u(v_1, \dots, v_k)$. We denote by c_n^T and by c_n^B the corollae with n leaves fully painted with the upper and the lower color respectively; we denote by c_n the corolla with n leaves with frontier color at the vertex. It is straightforward to see that these two grafting operations on corollae generate all the 2-colored trees of codimension 1: we call (B), for “bottom”, the first type of 2-colored trees $c_{p+1+r} \circ_{p+1} c_q^T$, with $p + q + r = n$

and $2 \leq q \leq n$, and we call (T), for “top”, the second type of 2-colored trees $c_k^B(c_1, \dots, c_k)$, with $i_1 + \dots + i_k = n$, $i_1, \dots, i_k \geq 1$, and $k \geq 2$.




FIGURE 2. Examples of 2-colored trees of type (B) and (T) respectively.

1.1.3. Multiplihedra.

Definition 1.7 (Multiplihedra). *For any $n \geq 1$, an $(n - 1)$ -dimensional multiplihedron is a polytope of dimension $(n - 1)$ whose face lattice is isomorphic to the lattice (CT_n, \subset) of 2-colored trees with n leaves.*




FIGURE 3. A 2-dimensional multiplihedron and the Tamari-type poset $(CMT_3, <)$ on its oriented 1-skeleton.

The dimension of a face labeled by a 2-colored tree is given by the sum of the degrees of its vertices defined by

$$\left| \begin{array}{c} 1 \dots k \\ \text{blue tree} \end{array} \right| = k - 2, \quad \left| \begin{array}{c} 1 \dots k \\ \text{red tree} \end{array} \right| = k - 2, \quad \left| \begin{array}{c} 1 \dots k \\ \text{blue tree with red root} \end{array} \right| = k - 1.$$

The codimension of a 2-colored tree is then equal to the number of blue and red vertices. In the example of the 2-colored tree depicted on the left of Figure 1, the dimension is equal to 4 and the codimension is equal to 5. As proven in [CP22, Proposition 117], the oriented 1-skeleton of a multiplihedron is the Hasse diagram of the Tamari-type poset.

1.2. Forcey–Loday realizations of the multiplihedra. Jean-Louis Loday gave in [Lod04] realizations of the associahedra in the form of polytopes with integer coordinates. Stefan Forcey generalized this construction in [For08] in order to give similar realizations for the multiplihedra.

Definition 1.8 (Weighted 2-colored maximal tree). *A weighted 2-colored maximal tree is a pair (t, ω) made up of a 2-colored maximal tree $t \in \text{CMT}_n$ with n leaves with a weight $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{R}_{>0}^n$. We call ω the weight and n the length of the weight ω .*

Let (t, ω) be a weighted 2-colored maximal tree with n leaves. We order its $n-1$ vertices from left to right. At the i^{th} vertex, we consider the sum α_i of the weights of the leaves supported by its left input and the sum β_i of the weights of the leaves supported by its right input. If the i^{th} vertex is colored by the upper color, we consider the product $\alpha_i \beta_i$ and if the i^{th} vertex is colored by the lower color, we consider the product $2\alpha_i \beta_i$. The associated string produces a point with integer coordinates $M(t, \omega) \in \mathbb{R}_{>0}^{n-1}$. For example, if only the first and last vertices of t are blue, we obtain a point of the form

$$M(t, \omega) = (2\alpha_1\beta_1, \alpha_2\beta_2, \dots, \alpha_{n-2}\beta_{n-2}, 2\alpha_{n-1}\beta_{n-1}) \in \mathbb{R}_{>0}^{n-1}.$$




FIGURE 4. Examples of points associated to 2-colored maximal trees, with standard weight.

Definition 1.9 (Forcey–Loday Realization). *The Forcey–Loday realization of weight ω of the $(n-1)$ -dimensional multiplihedron is the polytope*

$$J_\omega := \text{conv} \{M(t, \omega) \mid t \in \text{CMT}_n\} \subset \mathbb{R}^{n-1}.$$

The Forcey–Loday realization associated to the standard weight $(1, \dots, 1)$ will simply be denoted by J_n . By convention, we define the polytope J_ω with weight $\omega = (\omega_1)$ of length 1 to be made up of one point labeled by the 2-colored tree $i_B^T := \dagger$.




FIGURE 5. The Forcey–Loday realization of the multiplihedron J_4 .

Proposition 1.10. *The Forcey–Loday realization J_ω satisfies the following properties.*

(1) *Let $t \in \text{CMT}_n$ be a 2-colored maximal tree.*

For $p+q+r=n$, with $2 \leq q \leq n$, the point $M(t, \omega)$ is contained in the half-space defined by the inequality

$$(B) \quad x_{p+1} + \dots + x_{p+q-1} \geq \sum_{p+1 \leq a < b \leq p+q} \omega_a \omega_b,$$

with equality if and only if the 2-colored maximal tree t can be decomposed as $t = u \circ_{p+1} v$, where $u \in \text{CMT}_{p+1+r}$ and $v \in \text{PBT}_q$.

For $i_1 + \dots + i_k = n$, with $i_1, \dots, i_k \geq 1$ and $k \geq 2$, the point $M(t, \omega)$ is contained in the half-space defined by the inequality

$$(T) \quad x_{i_1} + x_{i_1+i_2} + \dots + x_{i_1+\dots+i_{k-1}} \leq 2 \sum_{1 \leq j < l \leq k} \omega_{I_j} \omega_{I_l},$$

where $I_j = [i_1 + \dots + i_{j-1} + 1, \dots, i_1 + \dots + i_j]$ and $\omega_{I_j} := \sum_{a \in I_j} \omega_a$, with equality if and only if the 2-colored maximal tree t can be decomposed as $t = u(v_1, \dots, v_k)$, where $u \in \text{PBT}_k$ and $v_j \in \text{CMT}_{i_j}$, for $1 \leq j \leq k$.

(2) *The polytope J_ω is the intersection of the half-spaces defined in (1).*

- (3) The face lattice $(\mathcal{L}(J_\omega), \subset)$ is isomorphic to the lattice (CT_n, \subset) of 2-colored trees with n leaves.
- (4) Any face of a Forcey–Loday realization of a multiplihedron is isomorphic to a product of a Loday realization of an associahedron with possibly many Forcey–Loday realizations of multiplihedra, via a permutation of coordinates.

Proof. Points (1)–(3) were proved in [For08]. We prove Point (4) by induction on n . It clearly holds true for $n = 1$. Let us suppose that it holds true up to $n - 1$ and let us prove it for the polytopes J_ω , for any weight ω of length n . We examine first facets. In the case of a facet of type (B) associated to $p + q + r = n$ with $2 \leq q \leq n - 1$, we consider the following two weights

$$\bar{\omega} := (\omega_1, \dots, \omega_p, \omega_{p+1} + \dots + \omega_{p+q}, \omega_{p+q+1}, \dots, \omega_n) \quad \text{and} \quad \tilde{\omega} := (\omega_{p+1}, \dots, \omega_{p+q})$$

and the isomorphism

$$\Theta_{p,q,r} : \mathbb{R}^{p+r} \times \mathbb{R}^{q-1} \xrightarrow{\cong} \mathbb{R}^{n-1} \\ (x_1, \dots, x_{p+r}) \times (y_1, \dots, y_{q-1}) \mapsto (x_1, \dots, x_p, y_1, \dots, y_{q-1}, x_{p+1}, \dots, x_{p+r}).$$

The image of the vertices of $J_{\bar{\omega}} \times K_{\tilde{\omega}}$ are sent to the vertices of the facet of J_ω labelled by the 2-colored tree $c_{p+1+r} \circ_{p+1} c_q^T$. In other words, the permutation of coordinates Θ sends bijectively $J_{\bar{\omega}} \times K_{\tilde{\omega}}$ to J_ω . Similarly, in the case of a facet of type (T) associated to $i_1 + \dots + i_k = n$ with $i_1, \dots, i_k \geq 1$ and $k \geq 2$, we consider the following weights

$$\bar{\omega} := (\sqrt{2}\omega_{i_1}, \dots, \sqrt{2}\omega_{i_k}) \quad \text{and} \quad \tilde{\omega}_j := (\omega_{i_1+\dots+i_{j-1}+1}, \dots, \omega_{i_1+\dots+i_{j-1}+i_j}), \text{ for } 1 \leq j \leq k,$$

and the isomorphism

$$\Theta^{i_1, \dots, i_k} : \mathbb{R}^{k-1} \times \mathbb{R}^{i_1-1} \times \dots \times \mathbb{R}^{i_k-1} \xrightarrow{\cong} \mathbb{R}^{n-1}$$

which sends

$$(x_1, \dots, x_{k-1}) \times (y_1^1, \dots, y_{i_1-1}^1) \times \dots \times (y_1^k, \dots, y_{i_k-1}^k)$$

to

$$(y_1^1, \dots, y_{i_1-1}^1, x_1, y_1^2, \dots, y_{i_2-1}^2, x_2, y_1^3, \dots, x_{k-1}, y_1^k, \dots, y_{i_k-1}^k).$$

The image of the vertices of $K_{\bar{\omega}} \times J_{\tilde{\omega}_1} \times \dots \times J_{\tilde{\omega}_k}$ are sent to the vertices of the facet of J_ω labelled by the 2-colored tree $c_k^B(c_1, \dots, c_k)$. In other words, the permutation of coordinates Θ sends bijectively $K_{\bar{\omega}} \times J_{\tilde{\omega}_1} \times \dots \times J_{\tilde{\omega}_k}$ to J_ω .

We can finally conclude the proof with these decompositions of facets of J_ω , the induction hypothesis, and Point (5) of [MTTV21, Proposition 1]. \square

1.3. Ardila–Doker realizations of the multiplihedra.

Definition 1.11 (Permutahedron). *The $(n - 1)$ -dimensional permutahedron is the polytope in \mathbb{R}^n equivalently defined as:*

- the convex hull of the points $\sum_{i=1}^n i e_{\sigma(i)}$ for all permutations $\sigma \in \mathbb{S}_n$, or
- the intersection of the hyperplane $\left\{ x \in \mathbb{R}^n \mid \sum_{i=1}^n x_i = \binom{n+1}{2} \right\}$ with the affine half-spaces $\left\{ x \in \mathbb{R}^n \mid \sum_{i \in I} x_i \geq \binom{|I|+1}{2} \right\}$ for all $\emptyset \neq I \subseteq [n]$.

For a face F of a polytope $P \subset \mathbb{R}^n$, the *normal cone* of F is the cone

$$\mathcal{N}_P(F) := \left\{ c \in (\mathbb{R}^n)^* \mid F \subseteq \{x \in P \mid cx = \max_{y \in P} cy\} \right\}.$$

The codimension of $\mathcal{N}_P(F)$ is equal to the dimension of F . The *normal fan* of P is the collection of the normal cones $\mathcal{N}_P := \{\mathcal{N}_P(F) \mid F \in \mathcal{L}(P) \setminus \emptyset\}$. We refer to [Zie95, Chapter 7] for more details.

Definition 1.12 (Generalized permutahedron). *A generalized permutahedron is a polytope equivalently defined as:*

- a polytope whose normal fan coarsens the one of the permutahedron, or
- the convex set

$$\left\{ x \in \mathbb{R}^n : \sum_{i=1}^n x_i = z_{[n]}, \sum_{i \in I} x_i \geq z_I \text{ for all } I \subseteq [n] \right\},$$

where $\{z_I\}_{I \subseteq [n]}$ are real numbers which satisfy the inequalities $z_I + z_J \leq z_{I \cup J} + z_{I \cap J}$ for all $I, J \subseteq [n]$, and where $z_\emptyset = 0$.

Generalized permutahedra were introduced by A. Postnikov in [Pos09]. Loday realizations of the associahedra are all generalized permutahedra (see [Pos09, Corollary 8.2]), while Forcey–Loday realizations of the multiplihedra are not. However, F. Ardila and J. Doker introduced in [AD13] realizations of the multiplihedra that are generalized permutahedra. They are obtained from the Loday realizations of the associahedra via the operation of *q-lifting*. We will consider the special case $q = 1/2$ of their construction.

Definition 1.13 (Lifting of a generalized permutahedron [AD13, Definition 2.3]). *For a generalized permutahedron $P \subset \mathbb{R}^n$, its $\frac{1}{2}$ -lifting $P\left(\frac{1}{2}\right) \subset \mathbb{R}^{n+1}$ is defined by*

$$P\left(\frac{1}{2}\right) := \left\{ x \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} x_i = z_{[n]}, \sum_{i \in I} x_i \geq \frac{1}{2} z_I, \sum_{i \in I \cup \{n+1\}} x_i \geq z_I \text{ for all } I \subseteq [n] \right\}.$$

Proposition 1.14 ([AD13, Proposition 2.4]). *The $\frac{1}{2}$ -lifting $P\left(\frac{1}{2}\right)$ of a generalized permutahedron is again a generalized permutahedron.*

Proposition 1.15. *The $\frac{1}{2}$ -lifting $K_\omega\left(\frac{1}{2}\right)$ of the Loday realization of weight ω of the associahedron is a realization of the multiplihedron.*

Proof. This is a particular case of [AD13, Corollary 4.10]. □

We call the lifting of the Loday associahedron $K_\omega\left(\frac{1}{2}\right)$ the *Ardila–Doker realization* of the multiplihedron. It is related to the Forcey–Loday realization via the projection $\pi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ which forgets the last coordinate.

Proposition 1.16. *The Forcey–Loday realization of the multiplihedron is the image under the projection π of the $\frac{1}{2}$ -lifting of the Loday realization of the associahedron, scaled by 2. That is, we have*

$$J_\omega = \pi\left(2K_\omega\left(\frac{1}{2}\right)\right).$$

Proof. This follows from the vertex description of $\frac{1}{2}$ -lifting given in [Dok11, Definition 3.5.3], together with the description of the projection from the permutahedron to the multiplihedron given in the proof of [Dok11, Theorem 3.3.6]. The coordinates of a vertex in $2K_\omega$ are of the form $(2\alpha_1\beta_1, \dots, 2\alpha_n\beta_n)$. A coordinate $2\alpha_i\beta_i$ is then multiplied by $1/2$ in the lifting if and only if its associated vertex in the 2-colored maximal tree is of the upper color. We thus recover the description of Definition 1.9. \square

In summary, we have the following diagram:

Loday associahedron		Ardila–Doker multiplihedron		Forcey–Loday multiplihedron
K_ω	\hookrightarrow	$K_\omega \left(\frac{1}{2} \right)$	$\xrightarrow{\pi^{(2\cdot)}}$	J_ω
\mathbb{R}^n	\hookrightarrow	\mathbb{R}^{n+1}	\rightarrow	\mathbb{R}^n
Gen. permutahedron		Gen. permutahedron		Not a gen. permutahedron

2. DIAGONAL OF THE MULTIPLIHEDRA

In this section, we define a cellular approximation of the diagonal of the Forcey–Loday realizations of the multiplihedra, and we endow them with an operadic bimodule structure over the Loday realizations of the associahedra in the category Poly. We use the methods of [MTTV21] and the general theory developed in [Lap22]. Our construction of the cellular approximation relies crucially on the fact that the Forcey–Loday multiplihedra, are obtained from the Ardila–Doker multiplihedra by projection (Proposition 1.16).

2.1. The monoidal category Poly. Let us recall the definition of the symmetric monoidal category (Poly, \times) from [MTTV21, Section 2.1].

OBJECTS: An object of Poly is a d -dimensional polytope P in the n -dimensional Euclidian space \mathbb{R}^n , for any $0 \leq d \leq n$.

MORPHISMS: A morphism in Poly is a continuous map $f : P \rightarrow Q$ which sends P homeomorphically to the underlying set $|\mathcal{D}|$ of a polytopal subcomplex $\mathcal{D} \subset \mathcal{L}(Q)$ of Q such that $f^{-1}(\mathcal{D})$ defines a polytopal subdivision of P .

We will use the notion of *operad*, *operadic bimodule* and *Hadamard product* of operads and operadic bimodules in the rest of this paper. For the sake of concision, we refer respectively to [Maz21a, Section 1.1.1], [Maz21a, Section 1.1.3] and [LV12, Section 5.1.12] for a complete definition of these notions. An operad will in particular be a non-symmetric operad in the language of [LV12, Section 5.2.8]. The fact that the category Poly is monoidal will moreover allow us to define operads and operadic bimodules in polytopes.

2.2. Positively oriented polytopes and diagonal maps. For a polytope P , we will denote by $\rho_z P := 2z - P$ its reflection with respect to a point $z \in P$.

Definition 2.1. A positively oriented polytope (P, \vec{v}) is a polytope $P \subset \mathbb{R}^n$ together with a vector $\vec{v} \in \mathbb{R}^n$ which is not perpendicular to any edge of $P \cap \rho_z P$, for any $z \in P$.

Any positively oriented polytope admits a diagonal map of the form

$$\begin{aligned} \Delta_{(P, \vec{v})} : P &\rightarrow P \times P \\ z &\mapsto (\text{bot}_{\vec{v}}(P \cap \rho_z P), \text{top}_{\vec{v}}(P \cap \rho_z P)) . \end{aligned}$$

Such a diagonal map is a morphism in Poly, coincides with the usual thin diagonal $x \mapsto (x, x)$ on vertices, and is fiber-homotopic to it, see [MTTV21, Proposition 5] and [Lap22, Proposition 1.1]. Its cellular image admits a combinatorial description in terms of the fundamental hyperplane arrangement of P , as we will now recall.

Definition 2.2 (Fundamental hyperplane arrangement). *An edge hyperplane of P is an hyperplane in \mathbb{R}^n which is orthogonal to the direction of an edge of $P \cap \rho_z P$ for some $z \in P$. The fundamental hyperplane arrangement \mathcal{H}_P of P is the collection of all edge hyperplanes of P .*

Recall that a face F of a polytope $P \subset \mathbb{R}^n$ is equal to the intersection of a family of facets $\{F_i\}$. If we choose an outward pointing normal vector \vec{F}_i for each facet F_i (see [Lap22, Definition 1.24]) and a basis $\{b_k\}$ of the orthogonal complement of the affine hull of P in \mathbb{R}^n , then the normal cone of F is given by $\mathcal{N}_P(F) = \text{Cone}(\{\vec{F}_i\} \cup \{b_k, -b_k\})$.

Proposition 2.3 ([Lap22, Theorem 1.23]). *Let (P, \vec{v}) be a positively oriented polytope in \mathbb{R}^n . For each $H \in \mathcal{H}_P$, we choose a normal vector \vec{d}_H such that $\langle \vec{d}_H, \vec{v} \rangle > 0$. We have*

$$(F, G) \in \text{Im } \Delta_{(P, \vec{v})} \iff \forall H \in \mathcal{H}_P, \exists i, \langle \vec{F}_i, \vec{d}_H \rangle < 0 \text{ or } \exists j, \langle \vec{G}_j, \vec{d}_H \rangle > 0 .$$

We finally recall general facts from [Lap22, Section 1.6].

Definition 2.4 (Coarsening projection). *Let P and Q be two polytopes in \mathbb{R}^n such that the normal fan of P refines the normal fan of Q . The coarsening projection from P to Q is the application $\theta : \mathcal{L}(P) \rightarrow \mathcal{L}(Q)$ which sends a face F of P to the face $\theta(F)$ of Q whose normal cone $\mathcal{N}_Q(\theta(F))$ is the minimal cone with respect to inclusion which contains $\mathcal{N}_P(F)$.*

Proposition 2.5. *Let P and Q be two polytopes such that the normal fan of P refines the one of Q . If P is positively oriented by \vec{v} , then so is Q . Moreover, the coarsening projection from P to Q commutes with the diagonal maps $\Delta_{(P, \vec{v})}$ and $\Delta_{(Q, \vec{v})}$, and we have*

$$(F, G) \in \text{Im } \Delta_{(Q, \vec{v})} \iff \forall H \in \mathcal{H}_P, \exists i, \langle \vec{F}_i, \vec{d}_H \rangle < 0 \text{ or } \exists j, \langle \vec{G}_j, \vec{d}_H \rangle > 0 .$$

We will apply Proposition 2.5 to P the permutahedron and Q the Ardila–Doker multiplihe-dron, in order to define a diagonal map on the Forcey–Loday multiplihedron and to compute an explicit formula for its cellular image in Theorem 2.

2.3. Good orientation vectors and generalized permutahedra. The projection $\pi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ forgetting the last coordinate defines an affine isomorphism between any hyperplane H of equation $\sum_{i=1}^{n+1} x_i = c \in \mathbb{R}$, and \mathbb{R}^n . The inverse map $(\pi|_H)^{-1}$ is given by the assignment

$$(x_1, \dots, x_n) \mapsto \left(x_1, \dots, x_n, c - \sum_{i=1}^n x_i \right) .$$

If a polytope P is contained in the hyperplane H , then the polytope $\pi(P)$ is affinely isomorphic to P , and the projection π defines a bijection between the faces of P and the faces of $\pi(P)$. Moreover, for every face F of P , we have $\dim F = \dim \pi(F)$.

However, the projection π does not preserve orthogonality in general, so if P is positively oriented by \vec{v} , the projection $\pi(P)$ might not be positively oriented by $\pi(\vec{v})$. We restrict our attention to a certain class of orientation vectors for which this property holds, in the case where P is a generalized permutahedron.

Definition 2.6. *A good orientation vector is a vector $\vec{v} = (v_1, \dots, v_{n+1}) \in \mathbb{R}^{n+1}$ satisfying*

$$v_i \geq 2v_{i+1}, \text{ for any } 1 \leq i \leq n, \quad \text{and} \quad v_{n+1} > 0.$$

Observe that the family of good orientation vectors is stable under the projection forgetting the last coordinate: if \vec{v} is a good orientation vector, then so is $\pi(\vec{v})$. Being a good orientation vector is a more restrictive condition than being a principal orientation vector in the sense of [Lap22, Definition 3.15]. Thus, a good orientation vector orients positively any generalized permutahedron.

Proposition 2.7. *Let $P \subset \mathbb{R}^{n+1}$ be a generalized permutahedron, and let $\vec{v} \in \mathbb{R}^{n+1}$ be a good orientation vector. Then, the polytope $\pi(P)$ is positively oriented by $\pi(\vec{v})$. Moreover, the projection π commutes with the diagonal maps of P and $\pi(P)$, that is $\Delta_{(\pi(P), \pi(\vec{v}))} = (\pi \times \pi)\Delta_{(P, \vec{v})}$.*

Proof. Since P is a generalized permutahedron, the direction of the edges of the intersection $P \cap \rho_z P$, for any $z \in P$, are vectors with coordinates equal to 0, 1 or -1 , and the same number of 1 and -1 (combine Proposition 1.27 and Proposition 3.4 of [Lap22]). The direction \vec{d} of such an edge satisfies $\langle \vec{d}, \vec{v} \rangle \neq 0$, since the first non-zero coordinate of \vec{d} will contribute a greater amount than the sum of the remaining coordinates in the scalar product. For the same reason, we have $\langle \pi(\vec{d}), \pi(\vec{v}) \rangle \neq 0$. As $\pi(P \cap \rho_z P) = \pi(P) \cap \rho_{\pi(z)} \pi(P)$, we have in particular that the image of the edges of $P \cap \rho_z P$ under π are the edges of $\pi(P) \cap \rho_{\pi(z)} \pi(P)$ and thus that $\pi(P)$ is positively oriented by $\pi(\vec{v})$. For the last part of the statement, observe that π preserves the orientation of the edges: if we have $\langle \vec{d}, \vec{v} \rangle > 0$, then we have $\langle \pi(\vec{d}), \pi(\vec{v}) \rangle > 0$. Hence, the image of the vertex $\text{top}_{\vec{v}}(P \cap \rho_z P)$, which maximizes $\langle -, \vec{v} \rangle$ over $P \cap \rho_z P$, under π is equal to the vertex $\text{top}_{\pi(\vec{v})}(\pi(P) \cap \rho_{\pi(z)} \pi(P))$ which maximizes $\langle -, \pi(\vec{v}) \rangle$ over $\pi(P) \cap \rho_{\pi(z)} \pi(P)$. The argument for the minimum $\text{bot}(P \cap \rho_z P)$ is the same. \square

Proposition 2.8. *Let $P \subset \mathbb{R}^{n+1}$ be a generalized permutahedron. Any two good orientation vectors \vec{v}, \vec{w} define the same diagonal maps on P and $\pi(P)$, that is, we have $\Delta_{(P, \vec{v})} = \Delta_{(P, \vec{w})}$ and $\Delta_{(\pi(P), \pi(\vec{v}))} = \Delta_{(\pi(P), \pi(\vec{w}))}$.*

Proof. Good orientation vectors are principal orientation vectors [Lap22, Definition 3.15]. Since all principal orientation vectors live in the same chamber of the fundamental hyperplane arrangement of the permutahedron, they all define the same diagonal on the permutahedron [Lap22, Proposition 1.21], and thus the same diagonal on any generalized permutahedron (Proposition 2.5). So, we have $\Delta_{(P, \vec{v})} = \Delta_{(P, \vec{w})}$. Finally, using Proposition 2.7, we have $\Delta_{(\pi(P), \pi(\vec{v}))} = (\pi \times \pi)\Delta_{(P, \vec{v})} = (\pi \times \pi)\Delta_{(P, \vec{w})} = \Delta_{(\pi(P), \pi(\vec{w}))}$. \square

2.4. Diagonal of the Forcey–Loday multiplihedra.

Definition 2.9. *A well-oriented realization of the multiplihedron is a positively oriented polytope which realizes the multiplihedron and such that the orientation vector induces the Tamari-type order on the set of vertices.*

Proposition 2.10. *Any good orientation vector induces a well-oriented realization (J_ω, \vec{v}) of the Forcey–Loday multiplihedron, for any weight ω .*

Proof. Using Definition 1.9, we can compute that any edge of the realization of the multiplihedron J_ω is directed, according to the Tamari type order, by either e_i or $e_i - e_j$, for $i < j$. Since \vec{v} has strictly decreasing coordinates, the scalar product is in each case positive. It remains to show that $P \cap \rho_z P$ is oriented by \vec{v} , for any $z \in P$. This follows directly from Proposition 2.7, and the fact that J_ω arises as the projection under π of a generalized permutahedron as shown in Proposition 1.16. \square

Any good orientation vector therefore defines a diagonal map $\Delta_\omega : J_\omega \rightarrow J_\omega \times J_\omega$, for any weight ω . These diagonal maps are all equivalent up to isomorphism in the category Poly.

Proposition 2.11. *For any pair of weights ω and θ of length n , there exists a unique isomorphism $\text{tr} = \text{tr}_\omega^\theta : J_\omega \rightarrow J_\theta$ in the category Poly, which preserves homeomorphically the faces of the same type and which commutes with the respective diagonals.*

Proof. The arguments of [MTTV21, Sections 3.1-3.2] hold in the present case using Proposition 1.10. We note that the crucial condition above is that the map tr commutes with the respective diagonals: this makes the map tr unique and highly non-trivial to construct, see the proof of [MTTV21, Proposition 7]. \square

Definition 2.12. *We define $\Delta_n : J_n \rightarrow J_n \times J_n$ to be the diagonal induced by any good orientation vector for the Forcey–Loday realization of standard weight $\omega = (1, \dots, 1)$.*

2.5. Operadic bimodule structure on the Forcey–Loday multiplihedra. We will use the transition maps tr of Proposition 2.11 above to endow the family of standard weight Forcey–Loday multiplihedra with an operadic bimodule structure over the standard weight Loday associahedra. The uniqueness property of the map tr will be used in a crucial way.

Definition 2.13 (Action-composition maps). *For any $n, m \geq 1$ and any $1 \leq i \leq m$, for any $k \geq 2$ and any $i_1, \dots, i_k \geq 1$, we define the action-composition maps by*

$$\begin{aligned} \circ_{p+1} : J_{p+1+r} \times K_q &\xrightarrow{\text{tr} \times \text{id}} J_{(1, \dots, q, \dots, 1)} \times K_q \xleftarrow{\Theta_{p,q,r}} J_n \text{ and} \\ \gamma_{i_1, \dots, i_k} : K_k \times J_{i_1} \times \dots \times J_{i_k} &\xrightarrow{\text{tr} \times \text{id}} K_{(i_1, \dots, i_k)} \times J_{i_1} \times \dots \times J_{i_k} \xleftarrow{\Theta^{i_1, \dots, i_k}} J_{i_1 + \dots + i_k}, \end{aligned}$$

where the last inclusions are given by the block permutations of the coordinates introduced in the proof of Proposition 1.10.

Recall from [MTTV21, Theorem 1] that the diagonal maps $\Delta_n : K_n \rightarrow K_n \times K_n$ define a morphism of operads, where the operad $\{K_n \times K_n\}$ is to be understood as the Hadamard product $\{K_n\} \times \{K_n\}$. The next proposition shows that the diagonal maps $\Delta_n : K_n \rightarrow K_n \times K_n$ and $\Delta_n : J_n \rightarrow J_n \times J_n$ are compatible with the action-composition maps introduced in Definition 2.13.

Proposition 2.14. *The diagonal maps Δ_n commute with the maps Θ .*

Proof. First observe that a good orientation vector has decreasing coordinates, thereby induces the diagonal maps $\Delta_n : K_n \rightarrow K_n \times K_n$ and the operad structure on $\{K_n\}$ defined in [MTTV21]. Following [Lap22, Proposition 4.14], to prove the claim it suffices to show that the preimage

under Θ^{-1} of a good orientation vector is still a good orientation vector for each associahedron and multiplihedron. This is easily seen to be the case from the definition of Θ , in the proof of Proposition 1.10. \square

Theorem 1.

- (1) The collection $\{J_n\}_{n \geq 1}$ together with the action-composition maps \circ_i and γ_{i_1, \dots, i_k} form an operadic bimodule over the operad $\{K_n\}$ in the category Poly.
- (2) The maps $\{\Delta_n : J_n \rightarrow J_n \times J_n\}_{n \geq 1}$ form a morphism of $(\{K_n\}, \{K_n\})$ -operadic bimodules in the category Poly.

Proof. Using Proposition 2.14, we can apply the proof of [MTTV21, Theorem 1] *mutatis mutandis*. The uniqueness of the transition map tr is the key argument, as it forces the operadic axioms to hold. We also point out that $\{J_n \times J_n\}$ is to be understood as the Hadamard product $\{J_n\} \times \{J_n\}$, and that its $(\{K_n\}, \{K_n\})$ -operadic bimodule structure is defined as the pullback of its natural $(\{K_n \times K_n\}, \{K_n \times K_n\})$ -operadic bimodule structure under the diagonal maps $\{\Delta_n : K_n \rightarrow K_n \times K_n\}$. \square

Point (1) of Theorem 1 was already mentioned in [Maz21a, Section 1.2], where associahedra and multiplihedra are realized as compactifications of moduli spaces of metric trees and used to construct A_∞ -structures on the Morse cochains of a closed manifold.

3. CELLULAR FORMULA FOR THE DIAGONAL OF THE MULTIPLIHEDRA

We compute in Theorem 2 an explicit cellular formula for the diagonal of the Forcey–Loday multiplihedra, using again the key fact that the Ardila–Doker multiplihedron is a generalized permutahedron to which one can apply Proposition 2.5 and the results of [Lap22]. We then explain geometrically why this formula necessarily has to differ from the "magical formula" computed for the associahedra in [MTTV21].

3.1. 2-colored nested linear graphs. Let ℓ be a *linear graph* with n vertices, as represented in Figure 6. We respectively write $V(\ell)$ and $E(\ell)$ for its sets of vertices and edges. Any subset of edges $N \subset E(\ell)$ defines a subgraph of ℓ whose edges are N and whose vertices are all the vertices adjacent to an edge in N . We call this graph the *closure* of N .

Definition 3.1 (Nest and nesting).

- A nest of a linear graph ℓ with n vertices is a non-empty set of edges $N \subset E(\ell)$ whose closure is a connected subgraph of ℓ .
- A nesting of a linear graph ℓ is a set $\mathcal{N} = \{N_i\}_{i \in I}$ of nests such that
 - (1) the trivial nest $E(\ell)$ is in \mathcal{N} ,
 - (2) for every pair of nests $N_i \neq N_j$, we have either $N_i \subsetneq N_j$, $N_j \subsetneq N_i$ or $N_i \cap N_j = \emptyset$, and
 - (3) if $N_i \cap N_j = \emptyset$ then no edge of N_i is adjacent to an edge of N_j .

Two nests that satisfy Conditions (2) and (3) are said to be *compatible*. We denote the set of nestings of ℓ by $\mathcal{N}(\ell)$. We naturally represent a nesting by circling the closure of each nest as in Figure 6. A nesting is moreover *maximal* if it has maximal cardinality $|\mathcal{N}| = |E(\ell)|$.

Definition 3.2 (2-colored nesting). A 2-colored nesting is a nesting where each nest is either colored in blue, red or both red and blue (that is, purple), and which satisfy the following properties:

- (1) if a nest N is blue or purple, then all nests contained in N are blue, and

(2) if a nest N is red or purple, then all nests that contain N are red.

We call *monochrome* the nests that are either blue or red, and *bicolored* the purple nests. We denote by $\text{mono}(\mathcal{N})$ the set of monochrome nests of a 2-colored nesting \mathcal{N} , and by $\mathcal{N}_2(\ell)$ the set of 2-colored nestings of ℓ . A 2-colored nesting is moreover *maximal* if it has maximal cardinality, and it is made of monochrome nests only.

REMARK 3.3. The data of a 2-colored nesting on a graph is equivalent to the data of a marked tubing on its line graph, as defined in [DF08]. See also [Lap22, Remark 2.4].

Lemma 3.4. *There is a bijection between (2-colored) trees with n leaves and (2-colored) nested linear graphs with n vertices. Under this map, (2-colored) maximal trees are in bijection with maximal (2-colored) nested linear graphs.*

Under this bijection, vertices of 2-colored trees correspond to nests, and their colors agree under the previous conventions.




FIGURE 6. Bijections between 2-colored trees, 2-colored nested linear graphs, and 2-colored parenthesizations.

3.2. Cellular formula for the diagonal.

Definition 3.5. *Let (ℓ, \mathcal{N}) be a nested linear graph. We respectively denote by $B(\mathcal{N})$, $P(\mathcal{N})$ and $R(\mathcal{N})$ the set of blue, purple and red nests of \mathcal{N} . We define $Q(\mathcal{N})$ to be the set whose elements are the unions of nests*

$$\bigcup_{i=1}^k R_i \cup \bigcup_{B \in B(\mathcal{N})} B \cup \bigcup_{P \in P(\mathcal{N})} P$$

where $R_1, \dots, R_k \in R(\mathcal{N})$, the case $\bigcup R_i = \emptyset$ being allowed, and where two unions that result in the same set are identified.

We number the edges of the linear graph with n vertices from bottom to top as represented in Figure 6, starting at 1 and ending at $n - 1$. To each blue nest $B \in B(\mathcal{N})$ in a 2-colored nesting \mathcal{N} of a linear graph with n vertices, we associate the *characteristic vector* $\vec{B} \in \mathbb{R}^n$ which has a 1 in position i if $i \in B$, 0 in position i if $i \notin B$ and 0 in position n . To each union of nests $Q \in Q(\mathcal{N})$, we associate the characteristic vector $\vec{Q} \in \mathbb{R}^n$ which has a 1 in position i if $i \in Q$, 0 in position i if $i \notin Q$ and 1 in position n . We denote moreover by \vec{n} the vector $(1, \dots, 1) \in \mathbb{R}^n$.

Lemma 3.6. *The normal cone of the face of the Ardila–Doker realization of the multiplihedron labeled by the 2-colored nesting \mathcal{N} is given by*

$$\text{Cone} \left(\{-\vec{B}\}_{B \in B(\mathcal{N})} \cup \{-\vec{Q}\}_{Q \in Q(\mathcal{N})} \cup \{\vec{n}, -\vec{n}\} \right).$$

Proof. This follows from the description of the Ardila–Doker multiplihedron as a generalized permutahedron: the normal cone of a face of the multiplihedron is a union of normal cones of faces of the permutahedron, and these faces can be easily determined from the projection from the permutahedron to the multiplihedron, written down explicitly in the proof of [Dok11, Theorem 3.3.6]. \square

We are now ready to compute the cellular formula for the diagonal of the Forcey–Loday multiplihedra. We introduce

$$D(n) := \{(I, J) \mid I, J \subset \{1, \dots, n\}, |I| = |J|, I \cap J = \emptyset, \min(I \cup J) \in I\}.$$

We number again the edges of the linear graph with n vertices from bottom to top, starting at 1 and ending at $n - 1$. Blue nests and unions of blue, purple and red nests can then in particular be seen as subsets of $\{1, \dots, n - 1\}$, hence of $\{1, \dots, n\}$.

Theorem 2. *The cellular image of the diagonal map $\Delta_n : \mathbb{J}_n \rightarrow \mathbb{J}_n \times \mathbb{J}_n$ introduced in Definition 2.12 admits the following description. For \mathcal{N} and \mathcal{N}' two 2-colored nestings of the linear graph with n vertices, we have that*

$$\begin{aligned} (\mathcal{N}, \mathcal{N}') \in \text{Im } \Delta_n &\iff \forall (I, J) \in D(n), \\ &\quad \exists B \in B(\mathcal{N}), |B \cap I| > |B \cap J| \text{ or} \\ &\quad \exists Q \in Q(\mathcal{N}), |(Q \cup \{n\}) \cap I| > |(Q \cup \{n\}) \cap J| \text{ or} \\ &\quad \exists B' \in B(\mathcal{N}'), |B' \cap I| < |B' \cap J| \text{ or} \\ &\quad \exists Q' \in Q(\mathcal{N}'), |(Q' \cup \{n\}) \cap I| < |(Q' \cup \{n\}) \cap J|. \end{aligned}$$

Proof. The essential ingredient is the computation of the fundamental hyperplane arrangement of the permutahedron, which was done in [Lap22, Section 3.1]. The result follows in three steps:

- (1) Since a good orientation vector \vec{v} is also a principal orientation vector [Lap22, Definition 3.15], it orients positively the permutahedron.
- (2) Using Proposition 2.5 and the description of the normal cones of the faces of the multiplihedron in Lemma 3.6, we get the above formula for the Ardila–Doker realizations of the multiplihedra.
- (3) Proposition 2.7 guarantees that this formula holds for the Forcey–Loday realizations, which completes the proof. \square

We now make this formula explicit in dimension 1, 2 and 3. We write 2-colored nestings of a linear graph with n vertices as 2-colored parenthesizations of a word with n symbols \bullet , which are easier to read and shorter to type, see Figure 6. We moreover only write pairs of faces (F, G) such that $\dim F + \dim G = \dim P$.

$$\Delta_2((\bullet\bullet)) = (\bullet\bullet) \times (\bullet\bullet) \cup (\bullet\bullet) \times (\bullet\bullet)$$

$$\begin{aligned}\Delta_3((\bullet\bullet\bullet)) &= ((\bullet\bullet)\bullet) \times (\bullet\bullet\bullet) \cup (\bullet\bullet\bullet) \times (\bullet(\bullet\bullet)) \cup (\bullet\bullet\bullet) \times (\bullet(\bullet\bullet)) \\ &\cup (\bullet\bullet\bullet) \times (\bullet(\bullet\bullet)) \cup (\bullet(\bullet\bullet)) \times (\bullet(\bullet\bullet)) \cup ((\bullet\bullet)\bullet) \times ((\bullet\bullet)\bullet) \\ &\cup ((\bullet\bullet)\bullet) \times (\bullet\bullet\bullet) \cup ((\bullet\bullet)\bullet) \times (\bullet\bullet\bullet)\end{aligned}$$

$$\begin{aligned}\Delta_4((\bullet\bullet\bullet\bullet)) &= \\ &(((\bullet\bullet)\bullet)\bullet) \times (\bullet\bullet\bullet\bullet) \cup (\bullet\bullet\bullet\bullet) \times (\bullet(\bullet(\bullet\bullet))) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \\ \cup &((\bullet\bullet)(\bullet\bullet)) \times (\bullet\bullet(\bullet\bullet)) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup ((\bullet\bullet)\bullet\bullet) \times (\bullet\bullet(\bullet\bullet)) \\ \cup &(\bullet(\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \cup ((\bullet\bullet)\bullet\bullet) \times (\bullet\bullet(\bullet\bullet)) \\ \cup &((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup (((\bullet\bullet)\bullet)\bullet) \times ((\bullet\bullet\bullet)\bullet) \cup (\bullet\bullet(\bullet\bullet)) \times (\bullet(\bullet(\bullet\bullet))) \\ \cup &((\bullet\bullet)(\bullet\bullet)) \times ((\bullet\bullet)(\bullet\bullet)) \cup (\bullet(\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \cup ((\bullet\bullet)\bullet\bullet) \times ((\bullet\bullet)(\bullet\bullet)) \\ \cup &(\bullet(\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup (\bullet((\bullet\bullet)\bullet)) \times (\bullet(\bullet\bullet\bullet)) \cup ((\bullet\bullet)\bullet\bullet) \times ((\bullet\bullet)(\bullet\bullet)) \\ \cup &(\bullet(\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \cup ((\bullet\bullet)\bullet\bullet) \times (\bullet\bullet(\bullet\bullet)) \\ \cup &(((\bullet\bullet)\bullet)\bullet) \times ((\bullet\bullet)\bullet\bullet) \cup (\bullet(\bullet\bullet\bullet)) \times (\bullet(\bullet(\bullet\bullet))) \cup (((\bullet\bullet)\bullet)\bullet) \times ((\bullet\bullet)\bullet\bullet) \\ \cup &(\bullet(\bullet\bullet\bullet)) \times (\bullet(\bullet(\bullet\bullet))) \cup (((\bullet\bullet)\bullet)\bullet) \times (\bullet\bullet\bullet\bullet) \cup (\bullet\bullet\bullet\bullet) \times (\bullet(\bullet(\bullet\bullet))) \\ \cup &(((\bullet\bullet)\bullet)\bullet) \times (\bullet\bullet\bullet\bullet) \cup (\bullet\bullet\bullet\bullet) \times (\bullet(\bullet(\bullet\bullet))) \cup ((\bullet\bullet)(\bullet\bullet)) \times (\bullet\bullet(\bullet\bullet)) \\ \cup &((\bullet\bullet\bullet)\bullet) \times ((\bullet(\bullet\bullet)\bullet)) \cup ((\bullet\bullet)(\bullet\bullet)) \times (\bullet\bullet(\bullet\bullet)) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \\ \cup &((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \cup ((\bullet\bullet)\bullet\bullet) \times (\bullet\bullet(\bullet\bullet)) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet\bullet)) \\ \cup &((\bullet(\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \cup ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \cup (((\bullet\bullet)\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet)\end{aligned}$$

We also compute in Figure 7 the number of faces of complementary dimensions and the number of pairs of vertices in the cellular image of the diagonal of the multiplihedra in dimensions 0 to 6. They are compared with the diagonals induced by the same orientation vector on the Loday associahedra and the permutahedra. The two sequences of numbers that we obtain did not appear before in [OEI22].

Pairs $(F, G) \in \text{Im } \Delta_{(P, \vec{v})}$	Polytopes	0	1	2	3	4	5	6	[OEI22]
$\dim F + \dim G = \dim P$	Associahedra	1	2	6	22	91	408	1938	A000139
	Multiplihedra	1	2	8	42	254	1678	11790	to appear
	Permutahedra	1	2	8	50	432	4802	65536	A007334
$\dim F = \dim G = 0$	Associahedra	1	3	13	68	399	2530	16965	A000260
	Multiplihedra	1	3	17	122	992	8721	80920	to appear
	Permutahedra	1	3	17	149	1809	28399	550297	A213507

FIGURE 7. Number of pairs of faces in the cellular image of the diagonal of the associahedra, multiplihedra and permutahedra of dimension $0 \leq \dim P \leq 6$, induced by any good orientation vector.

3.3. About the cellular formula. Given a face F of a positively oriented polytope (P, \vec{v}) , the orientation vector \vec{v} defines a unique vertex $\text{top } F$ (resp. $\text{bot } F$) which maximizes (resp. minimizes) the scalar product $\langle -, \vec{v} \rangle$ over F . By [Lap22, Proposition 1.15], any pair of faces $(F, G) \in \text{Im } \Delta_{(P, \vec{v})}$ satisfies $\text{top } F \leq \text{bot } G$. In the case of the simplices, the cubes and the associahedra, the converse also holds: the image of the diagonal is given by the "magical formula"

$$(1) \quad (F, G) \in \text{Im } \Delta_n \iff \text{top } F \leq \text{bot } G .$$

This formula, however, does not hold for the diagonal of the Forcey–Loday multiplihedra.

Proposition 3.7. *The diagonal on the multiplihedron J_4 is such that*

$$\text{Im } \Delta_4 \subseteq \{(F, G), \text{top } F \leq \text{bot } G\} .$$

Proof. The pairs of faces (F, G) that satisfy $\dim F + \dim G = 3$ and $\text{top } F \leq \text{bot } G$ include the four pairs

$$(2) \quad \begin{array}{cc} ((\bullet\bullet\bullet)\bullet) \times ((\bullet(\bullet\bullet))\bullet) & ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \\ ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) & ((\bullet(\bullet\bullet))\bullet) \times (\bullet(\bullet\bullet)\bullet) \end{array}$$

and the four pairs

$$(3) \quad \begin{array}{cc} ((\bullet\bullet\bullet)\bullet) \times ((\bullet(\bullet\bullet))\bullet) & ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) \\ ((\bullet\bullet\bullet)\bullet) \times (\bullet(\bullet\bullet)\bullet) & ((\bullet(\bullet\bullet))\bullet) \times (\bullet(\bullet\bullet)\bullet) \end{array} .$$

While the image $\text{Im } \Delta_4$ contains the four pairs in (2), it does *not* include the four pairs in (3), as can be checked directly from Theorem 2. \square

REMARK 3.8. We point out that Formula (1) also does not hold neither for the permutahedra nor the operahedra in general, as proven in [Lap22, Section 3.2].

The diagonal Δ_n being a section of the projection $\pi : J_n \times J_n \rightarrow J_n, (x, y) \mapsto (x + y)/2$ [Lap22, Proposition 1.1], one can in fact represent its cellular image by projecting it to J_n : for each pair of faces $(F, G) \in \text{Im } \Delta_n$, one draws the polytope $(F + G)/2$ in J_n . This defines a polytopal subdivision of J_n . The polytopal subdivision of J_3 can be found in [Lap22, Figure 3], while the polytopal subdivision of J_4 is illustrated on the first page of this article.

Proposition 3.7 can then be illustrated geometrically as follows. There are two distinct diagonals on J_4 which agree with the Tamari-type order on the vertices. The first one, corresponding to the diagonal defined in this paper, is induced by the choice of any orientation vector $\vec{v} = (v_1, v_2, v_3, v_4)$ satisfying $v_1 > v_2 > v_3 > v_4$ and $v_1 + v_4 > v_2 + v_3$ (here we work with the Ardila–Doker realization of the multiplihedron). Changing the last condition to $v_1 + v_4 < v_2 + v_3$ gives the second choice of diagonal, which is in fact exactly the diagonal of Saneblidze–Umble [SU04, Section 5]. These two diagonals on J_4 then differ by four pairs of faces, as represented in Figure 8: the first diagonal includes the pairs of (2), while the second diagonal includes the pairs of (3). Under the projection $\pi : J_4 \times J_4 \rightarrow J_4, (x, y) \mapsto (x + y)/2$, these two families of faces induce two distinct polytopal subdivisions of the same "diamond" inside J_4 , represented in Figure 9. We also refer to the last paragraph of Section 4.2.3 for an algebraic counterpart of Proposition 3.7.

REMARK 3.9. The two previous families of orientation vectors correspond to two adjacent chambers in the fundamental hyperplane arrangement of the permutahedron [Lap22, Theorem 3.6], separated by the hyperplane $x_1 + x_4 = x_2 + x_3$, pictured in blue in [Lap22, Figure 12]. A way to relate the diagonal constructed in this article to the diagonal of [SU04, Section 5] would possibly be to find further choices of chambers in the fundamental hyperplane arrangements of the permutahedra (or the multiplihedra) in all dimensions $n \geq 4$ recovering the latter diagonal, see also [Lap22, Remark 3.18].




FIGURE 8. The four pairs of (2) represented in blue on the two top copies of J_4 and the four pairs of (3) represented in red on the two bottom copies of J_4 . The minimal (top right) and maximal (bottom left) vertices for the Tamari-type order are drawn in black, in the top left copy.

4. TENSOR PRODUCT OF A_∞ -MORPHISMS AND A_∞ -FUNCTORS

We begin by proving that for a certain choice of cellular orientation, the cellular chains functor maps the Loday associahedra to the operad A_∞ encoding A_∞ -algebras and the Forcey–Loday multiplihedra to the operadic bimodule M_∞ encoding A_∞ -morphisms between them. It then maps the respective geometric diagonals to algebraic ones, which can be used to define compatible tensor products of A_∞ -algebras and A_∞ -morphisms (with signs). Tensor product of A_∞ -categories and A_∞ -functors are defined in a similar fashion, and we relate them to the different notions of A_∞ -categories with identities. We finally study coassociativity, cocommutativity and compatibility with composition of A_∞ -morphisms for these diagonals.




FIGURE 9. The two distinct subdivisions of the same "diamond" in J_4 , respectively induced by the pairs of (2) and (3).

We show that these properties are always satisfied up to homotopy, hinting at the idea that the category $\infty\text{-A}_\infty\text{-alg}$ should possess some kind of *homotopy* symmetric monoidal structure.

4.1. A_∞ -algebras and A_∞ -morphisms.

4.1.1. *Definitions.* We work in the rest of this article with homological convention. We will refer to chain complexes as *dg modules*, where the abbreviation dg stands for "differential graded", and their differential will always have degree -1 .

Definition 4.1 (A_∞ -algebra). *An A_∞ -algebra is the data of a dg module (A, ∂) together with operations*

$$m_n : A^{\otimes n} \rightarrow A, \quad n \geq 2$$

of degree $|m_n| = n - 2$, satisfying the equations

$$[\partial, m_n] = - \sum_{\substack{p+q+r=n \\ 2 \leq q \leq n-1}} (-1)^{p+qr} m_{p+1+r} (\text{id}^{\otimes p} \otimes m_q \otimes \text{id}^{\otimes r}), \quad n \geq 2.$$

Definition 4.2 (A_∞ -morphism). *An A_∞ -morphism $F : A \rightsquigarrow B$ between two A_∞ -algebras $(A, \{m_n\})$ and $(B, \{m'_n\})$ is a family of linear maps*

$$f_n : A^{\otimes n} \rightarrow B, \quad n \geq 1$$

of degree $|f_n| = n - 1$, satisfying the equations

$$[\partial, f_n] = \sum_{\substack{p+q+r=n \\ q \geq 2}} (-1)^{p+qr} f_{p+1+r} (\text{id}^{\otimes p} \otimes m_q \otimes \text{id}^{\otimes r}) - \sum_{\substack{i_1+\dots+i_k=n \\ k \geq 2}} (-1)^\varepsilon m'_k (f_{i_1} \otimes \dots \otimes f_{i_k}), \quad n \geq 1,$$

where $\varepsilon = \sum_{u=1}^k (k-u)(1-i_u)$.

For three A_∞ -algebras A, B, C and two A_∞ -morphisms $F : A \rightsquigarrow B, B \rightsquigarrow C$, their composition $G \circ F : A \rightsquigarrow C$ is the A_∞ -morphism whose operation of arity n is given by the formula

$$(G \circ F)_n := \sum_{i_1 + \dots + i_k = n} (-1)^\varepsilon g_k(f_{i_1} \otimes \dots \otimes f_{i_k}) .$$

This composition is associative. We moreover point out that a standard *dg (associative) algebra* can be defined as an A_∞ -algebra whose higher operations m_n vanish for $n \geq 3$. For more details on these notions, we refer to [LV12, Chapter 9].

Definition 4.3. *We denote by $\infty\text{-}A_\infty\text{-alg}$ the category of A_∞ -algebras with A_∞ -morphisms.*

Representing the operations m_n as corollae $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$ of arity n , the equations of Definition 4.1 read as

$$(4) \quad [\partial, \begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}] = - \sum_{\substack{p+q+r=n \\ 2 \leq q \leq n-1}} (-1)^{p+qr} \begin{smallmatrix} & q \\ \diagup & \diagdown \\ \diagdown & \diagup \\ & p \quad r \end{smallmatrix} .$$

Representing the operations m_n in blue $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$, the operations m'_n in red $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$ and the operations f_n by $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$, the equations of Definition 4.2 can be rewritten as

$$(5) \quad [\partial, \begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}] = \sum_{\substack{p+q+r=n \\ q \geq 2}} (-1)^{p+qr} \begin{smallmatrix} & q \\ \diagup & \diagdown \\ \diagdown & \diagup \\ & p \quad r \end{smallmatrix} - \sum_{\substack{i_1 + \dots + i_k = n \\ k \geq 2}} (-1)^\varepsilon \begin{smallmatrix} i_1 & & i_k \\ \diagup & \dots & \diagdown \\ \diagdown & \dots & \diagup \end{smallmatrix} .$$

Finally, representing the operations f_n by $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$ and the operations g_n by $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$, the formula for the composition of A_∞ -morphisms reads as

$$(6) \quad \sum_{i_1 + \dots + i_k = n} (-1)^\varepsilon \begin{smallmatrix} i_1 & & i_k \\ \diagup & \dots & \diagdown \\ \diagdown & \dots & \diagup \end{smallmatrix} .$$

4.1.2. The operad A_∞ and the operadic bimodule M_∞ .

Definition 4.4 (Operad A_∞). *The operad A_∞ is the quasi-free dg operad generated in arity $n \geq 2$ by one operation $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$ of degree $n - 2$*

$$A_\infty := (\mathcal{T}(\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}, \begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}, \dots), \partial) ,$$

and whose differential is defined by Equations (4).

Definition 4.5 (Operadic bimodule M_∞). *The operadic bimodule M_∞ is the quasi-free (A_∞, A_∞) -operadic bimodule generated in arity $n \geq 1$ by one operation $\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}$ of degree $n - 1$*

$$M_\infty := (\mathcal{T}^{A_\infty, A_\infty}(\begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}, \begin{smallmatrix} \diagup & \diagdown \\ \diagdown & \diagup \end{smallmatrix}, \dots), \partial) ,$$

and whose differential is defined by Equations (5).

We denote by End_A the *endomorphism operad* of a dg module A , i.e. the operad whose dg module of operations of arity n is $\text{End}_A(n) := \text{Hom}(A^{\otimes n}, A)$. An A_∞ -algebra structure on A is then equivalent to the datum of a morphism of operads $A_\infty \rightarrow \text{End}_A$. We denote similarly by Hom_B^A the $(\text{End}_B, \text{End}_A)$ -operadic bimodule defined by $\text{Hom}_B^A(n) := \text{Hom}(A^{\otimes n}, B)$. An A_∞ -morphism between two A_∞ -algebras A and B is then equivalent to the datum of a morphism of operadic bimodules $M_\infty \rightarrow \text{Hom}_B^A$.

Composition of A_∞ -morphisms can also be formulated at the level of the operadic bimodule M_∞ as a morphism of (A_∞, A_∞) -operadic bimodules $M_\infty \rightarrow M_\infty \circ_{A_\infty} M_\infty$, where the notation \circ_{A_∞} denotes the *relative composite product* [LV12, Section 11.2.1]. We write the first factor of $M_\infty \circ_{A_\infty} M_\infty$ using green for the color above the gauge and red for the color below the gauge,

$$M_\infty := \mathcal{T}^{A_\infty, A_\infty}(\text{green}, \text{green}, \text{green}, \text{green}, \dots),$$

and its second factor using blue for the color above the gauge and green for the color below the gauge

$$M_\infty := \mathcal{T}^{A_\infty, A_\infty}(\text{blue}, \text{blue}, \text{blue}, \text{blue}, \dots).$$

Definition 4.6 (Composition morphism). *The composition morphism is defined to be the morphism of (A_∞, A_∞) -operadic bimodules $\text{comp} : M_\infty \rightarrow M_\infty \circ_{A_\infty} M_\infty$ given on the generating operations of M_∞ by*

$$\text{comp}(\text{blue}) = \sum_{i_1 + \dots + i_k = n} (-1)^\varepsilon \text{diagram}.$$

The composition of two A_∞ -morphisms $A \rightsquigarrow B$ and $B \rightsquigarrow C$ is then equivalent to the following composition of morphisms of operadic bimodules

$$M_\infty \xrightarrow{\text{comp}} M_\infty \circ_{A_\infty} M_\infty \longrightarrow \text{Hom}_C^B \circ_{\text{End}_B} \text{Hom}_B^A \longrightarrow \text{Hom}_C^A.$$

4.1.3. *The Forcey–Loday multiplihedra realize the operadic bimodule M_∞ .*

Definition 4.7 (Cellular orientation). *Let $P \subset \mathbb{R}^n$ be a polytope, and let F be a face of P . A cellular orientation of F is a choice of orientation of its linear span. A cellular orientation of P is a choice of cellular orientation for each face F of P .*

We respectively denote by CW and dg-mod the symmetric monoidal categories of CW complexes and of dg modules over \mathbb{Z} , and by $C_\bullet^{\text{cell}} : \text{CW} \rightarrow \text{dg-mod}$ the cellular chains functor. A choice of a cellular orientation for every polytope $P \in \text{Poly}$ defines an inclusion $\text{Poly} \subset \text{CW}$. Then, the strong symmetric monoidal functor C_\bullet^{cell} respectively sends operads and operadic bimodules in polytopes to dg operads and dg operadic bimodules.

Definition 4.8 (Left-levelwise order). *Let t be a (2-colored) tree t . The left-levelwise order on the vertices of t is defined by ordering them from bottom to top and from left to right, proceeding one level at a time.*

Given a tree t , there is a unique decomposition $t = (\dots((c_{n_1} \circ_{i_1} c_{n_2}) \circ_{i_2} c_{n_3}) \dots \circ_{i_k} c_{n_{k+1}})$ where the corollae c_n are grafted according to this total order. Using the grafting operations defined in Section 1.1.2, a 2-colored tree admits similarly a unique decomposition as a sequence of blue corollae, red corollae and 2-colored corollae ordered according to this total order. We



FIGURE 10. The tree on the left decomposes as $(c_4 \circ_3 c_4) \circ_3 c_3$ and the orientation on the face it labels is determined by the product $K_4 \times K_4 \times K_3$. The tree on the right decomposes as $(c_4 \circ_1 c_3) \circ_6 c_4$ and defines the orientation determined by the product $K_4 \times K_3 \times K_4$.

can then make the same choices of cellular orientations as in [Maz21a, Section 1.4], illustrated in Figure 10 :

- For the Loday associahedra $K_n \subset \mathbb{R}^{n-1}$ of [MTTV21], we choose the basis $\{e_1 - e_{j+1}\}_{1 \leq j \leq n-2}$ as positively oriented basis of the top dimensional cell \searrow . We then choose the orientation of any other face t of K_n to be the image of the positively oriented bases of the top cells of the polytopes K_{n_i} under the sequence of partial compositions following the left-levelwise order on t .
- We choose the basis $\{-e_j\}_{1 \leq j \leq n-1}$ as positively oriented basis of the top dimensional cell \swarrow of the Forcey–Loday multiplihedra $J_n \subset \mathbb{R}^{n-1}$. We then choose the orientation of any other face t of J_n to be the image of the positively oriented bases of the top cells of the polytopes K_{n_i} and J_{n_j} under the sequence of action-compositions maps, following the left-levelwise order on t .

Proposition 4.9. *These cellular orientations on the Loday associahedra and the Forcey–Loday multiplihedra provide an isomorphism of dg operads $C_\bullet^{\text{cell}}(\{K_n\}) \cong A_\infty$ and an isomorphism of dg operadic bimodules $C_\bullet^{\text{cell}}(\{J_n\}) \cong M_\infty$.*

Proof. The choice of a cellular orientation endows the K_n and J_n with a natural CW structure (see [Lap22, Proposition 4.22]). The choice of the left-levelwise order on trees ensures that we recover precisely the usual sign conventions for the partial compositions of the quasi-free operad A_∞ and for the action-composition maps of the quasi-free operadic bimodule M_∞ . The signs for the respective differentials were computed in [Maz21a, Section 1.4]. \square

4.2. Tensor product of A_∞ -algebras and A_∞ -morphisms.

4.2.1. Diagonals on the operad A_∞ and on the operadic bimodule M_∞ .

Definition 4.10 (Operadic diagonals).

- (1) A diagonal on the operad A_∞ is a morphism of dg operads $\Delta : A_\infty \rightarrow A_\infty \otimes A_\infty$ which satisfies $\Delta(\searrow) = \searrow \otimes \searrow$.
- (2) Given a diagonal on the operad A_∞ , a diagonal on the operadic bimodule M_∞ is a morphism of operadic bimodules $\Delta : M_\infty \rightarrow M_\infty \otimes M_\infty$ which satisfies $\Delta(\swarrow) = \swarrow \otimes \swarrow$, and where $M_\infty \otimes M_\infty$ is endowed with its (A_∞, A_∞) -operadic bimodule structure induced by the diagonal on A_∞ .

Diagonals provide an adapted framework to define tensor products of A_∞ -algebras and A_∞ -morphisms. Given a diagonal $A_\infty \rightarrow A_\infty \otimes A_\infty$ and two A_∞ -algebras A and B , one can define

an A_∞ -algebra structure on $A \otimes B$ by considering the following composition

$$A_\infty \longrightarrow A_\infty \otimes A_\infty \longrightarrow \text{End}_A \otimes \text{End}_B \longrightarrow \text{End}_{A \otimes B} .$$

Given similarly a diagonal $M_\infty \rightarrow M_\infty \otimes M_\infty$ and two A_∞ -morphisms $F_1 : A_1 \rightsquigarrow B_1$ and $F_2 : A_2 \rightsquigarrow B_2$, one can define an A_∞ -morphism $F_1 \otimes F_2 : A_1 \otimes A_2 \rightsquigarrow B_1 \otimes B_2$ by the following composition

$$M_\infty \rightarrow M_\infty \otimes M_\infty \rightarrow \text{Hom}_{B_1}^{A_1} \otimes \text{Hom}_{B_2}^{A_2} \rightarrow \text{Hom}_{B_1 \otimes B_2}^{A_1 \otimes A_2} .$$

We moreover point out that the conditions $\Delta(\curlyvee) = \curlyvee \otimes \curlyvee$ and $\Delta(\curlywedge) = \curlywedge \otimes \curlywedge$ respectively imply that these constructions recover the standard tensor product of dg algebras and the standard tensor product of ordinary morphisms between dg algebras.

4.2.2. Admissible edges and permutations. We fix a (2-colored) nested linear graph (ℓ, \mathcal{N}) . We denote by N_i the unique minimal nest of \mathcal{N} with respect to nest inclusion, which contains the edge i .

Definition 4.11 (Admissible edge). *For a nested linear graph (ℓ, \mathcal{N}) , an edge i is admissible with respect to \mathcal{N} if $i \neq \min N_i$. For a 2-colored nested linear graph (ℓ, \mathcal{N}) , an edge i is admissible with respect to \mathcal{N} when N_i is bicolored, or if $i \neq \min N_i$ when N_i is monochrome. We denote the set of admissible edges of \mathcal{N} by $\text{Ad}(\mathcal{N})$.*

Definition 4.12 (Left-levelwise order). *The left-levelwise order on \mathcal{N} is defined by ordering the nests by decreasing order of cardinality, and ordering two nests of the same cardinality according to the increasing order on their minimal elements.*

Under the bijection of Lemma 3.4, the left-levelwise order on the nesting of a nested linear graph is equivalent to the left-levelwise order on the vertices of the corresponding tree t , as defined in Definition 4.8 .

Consider the left-levelwise order $N^1 < N^2 < \dots < N^k$ on the nesting $\mathcal{N} = \{N^j\}_{1 \leq j \leq k}$. We endow the set $\text{Ad}(\mathcal{N})$ with a total order, by ordering the admissible edges of $N_1 \setminus \cup_{2 \leq j \leq k} N_j$ in increasing order, then the admissible edges of $N_2 \setminus \cup_{3 \leq j \leq k} N_j$ in increasing order, and so on. Given two nestings $\mathcal{N}, \mathcal{N}'$ of ℓ , we endow the set $\text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}')$ with the total order given by following the total order on $\text{Ad}(\mathcal{N})$ and then the total order on $\text{Ad}(\mathcal{N}')$. We denote by Δ^K and Δ^J the algebraic diagonals obtained from the polytopal ones by applying the cellular chains functor, see Propositions 4.15 and 4.18 below. The proofs of these two propositions include the proofs of the following two lemmas.

Lemma 4.13. *For a pair of nestings of complementary dimensions $(\mathcal{N}, \mathcal{N}') \in \text{Im } \Delta^K$, the function $\sigma_{\mathcal{N}, \mathcal{N}'} : \text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}') \rightarrow \{1, 2, \dots, |\text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}')|\}$ defined on $i \in \text{Ad}(\mathcal{N})$ by*

$$\sigma_{\mathcal{N}, \mathcal{N}'}(i) = \begin{cases} \min N_i - 1 & \text{if } i \in \text{Ad}(\mathcal{N}) \cap \text{Ad}(\mathcal{N}') \text{ and } 1 \neq \min N_i < \min N'_i \\ i - 1 & \text{otherwise,} \end{cases}$$

and similarly on $i \in \text{Ad}(\mathcal{N}')$ by reversing the roles of \mathcal{N} and \mathcal{N}' , induces a permutation of the set $\{1, 2, \dots, |\text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}')|\}$ that we will still denote by $\sigma_{\mathcal{N}, \mathcal{N}'}$.

Lemma 4.14. *For a pair of 2-colored nestings of complementary dimensions $(\mathcal{N}, \mathcal{N}') \in \text{Im } \Delta^J$, the function $\sigma_{\mathcal{N}, \mathcal{N}'} : \text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}') \rightarrow (1, 2, \dots, |\text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}')|)$ defined on $i \in \text{Ad}(\mathcal{N})$ by*

$$\sigma_{\mathcal{N}, \mathcal{N}'}(i) = \begin{cases} \min N_i & \text{if } i \in \text{Ad}(\mathcal{N}) \cap \text{Ad}(\mathcal{N}'), N_i \text{ is monochrome and } N'_i \text{ is not} \\ \min N_i & \text{if } i \in \text{Ad}(\mathcal{N}) \cap \text{Ad}(\mathcal{N}'), N_i \text{ and } N'_i \text{ are monochrome} \\ & \text{and } \min N_i < \min N'_i, \\ i & \text{otherwise,} \end{cases}$$

and similarly on $i \in \text{Ad}(\mathcal{N}')$ by reversing the roles of \mathcal{N} and \mathcal{N}' , induces a permutation of the set $\{1, 2, \dots, |\text{Ad}(\mathcal{N}) \sqcup \text{Ad}(\mathcal{N}')|\}$ that we will still denote by $\sigma_{\mathcal{N}, \mathcal{N}'}$.

4.2.3. The polytopal diagonals on A_∞ and M_∞ . We use nested linear graphs introduced in Section 3.1 to work with the operad A_∞ and the operadic bimodule M_∞ . The generating operation of arity n of A_∞ corresponds to the trivial nested linear graph with n vertices $(\bullet \cdots \bullet)$, while the generating operation of arity n of M_∞ is represented by the trivial 2-colored nested linear graph with n vertices $(\bullet \cdots \bullet)$.

Proposition 4.15. *The image under the functor C_\bullet^{cell} of the diagonal of the Loday associahedra constructed in [MTTV21] defines a diagonal on the operad A_∞ , that we denote Δ^K . It is determined by the formula*

$$\Delta^K((\bullet \cdots \bullet)) = \sum_{\substack{\mathcal{N}, \mathcal{N}' \in \mathcal{N}_n \\ \text{top}(\mathcal{N}) \leq \text{bot}(\mathcal{N}') \\ |\mathcal{N}| + |\mathcal{N}'| = n}} (-1)^{|\text{Ad}(\mathcal{N}) \cap \text{Ad}(\mathcal{N}')|} \text{sgn}(\sigma_{\mathcal{N}, \mathcal{N}'}) \mathcal{N} \otimes \mathcal{N}',$$

where $\bullet \cdots \bullet$ stands for the linear graph with n vertices.

Proof. The image of the diagonal on the Loday associahedra under the functor C_\bullet^{cell} defines a diagonal on the operad A_∞ as this functor is strong monoidal. This diagonal $\Delta^K : A_\infty \rightarrow A_\infty \otimes A_\infty$ is determined by the image of the generating operations of the quasi-free operad A_∞ , which are the trivially nested linear graphs. The signs arise from the choices of cellular orientations on the Loday associahedra made in Section 4.1.3 as follows. As explained in the proof of [Lap22, Proposition 4.27], the computation of the signs boils down to the computation of the determinant of the bases e_j^F, e_j^G determining the cellular orientations of the faces F and G associated to the nestings \mathcal{N} and \mathcal{N}' , expressed in the basis e_j of the top dimensional cell of K_n . The second part of the proof of [Lap22, Theorem 1.26] shows that $\dim(F \cap \rho_z G) = 0$, for any $z \in (\hat{F} + \hat{G})/2$. Combined with the fact that $\dim F + \dim G = \dim K_n$, this implies that the two bases e_j^F, e_j^G form together a basis of the linear span of K_n . Writing horizontally the e_j^F and then the e_j^G in the basis e_j defines a square matrix. The positions of the rightmost non-zero entries of each line are given by the admissible edges of \mathcal{N} and \mathcal{N}' . The permutation $\sigma_{\mathcal{N}, \mathcal{N}'}$ corresponds to a permutation of the lines of this matrix, sending these rightmost entries to the diagonal, except for one case: when \mathcal{N} and \mathcal{N}' share the same admissible edge. In this case, linear independence guarantees that the two vectors differ in another place. We moreover point out that the -1 term in the definition of the permutation $\sigma_{\mathcal{N}, \mathcal{N}'}$ in Lemma 4.13 stems from the fact that K_n is defined in \mathbb{R}^{n-1} but has dimension $n - 2$. \square

We compute in particular

$$\begin{aligned}
\Delta^K((\bullet\bullet)) &= (\bullet\bullet) \otimes (\bullet\bullet), \\
\Delta^K((\bullet\bullet\bullet)) &= ((\bullet\bullet)\bullet) \otimes (\bullet\bullet\bullet) + (\bullet\bullet\bullet) \otimes (\bullet(\bullet\bullet)), \\
\Delta^K((\bullet\bullet\bullet\bullet)) &= (\bullet\bullet\bullet\bullet) \otimes (\bullet(\bullet(\bullet\bullet))) + (((\bullet\bullet)\bullet)\bullet) \otimes (\bullet\bullet\bullet\bullet) \\
&\quad - ((\bullet\bullet)\bullet\bullet) \otimes (\bullet\bullet(\bullet\bullet)) + ((\bullet\bullet\bullet)\bullet) \otimes (\bullet(\bullet\bullet)\bullet) \\
&\quad + ((\bullet\bullet\bullet)\bullet) \otimes (\bullet(\bullet\bullet\bullet)) + (\bullet(\bullet\bullet)\bullet) \otimes (\bullet(\bullet\bullet\bullet)).
\end{aligned}$$

REMARK 4.16. Proposition 4.15 completes the work of [MTTV21], by explicitly computing the signs for the polytopal diagonal on the dg level. This formula corresponds in fact to the formula originally computed in [MS06] (up to signs verification). We also conjecture that this diagonal is equal to the diagonal constructed in [SU04].

Definition 4.17 (Tensor product of A_∞ -algebras). *Given A and B two A_∞ -algebras, their tensor product as A_∞ -algebras is defined to be the dg module $A \otimes B$ endowed with the A_∞ -algebra structure induced by the diagonal Δ^K .*

Proposition 4.18. *The image under the functor C_\bullet^{cell} of the diagonal on the Forcey–Loday multiplihedra constructed in this paper defines a diagonal on the operadic bimodule M_∞ , that we denote Δ^J . It is determined by the formula*

$$\Delta^J((\bullet \cdots \bullet)) = \sum_{\mathcal{N}, \mathcal{N}'} (-1)^{|\text{Ad}(\mathcal{N}) \cap \text{Ad}(\mathcal{N}')|} \text{sgn}(\sigma_{\mathcal{N}\mathcal{N}'}) \mathcal{N} \otimes \mathcal{N}',$$

where the sum runs over the pairs $\mathcal{N}, \mathcal{N}' \in \mathcal{N}_n^2$ such that $|\text{mono}(\mathcal{N})| + |\text{mono}(\mathcal{N}')| = n - 1$ and which satisfy the conditions in Theorem 2.

Proof. The proof is similar to the proof of Proposition 4.15. Note that in this case, there is no -1 term in the definition of the permutation $\sigma_{\mathcal{N}\mathcal{N}'}$ in Lemma 4.14 since J_n is full-dimensional. \square

We compute in particular

$$\begin{aligned}
\Delta^J((\bullet\)) &= (\bullet) \otimes (\bullet), \\
\Delta^J((\bullet\bullet)) &= (\bullet\bullet) \otimes (\bullet\bullet) + (\bullet\bullet) \otimes (\bullet(\bullet)), \\
\Delta^J((\bullet\bullet\bullet)) &= ((\bullet\bullet)\bullet) \otimes (\bullet\bullet\bullet) + (\bullet\bullet\bullet) \otimes (\bullet(\bullet\bullet)) \\
&\quad - (\bullet\bullet\bullet) \otimes (\bullet(\bullet\bullet)) - (\bullet\bullet\bullet) \otimes (\bullet(\bullet\bullet)) \\
&\quad + (\bullet(\bullet\bullet)) \otimes (\bullet(\bullet\bullet)) - ((\bullet\bullet)\bullet) \otimes (\bullet(\bullet\bullet)\bullet) \\
&\quad + ((\bullet\bullet)\bullet) \otimes (\bullet\bullet\bullet) + ((\bullet\bullet)\bullet) \otimes (\bullet\bullet\bullet).
\end{aligned}$$

Definition 4.19 (Tensor product of A_∞ -morphisms). *Let $F_1 : A_1 \rightsquigarrow B_1$ and $F_2 : A_2 \rightsquigarrow B_2$ be two A_∞ -morphisms between A_∞ -algebras. Their tensor product is defined to be the A_∞ -morphism $F_1 \otimes F_2 : A_1 \otimes A_2 \rightsquigarrow B_1 \otimes B_2$ induced by the diagonal Δ^J on M_∞ .*

One can ask whether the dg "magical formula" for the diagonal on the operad A_∞ also defines a diagonal on the operadic bimodule M_∞ , i.e. if by relaxing the conditions of Theorem 2 to the condition $\text{top}(\mathcal{N}) \leq \text{bot}(\mathcal{N}')$, the formula of Proposition 4.18 still defines a diagonal on M_∞ . A simple computation in arity 4 shows that the answer to this question is negative. In other words, it is not possible to naively extend the "magical formula" for the tensor product of A_∞ -algebras to define a tensor product of A_∞ -morphisms, see also Section 3.3.

4.3. Categorification.

4.3.1. Tensor product of A_∞ -categories and A_∞ -functors. The horizontal categorifications of the notions of A_∞ -algebra and A_∞ -morphism are the notions of A_∞ -category and A_∞ -functor, respectively. We refer to [Sei08, Chapter 1] for the definitions of these two notions. We borrow the notations from [Sei08] and will moreover use the sign conventions of Section 4.1.

Definition 4.20 (Tensor product of A_∞ -categories). *The tensor product of two A_∞ -categories \mathcal{A} and \mathcal{B} is given by*

- *the set of objects $\text{Ob}(\mathcal{A} \otimes \mathcal{B}) := \text{Ob}(\mathcal{A}) \times \text{Ob}(\mathcal{B})$,*
- *for each pair of objects $X_1 \times Y_1, X_2 \times Y_2 \in \text{Ob}(\mathcal{A} \otimes \mathcal{B})$, the dg module of morphisms*

$$\mathcal{A} \otimes \mathcal{B}(X_1 \times Y_1, X_2 \times Y_2) := \mathcal{A}(X_1, X_2) \otimes \mathcal{B}(Y_1, Y_2) ,$$

and by defining the higher compositions m_n as in Proposition 4.15.

Definition 4.21 (Tensor product of A_∞ -functors). *The tensor product of two A_∞ -functors $\mathcal{F} : \mathcal{A}_1 \rightsquigarrow \mathcal{B}_1$ and $\mathcal{G} : \mathcal{A}_2 \rightsquigarrow \mathcal{B}_2$ is given by the function*

$$\text{Ob}(\mathcal{F} \otimes \mathcal{G}) := \text{Ob}(\mathcal{F}) \times \text{Ob}(\mathcal{G}) : \text{Ob}(\mathcal{A}_1 \otimes \mathcal{B}_1) \rightarrow \text{Ob}(\mathcal{A}_2 \otimes \mathcal{B}_2) ,$$

and by defining the operations $(\mathcal{F} \otimes \mathcal{G})_n$ as in Proposition 4.18.

4.3.2. Identities. The category $H_*(\mathcal{A})$ associated to an A_∞ -category \mathcal{A} does not necessarily have identity morphisms. As explained in [Sei08, Section 1.2], there exist three notions of A_∞ -category with identity morphisms : *strictly unital A_∞ -category*, *cohomologically unital A_∞ -category* and *homotopy unital A_∞ -category*.

- (1) A *cohomologically unital A_∞ -category* is an A_∞ -category \mathcal{A} which is such that $H_*(\mathcal{A})$ has identity morphisms.
- (2) A *strictly unital A_∞ -category* is an A_∞ -category together with an element $e_X \in \mathcal{A}(X, X)$ for every $X \in \text{Ob}(\mathcal{A})$ such that $\partial(e_X) = 0$, $m_2(e, \cdot) = m_2(\cdot, e) = \text{id}$ and $m_n(\cdots, e, \cdots) = 0$ for $n \geq 3$.
- (3) A *homotopy unital A_∞ -category* is defined to be an A_∞ -category together with elements $e_X \in \mathcal{A}(X, X)$ and endowed with additional operations encoding the fact that the previous relations on the m_n and the e_X are satisfied only up to higher coherent homotopies, see also [HM12, Section 6.1].

We have in particular that

$$\text{unital} \Rightarrow \text{homotopy unital} \Rightarrow \text{cohomologically unital} .$$

The proof of the following proposition is straightforward.

Proposition 4.22.

- (1) *If \mathcal{A} and \mathcal{B} are cohomologically unital A_∞ -categories, the tensor A_∞ -category $\mathcal{A} \otimes \mathcal{B}$ is again cohomologically unital.*
- (2) *If \mathcal{A} and \mathcal{B} are unital A_∞ -categories, the tensor A_∞ -category $\mathcal{A} \otimes \mathcal{B}$ is again unital, with identity morphisms $e_{X \times Y} := e_X \otimes e_Y$ for $X \in \text{Ob}(\mathcal{A})$ and $Y \in \text{Ob}(\mathcal{B})$.*

If \mathcal{A} and \mathcal{B} are homotopy unital A_∞ -categories, we have to define the additional operations associated to the fact that the elements $e_X \otimes e_Y$ are identity morphisms up to homotopy in order to endow the A_∞ -category $\mathcal{A} \otimes \mathcal{B}$ with a homotopy unital A_∞ -category structure. In other words, we have to define a diagonal on the operad $\text{u}A_\infty$ encoding homotopy unital A_∞ -algebras, which has not been done yet to the authors knowledge. An idea would be to

define a diagonal on the unital associahedra, which are CW-complexes constructed by Muro and Tonks in [MT14] and which form an operad whose image under the cellular chains is the operad uA_∞ . However, not all unital associahedra are polytopes, meaning that the present techniques cannot be directly applied to them.

4.4. Homotopy properties of diagonals on A_∞ and M_∞ .

4.4.1. *The 2-colored viewpoint.* The operad A_∞ together with the operadic bimodule M_∞ define the quasi-free 2-colored operad

$$A_\infty^2 := (\mathcal{T}(\text{red } \vee, \text{red } \wedge, \text{red } \times, \dots, \text{blue } \vee, \text{blue } \wedge, \text{blue } \times, \dots, \text{red } \oplus, \text{red } \otimes, \text{red } \boxtimes, \text{red } \boxtimes, \dots), \partial) ,$$

whose differential is given by the equations of Definition 4.1 and Definition 4.2. We refer to [Yau16, Section 11] for a complete definition of a 2-colored operad. The data of A_∞ -algebra structures on two dg modules A and B together with an A_∞ -morphism $A \rightsquigarrow B$ between them is equivalent to a morphism of 2-colored operads $A_\infty^2 \rightarrow \text{End}(A; B)$, where $\text{End}(A; B)$ is the *endomorphism 2-colored operad* naturally associated to A and B . The data of a diagonal on the operad A_∞ and of a diagonal on the operadic bimodule M_∞ is moreover equivalent to the datum of a morphism of 2-colored operads $A_\infty^2 \rightarrow A_\infty^2 \otimes A_\infty^2$, while the composition of A_∞ -morphisms can be defined by a morphism of 2-colored operads $A_\infty^2 \rightarrow A_\infty^2 \circ_{A_\infty} A_\infty^2$.

4.4.2. *Coassociativity and cocommutativity.* First, we would like to know whether given three A_∞ -algebras A , B and C , the two A_∞ -algebra structures $(A \otimes B) \otimes C$ and $A \otimes (B \otimes C)$ on the dg module $A \otimes B \otimes C$ are the same. In operadic terms, this amounts to ask if the diagonal on A_∞ is coassociative.

Proposition 4.23.

- (1) *There is no diagonal on the operad A_∞ which is coassociative.*
- (2) *There is no diagonal on the operadic bimodule M_∞ which is coassociative.*

Proof. The non-existence of a coassociative diagonal on the operad A_∞ was already proven in [MS06, Section 6]. The non-existence of a coassociative diagonal on the operad A_∞ implies the non-existence of a coassociative diagonal on the operad M_∞ . Given indeed diagonals Δ^{A_∞} and Δ^{M_∞} , it is not possible to compare the two morphisms of dg operadic bimodules $(\Delta^{M_\infty} \otimes \text{id}^{M_\infty}) \Delta^{M_\infty}$ and $(\text{id}^{M_\infty} \otimes \Delta^{M_\infty}) \Delta^{M_\infty}$, as the (A_∞, A_∞) -operadic bimodule structures induced on $M_\infty^{\otimes 3}$ by $(\Delta^{A_\infty} \otimes \text{id}^{A_\infty}) \Delta^{A_\infty}$ and $(\text{id}^{A_\infty} \otimes \Delta^{A_\infty}) \Delta^{A_\infty}$ do not coincide. We can in fact prove a stronger result: for any diagonal $\Delta : M_\infty \rightarrow M_\infty \otimes M_\infty$, we have that

$$((\text{id} \otimes \Delta) \Delta - (\Delta \otimes \text{id}) \Delta) ((\bullet \bullet \bullet)) \neq 0 .$$

The proof of this result involves computations identical to the ones of [MS06, Section 6], that we do not include for the sake of concision. \square

This proposition implies in particular that a diagonal on the 2-colored operad A_∞^2 is never coassociative. In the specific cases of Δ^K and Δ^J we compute moreover that

$$\begin{aligned} & ((\text{id} \otimes \Delta^K) \Delta^K - (\Delta^K \otimes \text{id}) \Delta^K) ((\bullet \bullet \bullet \bullet)) \\ &= -\partial(((\bullet \bullet \bullet) \bullet) \otimes (\bullet (\bullet \bullet) \bullet) \otimes (\bullet (\bullet \bullet \bullet))) , \end{aligned}$$

and that

$$\begin{aligned} & \left((\text{id} \otimes \Delta^J) \Delta^J - (\Delta^J \otimes \text{id}) \Delta^J \right) ((\bullet \bullet \bullet)) \\ &= \partial \left((\bullet \bullet \bullet) \otimes (\bullet (\bullet \bullet)) \otimes (\bullet (\bullet \bullet)) - ((\bullet \bullet) \bullet) \otimes ((\bullet \bullet) \bullet) \otimes (\bullet \bullet \bullet) \right). \end{aligned}$$

Given two A_∞ -algebras A and B , we would also like to know whether the A_∞ -algebra structure on $B \otimes A$ can simply be obtained from the maps defining the A_∞ -algebra structure on $A \otimes B$

$$m_n^{A \otimes B} : (A \otimes B)^{\otimes n} \rightarrow A \otimes B$$

by rearranging $(A \otimes B)^{\otimes n}$ into $(B \otimes A)^{\otimes n}$ and $A \otimes B$ into $B \otimes A$. In operadic terms, this amounts to ask if the diagonal on A_∞ is cocommutative or not.

Proposition 4.24. *The diagonals Δ^K and Δ^J are not cocommutative.*

Proof. We compute indeed that

$$\left(\Delta^K - \tau \Delta^K \right) ((\bullet \bullet \bullet)) = \partial \left((\bullet \bullet \bullet) \otimes (\bullet \bullet \bullet) \right),$$

where τ acts by the permutation $(1\ 2)$ on the operad $A_\infty \otimes A_\infty$. We also compute that

$$\left(\Delta^J - \tau \Delta^J \right) ((\bullet \bullet \bullet)) = \partial \left((\bullet \bullet \bullet) \otimes (\bullet \bullet \bullet) \right).$$

□

We conjecture in fact that Proposition 4.24 holds for any diagonal on the operad A_∞ and for any diagonal on the operadic bimodule M_∞ .

4.4.3. Compatibility with the composition. We would finally like to know whether the tensor product is functorial with respect to the composition of A_∞ -morphisms. In other words, if given four A_∞ -morphisms $F_1 : A_1 \rightsquigarrow B_1$, $G_1 : B_1 \rightsquigarrow C_1$, $F_2 : A_2 \rightsquigarrow B_2$ and $G_2 : B_2 \rightsquigarrow C_2$ they satisfy the following equality

$$(G_1 \otimes F_1) \circ (G_2 \otimes F_2) = (G_1 \otimes G_2) \circ (F_1 \otimes F_2).$$

In operadic terms, this amounts to ask if the diagonal Δ on M_∞ together with the composition morphism comp of Section 4.1.2 satisfy the following equality

$$(\text{comp} \otimes \text{comp}) \Delta = (\Delta \circ_{A_\infty} \Delta) \text{comp}.$$

Proposition 4.25. *There is no diagonal on the operadic bimodule M_∞ which is compatible with the composition of A_∞ -morphisms.*

Proof. Let Δ be a diagonal $M_\infty \rightarrow M_\infty \otimes M_\infty$. The compatibility with the differential implies that Δ is necessarily of the form

$$\Delta((\bullet \bullet)) = (\bullet \bullet) \otimes (\bullet \bullet)$$

and

$$\begin{aligned} \Delta((\bullet \bullet \bullet)) &= \alpha \left((\bullet \bullet \bullet) \otimes (\bullet \bullet) + (\bullet \bullet \bullet) \otimes (\bullet \bullet) \right) \\ &\quad + (1 - \alpha) \left((\bullet \bullet \bullet) \otimes (\bullet \bullet) + (\bullet \bullet \bullet) \otimes (\bullet \bullet) \right), \end{aligned}$$

where $\alpha \in \mathbb{Z}$. We compute that if the equality

$$(\text{comp} \otimes \text{comp}) \Delta((\bullet \bullet \bullet)) = (\Delta \circ_{A_\infty} \Delta) \text{comp}((\bullet \bullet \bullet))$$

holds, we necessarily have that $\alpha = 0$ and that $\alpha = 1$, which is not possible. □

In the case of the diagonals Δ^K and Δ^J , we compute that

$$\left(\text{comp} \circ \Delta^J - (\Delta^J \circ_{A_\infty} \Delta^J) \circ \text{comp} \right) (\text{diagram}) = \partial \left(\text{diagram} \otimes \text{diagram} \right).$$

4.4.4. Homotopy properties. While coassociativity, cocommutativity and compatibility with the composition are not satisfied by the diagonals Δ^K and Δ^J , we will now prove that a diagonal on the 2-colored operad A_∞^2 always satisfies these properties up to homotopy. We use the notion of homotopy between morphisms of 2-colored operads as defined in [MSS02, Section 3.10].

Proposition 4.26. *Let Δ be a diagonal on the 2-colored operad A_∞^2 .*

- (1) *The morphisms of operads $(\Delta \otimes \text{id})\Delta$ and $(\text{id} \otimes \Delta)\Delta$ are homotopic. In other words, a diagonal on A_∞^2 is always coassociative up to homotopy.*
- (2) *The morphisms of operads Δ and $\tau\Delta$ are homotopic. In other words, a diagonal on A_∞^2 is always cocommutative up to homotopy.*
- (3) *The morphisms of operads $\text{comp} \circ \Delta^J$ and $(\Delta^J \circ_{A_\infty} \Delta^J) \circ \text{comp}$ are homotopic. In other words, a diagonal on A_∞^2 is always compatible with the composition of A_∞ -morphisms up to homotopy.*

Proof. The proof of this proposition is a simple adaptation of the results of [MS06, Section 2] in the context of 2-colored dg operads, applied to the minimal model A_∞^2 for the 2-colored dg operad As^2 encoding pairs of dg algebras together with morphisms between them. \square

While Proposition 4.25 shows that it is not possible to endow the category $\infty\text{-}A_\infty\text{-alg}$ with a symmetric monoidal category structure using the viewpoint of diagonals, Proposition 4.26 exhibits a first level of homotopies that could be involved in the definition of some kind of *homotopy symmetric monoidal* category structure on $\infty\text{-}A_\infty\text{-alg}$. This question will be studied in a future work by D. Poliakova and the two authors of this paper. As a first step towards solving that problem, we will inspect in particular which higher coherent homotopies arise from the lack of coassociativity of Δ^{K_n} and Δ^{J_n} on the level of polytopes.

5. FURTHER APPLICATIONS

We first prove that a diagonal on the dg operad A_∞ is equivalent to a retraction of the bar-cobar resolution AA_∞ onto the operad A_∞ . We then explain how to associate a convolution A_∞ -algebra to an A_∞ -coalgebra and an A_∞ -algebra, as well as A_∞ -morphisms between convolution A_∞ -algebras, using diagonals on A_∞ and M_∞ . We finally describe two possible applications of our results in symplectic topology: in the context of Heegard Floer homology, and to study tensor products of Fukaya categories/algebras and A_∞ -functors between them.

5.1. Retractions and diagonals. Recall that the operad A_∞ is the minimal model $A_\infty = \Omega As^i$ of the dg operad As encoding associative algebras. Another cofibrant replacement of the operad As is given by the bar-cobar (or Boardman-Vogt) resolution $AA_\infty := \Omega BAs$, which is defined as the quasi-free operad

$$AA_\infty := (\mathcal{T}(\Upsilon, \Upsilon, \Upsilon, \Upsilon, \dots, \text{PT}_n, \dots), \partial),$$

where PT_n is the set of planar rooted trees of arity n and the degree of a tree is defined as the number of its internal edges. We refer to [LV12, Section 9.3] for a complete study of the operad AA_∞ , and in particular for a definition of its differential. There exists an explicit embedding

of dg operads $A_\infty \rightarrow AA_\infty$, as constructed in [MS06, Section 4] and in [Maz21a, Section 1.3.1.5]. The problem of the construction of an explicit morphism of dg operads $AA_\infty \rightarrow A_\infty$ is more complicated and is the subject of the following proposition.

Definition 5.1 (Retraction). *A morphism of dg operads $AA_\infty \rightarrow A_\infty$ sending \curlywedge to \curlywedge will be called a retraction of the operad AA_∞ onto the operad A_∞ .*

Proposition 5.2. *The datum of a diagonal on the operad A_∞ is equivalent to the datum of a retraction $r : AA_\infty \rightarrow A_\infty$.*

Proof. We apply the general theory of operadic twisting morphisms [LV12, Section 6.4] to prove the following sequence of isomorphisms:

$$\begin{aligned} \mathrm{Hom}_{\mathrm{Op}}(\Omega As^i, \Omega As^i \otimes \Omega As^i) &\cong \mathrm{Tw}(As^i, \Omega As^i \otimes \Omega As^i) \\ &\cong \mathrm{Tw}(BAs, \Omega As^i) \\ &\cong \mathrm{Hom}_{\mathrm{Op}}(\Omega BAs, \Omega As^i) . \end{aligned}$$

The first and last isomorphisms are given by the bar-cobar adjunction. We thus only need to explain the second isomorphism. A twisting morphism $As^i \rightarrow \Omega As^i \otimes \Omega As^i$ is by definition a Maurer–Cartan element in the convolution pre-Lie algebra associated to the convolution dg operad $\mathrm{Hom}(As^i, \Omega As^i \otimes \Omega As^i)$. This convolution dg operad is in turn isomorphic to the desuspension $\mathcal{S}^{-1}(\Omega As^i \otimes \Omega As^i)$. Since the cooperad As^i is 1-dimensional in every arity, and since the arity-wise linear dual dg cooperad of the desuspended dg operad $\mathcal{S}^{-1}(\Omega As^i)$ is isomorphic to the bar construction BAs , we have that the desuspension $\mathcal{S}^{-1}(\Omega As^i \otimes \Omega As^i)$ is isomorphic to the convolution dg operad $\mathrm{Hom}(BAs, \Omega As^i)$. We hence have the following isomorphisms of dg operads

$$\mathrm{Hom}(As^i, \Omega As^i \otimes \Omega As^i) \cong \mathcal{S}^{-1}(\Omega As^i \otimes \Omega As^i) \cong \mathrm{Hom}(BAs, \Omega As^i) .$$

This implies an isomorphism on the level of the Maurer–Cartan elements of the associated dg pre-Lie algebras, that is

$$\mathrm{Tw}(As^i, \Omega As^i \otimes \Omega As^i) \cong \mathrm{Tw}(BAs, \Omega As^i) .$$

We finally check that the condition $\Delta(\curlywedge) = \curlywedge \otimes \curlywedge$ is equivalent to the condition $r(\curlywedge) = \curlywedge$. \square

Proposition 5.2 clarifies in particular the construction of the diagonal on the operad A_∞ given in [MS06]. The operad AA_∞ can indeed be seen as the cellular chains on the cubical realization of the associahedra [LV12, Section 9.3.1]. It comes with an elementary diagonal $AA_\infty \rightarrow AA_\infty \otimes AA_\infty$ defined using the Serre cubical diagonal of [Ser51]. M. Markl and S. Shnider then define a retraction $r : AA_\infty \rightarrow A_\infty$ and deduce a diagonal on the operad A_∞ as the composite

$$A_\infty \longrightarrow AA_\infty \longrightarrow AA_\infty \otimes AA_\infty \xrightarrow{r \otimes r} A_\infty \otimes A_\infty .$$

Their choice of retraction recovers the diagonal constructed directly on the level of the associahedra in [MTTV21, Theorem 2]. A similar proof would however not adapt to the case of the multiplihedra, as they are not simple polytopes hence do not admit a cubical realization.

REMARK 5.3. As observed in [Lap22, Remark 1.6], the methods used to construct our cellular approximation of the diagonal could be related to the Fulton–Sturmfels formula [FS97, Theorem 4.2], appearing in the study of the intersection theory on toric varieties. We also expect an interpretation of Proposition 5.2 in terms of Morse theory, in the vein of [FMMS21, Fra07]. There should also be an interpretation in terms of discrete Morse theory as in [Tho18, Section 1.1.4] for the case of the standard simplices.

5.2. Convolution A_∞ -algebra.

5.2.1. *Standard convolution algebra.* Given a dg algebra A and a dg coalgebra C , recall from [LV12, Section 1.6] that one can define the *convolution algebra* of C and A as the dg algebra $(\text{Hom}(C, A), [\partial, \cdot], \star)$, where $\text{Hom}(C, A)$ is the dg module of maps $C \rightarrow A$, endowed with the convolution product $f \star g := \mu_A \circ (f \otimes g) \circ \Delta_C$. The convolution algebra construction is in fact functorial, i.e. fits into a bifunctor $(\text{dg-coalg})^{\text{op}} \times \text{dg-alg} \rightarrow \text{dg-alg}$ defined on objects as $(C, A) \mapsto \text{Hom}(C, A)$. A Maurer-Cartan element α of $\text{Hom}(C, A)$, i.e. a map $\alpha : C \rightarrow A$ such that $[\partial, \alpha] + \alpha \star \alpha = 0$, is then called a *twisting morphism*. Twisting morphisms define twisted differentials on the tensor product $C \otimes A$ via the formula

$$\partial_\alpha := \partial_{C \otimes A} + (\text{id} \otimes \mu_A)(\text{id} \otimes \alpha \otimes \text{id})(\Delta_C \otimes \text{id}) .$$

Twisted differentials appear in the computation of the singular homology of fiber spaces [Bro59]. Given a fibration $F \rightarrow X \rightarrow B$ satisfying some mild assumptions, the singular homology of X can then be computed as the homology of the tensor product $C_*(B) \otimes C_*(F)$ endowed with a twisted differential, where $C_*(F)$ is seen as a dg module over the dg algebra $C_*(\Omega B)$.

5.2.2. *Convolution A_∞ -algebra.* One defines an A_∞ -coalgebra structure on a dg module C to be a morphism of dg operads $A_\infty \rightarrow \text{coEnd}_C$, where $\text{coEnd}_C(n) = \text{Hom}(C, C^{\otimes n})$. Put differently, it is the structure dual to the structure of A_∞ -algebra, i.e. it corresponds to a collection of operations $c_n : C \rightarrow C^{\otimes n}$ of degree $n - 2$ satisfying the equations obtained by inverting inputs and outputs in the equations for A_∞ -algebras. The notion of an A_∞ -morphism between A_∞ -coalgebras is defined in a similar fashion: either in terms of operations $f_n : C \rightarrow D^{\otimes n}$ of degree $n - 1$ and satisfying the equations dual to the equations for A_∞ -morphisms, or equivalently as a morphism of dg operadic bimodules $M_\infty \rightarrow \text{coHom}_{C_2}^{C_1}$. Our results allow us to extend the convolution algebra construction when C is an A_∞ -coalgebra and A is an A_∞ -algebra.

Proposition 5.4.

- (1) Let C be an A_∞ -coalgebra and A be an A_∞ -algebra. A diagonal on the operad A_∞ yields an A_∞ -algebra structure on the dg module $(\text{Hom}(C, A), \partial)$. We call this A_∞ -algebra the convolution A_∞ -algebra of C and A .
- (2) Let $F : A_1 \rightsquigarrow A_2$ be an A_∞ -morphism between two A_∞ -algebras A_1 and A_2 and $G : C_2 \rightsquigarrow C_1$ be an A_∞ -morphism between two A_∞ -coalgebras C_2 and C_1 . A diagonal on the operad M_∞ yields an A_∞ -morphism between the convolution A_∞ -algebras $\text{Hom}(C_1, A_1)$ and $\text{Hom}(C_2, A_2)$.

Proof.

- (1) Given a diagonal $A_\infty \rightarrow A_\infty \otimes A_\infty$, the following composite of morphism of operads defines the A_∞ -algebra structure on $\text{Hom}(C, A)$:

$$A_\infty \rightarrow A_\infty \otimes A_\infty \rightarrow \text{coEnd}_C \otimes \text{End}_A \rightarrow \text{End}_{\text{Hom}(C, A)} ,$$

where the morphism of dg operads $\text{coEnd}_C \otimes \text{End}_A \rightarrow \text{End}_{\text{Hom}(C,A)}$ is straightforward to define.

- (2) Given a diagonal $M_\infty \rightarrow M_\infty \otimes M_\infty$, we consider in a similar fashion the composite of morphism of operadic bimodules

$$M_\infty \rightarrow M_\infty \otimes M_\infty \rightarrow \text{coHom}_{C_1}^{C_2} \otimes \text{Hom}_{A_2}^{A_1} \rightarrow \text{Hom}_{\text{Hom}(C_2,A_2)}^{\text{Hom}(C_1,A_1)}.$$

□

Proposition 5.5. *For any diagonal on A_∞ , and for any diagonal on M_∞ , the convolution A_∞ -algebra $\text{Hom}(C, A)$ does not define a bifunctor $(\infty\text{-}A_\infty\text{-cog})^{\text{op}} \times \infty\text{-}A_\infty\text{-alg} \rightarrow \infty\text{-}A_\infty\text{-alg}$.*

Proof. This is a direct corollary to Proposition 4.25. □

Proposition 5.4 implies in particular that for an A_∞ -coalgebra C and an A_∞ -algebra A , it is still possible to define the notion of a *twisting morphism* $\alpha : C \rightarrow A$ as a Maurer-Cartan element in the A_∞ -algebra $\text{Hom}(C, A)$, see [DSV18, Equation 1, p.8] for instance. It also implies that the A_∞ -morphism $\text{Hom}(C_1, A_1) \rightsquigarrow \text{Hom}(C_2, A_2)$ defined by the A_∞ -morphism $F : A_1 \rightsquigarrow A_2$ and $G : C_2 \rightsquigarrow C_1$, sends a twisting morphism $C_1 \rightarrow A_1$ to a twisting morphism $C_2 \rightarrow A_2$. We will use this key property in order to pursue the work of Brown [Bro59] and [Pro86] on the homology of fibered spaces in a forthcoming paper.

5.2.3. Diagonals as twisting morphisms. The results of Section 5.2.2 can be interpreted in a more general framework, developed by D. Robert-Nicoud and F. Wierstra in [RNW19b, RNW19a].

Proposition 5.6. *The datum of a diagonal on A_∞ is equivalent to the datum of a twisting morphism $\alpha \in \text{Tw}(BAS, \Omega As^i)$ sending \curlyvee to \curlyvee .*

Proof. This result was proven in the proof of Proposition 5.2. □

Setting $\mathcal{C} = BAS$ and $\mathcal{P} = \Omega As^i$ and working in the context of non-symmetric operads where the operad L_∞ of [RNW19b, RNW19a] is replaced by the operad A_∞ , we recover Proposition 5.6 (and thus Proposition 5.2) via [RNW19b, Theorem 7.1] and Point (1) of Proposition 5.4 via [RNW19b, Theorem 4.1]. We denote by $A_\infty\text{-alg}$ the category of A_∞ -algebras and their *strict* morphisms [LV12, Section 10.2.1]. It is shown in [RNW19b, Corollary 5.4] that the assignments

$$(7) \quad \text{Hom}(-, \text{id}) : (\infty\text{-}A_\infty\text{-cog})^{\text{op}} \times A_\infty\text{-alg} \rightarrow A_\infty\text{-alg}$$

$$(8) \quad \text{Hom}(\text{id}, -) : (A_\infty\text{-cog})^{\text{op}} \times \infty\text{-}A_\infty\text{-alg} \rightarrow A_\infty\text{-alg}$$

given by the convolution A_∞ -algebra extend to bifunctors. The authors also show that these two bifunctors do *not* extend to a bifunctor

$$(9) \quad \text{Hom}(-, -) : (\infty\text{-}A_\infty\text{-cog})^{\text{op}} \times \infty\text{-}A_\infty\text{-alg} \rightarrow \infty\text{-}A_\infty\text{-alg}$$

in general, since this assignment is not compatible with the composition of A_∞ -morphisms [RNW19b, Theorem 6.6]. Point (2) of Proposition 5.4 allows us to define the assignment (9) directly, and Proposition 5.5 can be seen as a stronger version of [RNW19b, Theorem 6.6], in the special case of A_∞ -algebras.

The main result of [RNW19a] says that if a twisting morphism $\alpha \in \text{Tw}(BAS, \Omega As^i)$ is Koszul, then the possible compositions of the two bifunctors (7) and (8) are homotopic and that they extend to a bifunctor on the level of the homotopy categories [RNW19a, Theorem 3.6 and

Corollary 3.8]. This should be seen as a statement analogous to Point (3) of Proposition 4.26. It would be interesting to know how the results of [RNW19b, RNW19a] can be interpreted from the viewpoint of diagonals, and if they admit an interpretation on the level of polytopes.

5.3. Diagonals in symplectic topology.

5.3.1. *The work of Lipshitz, Ozsváth and Thurston.* In [LOT20], R. Lipshitz, P. Ozsváth and D. Thurston also study diagonals on the dg operad A_∞ and on the dg operadic bimodule M_∞ . They however work exclusively on the dg level, constructing abstract diagonals by using the fact that A_∞ and M_∞ are contractible, and do not provide explicit formulae for these diagonals as in Proposition 4.15 and Proposition 4.18. The goal of their work is to study bordered Heegaard Floer homology of 3-manifolds. Given a 3-manifold Y with two boundary components, they aim to construct a *bimodule twisted complex* $CFDD^-(Y)$, also called a *type DD-bimodule*. The definition of such an object uses a diagonal on the dg operad A_∞ . A diagonal on M_∞ is then needed in order to relate the categories of bimodules defined with different diagonals on A_∞ , which in turn is needed for properties like the associativity of tensor products. They also expect that diagonals on M_∞ could be needed in a distant future to define A_∞ -morphisms between bimodule twisted complexes arising from a cobordism between 3-manifolds Y_1 and Y_2 . Thus, the explicit formula for the diagonal defined in this paper could be used to compute invariants of 3 and 4-manifolds, via implementation in a computer program for instance.

5.3.2. *Künneth theorems in Lagrangian Floer theory.* Let (M, ω) be a closed symplectic manifold, i.e. a closed manifold M together with a closed non-degenerate 2-form ω on M . The *Fukaya category* $\text{Fuk}(M, \omega)$ of (M, ω) is defined to be the (curved filtered unital) A_∞ -category whose objects are (unobstructed) Lagrangian submanifolds of M and higher compositions are defined by counting pseudo-holomorphic disks with Lagrangian boundary conditions and marked points on their boundary, as represented in Figure 11. We refer for instance to [Smi15] and [Aur14] for introductions to this subject. Given a closed spin Lagrangian submanifold $L \subset M$, K. Fukaya also constructs in [Fuk10] a strictly unital A_∞ -algebra $\mathcal{F}(L)$, the *Fukaya algebra* of the Lagrangian L , whose higher multiplications are again defined by counting pseudo-holomorphic disks.




FIGURE 11. On the left, a pseudo-holomorphic disk defining the A_∞ -category structure on $\text{Fuk}(M)$. On the right, a pseudo-holomorphic quilted disk defining an A_∞ -functor $\text{Fuk}(M_0) \rightsquigarrow \text{Fuk}(M_1)$

In [Amo17], L. Amorim shows that given two symplectic manifolds M_1 and M_2 together with Lagrangians $L_i \subset M_i$, the Fukaya algebra of the product Lagrangian $L_1 \times L_2$ is quasi-isomorphic to the tensor product of their Fukaya algebras, i.e. $\mathcal{F}(L_1 \times L_2) \simeq \mathcal{F}(L_1) \otimes \mathcal{F}(L_2)$. His proof relies on a theorem that he proves in [Amo16], giving a criterion for an A_∞ -algebra C to be quasi-isomorphic to the tensor A_∞ -algebra $A \otimes B$ (see Definition 4.17) of two commuting A_∞ -subalgebras $A \subset C$ and $B \subset C$, which he then applies to the two A_∞ -subalgebras $\mathcal{F}(L_1) \subset \mathcal{F}(L_1 \times L_2)$ and $\mathcal{F}(L_2) \subset \mathcal{F}(L_1 \times L_2)$. Fukaya generalizes this result in [Fuk17], working this time on the level of Fukaya categories. He proves that for two closed symplectic manifolds M_0 and M_1 there exists a unital A_∞ -functor

$$\mathrm{Fuk}(M_0) \otimes \mathrm{Fuk}(M_1) \longrightarrow \mathrm{Fuk}(M_0^- \times M_1)$$

which is a homotopy equivalence to its image.

Let now M_0 and M_1 be two compact symplectic manifolds. Define a *Lagrangian correspondence* from M_0 to M_1 to be a Lagrangian submanifold $\mathcal{L} \subset M_0^- \times M_1$. In [MWW18], S. Mau, K. Wehrheim and C. Woodward associate to a Lagrangian correspondence \mathcal{L} (with additional technical assumptions) an A_∞ -functor $\Phi_{\mathcal{L}} : \mathrm{Fuk}(M_0) \rightsquigarrow \mathrm{Fuk}(M_1)$. It is defined on objects as

$$\Phi_{\mathcal{L}}(L_0) := \pi_{M_1}(L_0 \times_{M_0} \mathcal{L}),$$

where π_{M_1} denotes the projection $M_0 \times M_0^- \times M_1 \rightarrow M_1$ and \times_{M_0} is the fiber product over M_0 . The operations of $\Phi_{\mathcal{L}}$ are defined by counting pseudo-holomorphic quilted disks with Lagrangian boundary conditions, seam condition on \mathcal{L} and marked points on their boundary, as represented in Figure 11. The tensor product of A_∞ -functors defined in the present paper allows one to consider the A_∞ -functor $\Phi_{\mathcal{L}_M} \otimes \Phi_{\mathcal{L}_N}$ associated to a pair of Lagrangian correspondences, raising the following question.

Problem. *Does the diagram*

$$\begin{array}{ccc} \mathrm{Fuk}(M_0) \otimes \mathrm{Fuk}(N_0) & \xrightarrow{\Phi_{\mathcal{L}_M} \otimes \Phi_{\mathcal{L}_N}} & \mathrm{Fuk}(M_1) \otimes \mathrm{Fuk}(N_1) \\ \downarrow \wr & & \downarrow \wr \\ \mathrm{Fuk}(M_0 \times N_0) & \xrightarrow{\Phi_{\tau(\mathcal{L}_M \times \mathcal{L}_N)}} & \mathrm{Fuk}(M_1 \times N_1) \end{array}$$

commute up to homotopy of A_∞ -functors?

In this diagram, $\mathcal{L}_M \subset M_0^- \times M_1$, $\mathcal{L}_N \subset N_0^- \times N_1$ and the symplectomorphism τ is defined by rearranging the factors of $M_0^- \times M_1 \times N_0^- \times N_1$ into the factors of $M_0^- \times N_0^- \times M_1 \times N_1$. In other words, we would like to know whether the *algebraic (tensor) product* of geometric A_∞ -functors between Fukaya categories defined in this paper is homotopic to the A_∞ -functor defined by the *geometric product* of the Lagrangian correspondences. We refer to [Fuk17, Section 13] for a discussion on two definitions of the notion of a homotopy between A_∞ -functors.

REFERENCES

- [AACD11] Camilo Arias Abad, Marius Crainic, and Benoit Dherin, *Tensor products of representations up to homotopy*, J. Homotopy Relat. Struct. **6** (2011), no. 2, 239–288.
- [AD13] Federico Ardila and Jeffrey Doker, *Lifted generalized permutahedra and composition polynomials*, Adv. in Appl. Math. **50** (2013), no. 4, 607–633.
- [Amo16] Lino Amorim, *Tensor product of filtered A_∞ -algebras*, J. Pure Appl. Algebra **220** (2016), no. 12, 3984–4016 (English).
- [Amo17] ———, *The Künneth theorem for the Fukaya algebra of a product of Lagrangians*, Int. J. Math. **28** (2017), no. 4, 38 (English), Id/No 1750026.
- [Aur14] Denis Auroux, *A beginner's introduction to Fukaya categories*, Contact and symplectic topology, Bolyai Soc. Math. Stud., vol. 26, János Bolyai Math. Soc., Budapest, 2014, pp. 85–136. MR 3220941
- [Bro59] Edgar H. Brown, Jr., *Twisted tensor products. I*, Ann. of Math. (2) **69** (1959), 223–246.
- [BS92] Louis J. Billera and Bernd Sturmfels, *Fiber polytopes*, Ann. of Math. (2) **135** (1992), no. 3, 527–549.
- [BV73] John M. Boardman and Rainer M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics, Vol. 347, Springer-Verlag, Berlin, 1973.
- [CP22] Frédéric Chapoton and Vincent Pilaud, *Shuffles of deformed permutahedra, multiplihedra, constrainahedra, and biasmahedra*, 2022, arXiv:2201.06896.
- [CZ12] Cesar Ceballos and Günter M. Ziegler, *Realizing the associahedron: mysteries and questions*, Associahedra, Tamari lattices and related structures, Progr. Math., vol. 299, Birkhäuser/Springer, Basel, 2012, pp. 119–127.
- [DF08] Satyan Devadoss and Stefan Forcey, *Marked tubes and the graph multiplihedron*, Algebr. Geom. Topol. **8** (2008), no. 4, 2081–2108.
- [Dok11] Jeffrey Samuel Doker, *Geometry of generalized permutohedra*, Ph.D. thesis, University of California, Berkeley, 2011.
- [DSV18] Vladimir Dotsenko, Sergey Shadrin, and Bruno Vallette, *The twisting procedure*, 2018, arXiv:1810.02941.
- [EML53] S. Eilenberg and S. Mac Lane, *On the groups of $H(\Pi, n)$. I*, Ann. of Math. (2) **58** (1953), 55–106.
- [FMMS21] Greg Friedman, Anibal M. Medina-Mardones, and Dev Sinha, *Flowing from intersection product to cup product*, ArXiv e-prints (2021), arXiv:2106.05986.
- [FOOO09a] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, *Lagrangian intersection Floer theory. Anomaly and obstruction. I*, vol. 46, Providence, RI: American Mathematical Society (AMS); Somerville, MA: International Press, 2009 (English).
- [FOOO09b] ———, *Lagrangian intersection Floer theory. Anomaly and obstruction. II*, vol. 46, Providence, RI: American Mathematical Society (AMS); Somerville, MA: International Press, 2009 (English).
- [For08] Stefan Forcey, *Convex hull realizations of the multiplihedra*, Topology Appl. **156** (2008), no. 2, 326–347.
- [Fra07] Martin Frankland, *Théorème de künneth en homologie de Morse*, Ann. Sci. Math. Québec **31** (2007), 31–39.
- [FS97] William Fulton and Bernd Sturmfels, *Intersection theory on toric varieties*, Topology **36** (1997), 335–353.
- [Fuk10] Kenji Fukaya, *Cyclic symmetry and adic convergence in Lagrangian Floer theory*, Kyoto J. Math. **50** (2010), no. 3, 521–590 (English).
- [Fuk17] Kenji Fukaya, *Unobstructed immersed Lagrangian correspondence and filtered A_∞ functor*, 2017, arXiv:1706.02131.
- [HM12] Joseph Hirsh and Joan Millès, *Curved Koszul duality theory*, Math. Ann. **354** (2012), no. 4, 1465–1520.
- [Lap22] Guillaume Laplante-Anfossi, *The diagonal of the operahedra*, Advances in Mathematics (2022), to appear.
- [LH03] Kenji Lefèvre-Hasegawa, *Sur les A_∞ -catégories*, Ph.D. thesis, Université Paris 7, UFR de Mathématiques, 2003.
- [Lod04] Jean-Louis Loday, *Realization of the Stasheff polytope*, Arch. Math. (Basel) **83** (2004), no. 3, 267–278.
- [LOT20] Robert Lipshitz, Peter Ozsváth, and Dylan P. Thurston, *Diagonals and A -infinity tensor products*, 2020, arXiv:2009.05222.

- [LV12] Jean-Louis Loday and Bruno Vallette, *Algebraic operads*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 346, Springer-Verlag, Berlin, 2012.
- [May72] J.P. May, *The geometry of iterated loop spaces*, Springer-Verlag, Berlin, 1972, Lectures Notes in Mathematics, Vol. 271.
- [Maz21a] Thibaut Mazuir, *Higher algebra of A_∞ and $\Omega B A_\infty$ -algebras in Morse theory I*, 2021, arXiv:2102.06654.
- [Maz21b] ———, *Higher algebra of A_∞ and $\Omega B A_\infty$ -algebras in Morse theory II*, 2021, arXiv:2102.08996.
- [MS06] Martin Markl and Steve Shnider, *Associahedra, cellular W -construction and products of A_∞ -algebras*, Trans. Amer. Math. Soc. **358** (2006), no. 6, 2353–2372 (electronic).
- [MSS02] Martin Markl, Steve Shnider, and James D. Stasheff, *Operads in algebra, topology and physics*, Mathematical Surveys and Monographs, vol. 96, American Mathematical Society, Providence, RI, 2002.
- [MT14] Fernando Muro and Andrew Tonks, *Unital associahedra*, Forum Math. **26** (2014), no. 2, 593–620. MR 3176644
- [MTTV21] Naruki Masuda, Hugh Thomas, Andy Tonks, and Bruno Vallette, *The diagonal of the associahedra*, J. Éc. Polytech., Math. **8** (2021), 121–146.
- [MW10] S. Ma'u and C. Woodward., *Geometric realizations of the multiplihedra*, Compos. Math. **146** (2010), no. 4, 1002–1028.
- [MWW18] S. Ma'u, K. Wehrheim, and C. Woodward, *A_∞ functors for Lagrangian correspondences*, Selecta Math. (N.S.) **24** (2018), no. 3, 1913–2002. MR 3816496
- [OEI22] OEIS, *The on-line encyclopedia of integer sequences*, Published electronically at <http://oeis.org>.
- [Pol20] Daria Poliakova, *Cellular chains on freehedra and operadic pairs*, 2020, arXiv:2011.11607.
- [Pos09] Alexander Postnikov, *Permutohedra, associahedra, and beyond*, Int. Math. Res. Not. IMRN **6** (2009), 1026–1106.
- [Pro86] Alain Prouté, *A_∞ -structures, modèle minimal de Baues-Lemaire et homologie des fibrations*, Ph.D. thesis, Université Paris 7, UFR de Mathématiques, 1986.
- [RNW19a] Daniel Robert-Nicoud and Felix Wierstra, *Convolution algebras and the deformation theory of infinity-morphisms*, Homology, Homotopy and Applications **21** (2019), 351–373.
- [RNW19b] ———, *Homotopy morphisms between convolution homotopy Lie algebras*, Journal of Noncommutative Geometry **13** (2019), 1435–1462.
- [Sei08] Paul Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [Ser51] Jean-Pierre Serre, *Homologie singulière des espaces fibrés. Applications*, Ann. of Math. (2) **54** (1951), 425–505.
- [Smi15] Ivan Smith, *A symplectic prolegomenon*, Bull. Amer. Math. Soc. (N.S.) **52** (2015), no. 3, 415–464. MR 3348443
- [Sta63] James D. Stasheff, *Homotopy associativity of H -spaces. I, II*, Trans. Amer. Math. Soc. **108** (1963), 275–292; ibid. **108** (1963), 293–312.
- [Sta70] ———, *H -spaces from a homotopy point of view*, Lecture Notes in Mathematics, Vol. 161, Springer-Verlag, Berlin, 1970.
- [SU04] Samson Sanedidze and Ronald Umble, *Diagonals on the permutahedra, multiplihedra and associahedra*, Homology Homotopy Appl. **6** (2004), no. 1, 363–411.
- [Tho18] Ryan George Thorngren, *Combinatorial topology and applications to quantum field theory*, Ph.D. thesis, UC Berkeley, 2018, Available online at <https://escholarship.org/uc/item/7r44w49f>.
- [Val20] Bruno Vallette, *Homotopy theory of homotopy algebras*, Ann. Inst. Fourier (Grenoble) **70** (2020), no. 2, 683–738. MR 4105949
- [Yau16] Donald Yau, *Colored operads*, Graduate Studies in Mathematics, vol. 170, American Mathematical Society, Providence, RI, 2016. MR 3444662
- [Zie95] Günter M. Ziegler, *Lectures on polytopes*, Graduate Texts in Mathematics, vol. 152, Springer-Verlag, New York, 1995.

LABORATOIRE ANALYSE, GÉOMÉTRIE ET APPLICATIONS, UNIVERSITÉ PARIS NORD 13, SORBONNE PARIS CITÉ,
CNRS, UMR 7539, 93430 VILLETANEUSE, FRANCE.

Email address: `laplante-anfossi@math.univ-paris13.fr`

SORBONNE UNIVERSITÉ, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE CNRS, CAMPUS PIERRE
ET MARIE CURIE, 75005 PARIS, FRANCE.

Email address: `thibaut.mazuir@imj-prg.fr`