

Quantum Field Theory, 2021/2022

Exercise sheet 4: Spin 1 fields & interactions

Hand-in: November 3, 2021

- 4.1. Consider the free electromagnetic field $A^\mu(x)$. In the Quantum Theory the Lorentz condition is substituted by a condition on the allowed physical states $|\Psi\rangle$

$$\partial_\mu A^{\mu+}(x)|\Psi\rangle = 0 \Rightarrow (a_{(3)\mathbf{k}} - a_{(0)\mathbf{k}})|\Psi\rangle = 0$$

- (a) From the commutation relations among $a_{(\lambda)\mathbf{k}}$, $a_{(\lambda)\mathbf{q}}^\dagger$, show that:

$$[a_{(3)\mathbf{k}} - a_{(0)\mathbf{k}}, a_{(3)\mathbf{q}}^\dagger - a_{(0)\mathbf{q}}^\dagger] = 0$$

- (b) Define

$$\alpha_i^\dagger = a_{(3)\mathbf{k}_i}^\dagger - a_{(0)\mathbf{k}_i}^\dagger$$

show that the most general allowed physical state which contains scalar and longitudinal photons can be written as:

$$|\Psi_{SL}\rangle = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots c(n_1, n_2, \dots) \prod_{i=1}^{\infty} (\alpha_i^\dagger)^{n_i} |0\rangle$$

Where the $c(n_1, n_2, \dots)$ are complex numbers.

- (c) Show that the norm of this state is given by just the coefficient with no photons present:

$$\langle \Psi_{SL} | \Psi_{SL} \rangle = |c(0, 0, \dots)|^2$$

- 4.2. Consider a real Klein-Gordon field with a cubic interaction Hamiltonian:

$$\mathcal{H}_{int} = \frac{\lambda}{3!} \phi^3$$

- (a) Compute the interaction Feynman rule for the 3-particle interaction vertex.
(b) Compute, **using the Dyson expansion**, the lowest order contribution to the transition matrix element for $p_A, p_B \rightarrow k_1, k_2$:

$$\langle k_1, k_2 | i\mathcal{T} | p_A, p_B \rangle$$

remember that only fully connected and amputated contributions have to be taken into account.