

Defining the Fourier Transform Using Complex Numbers

Valerio Velardo

Join the community!



thesoundofai.slack.com

Previously...

Previously...

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

Previously...

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

The intuition

- Use magnitude and phase as polar coordinates

The intuition

- Use magnitude and phase as polar coordinates
- Encode both coefficients in a single complex number

Complex Fourier transform coefficients

Complex Fourier transform coefficients

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

Complex Fourier transform coefficients

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

Complex Fourier transform coefficients

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

Complex Fourier transform coefficients

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$


$$d_f = \max_{\varphi \in [0,1)} \left(\int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$


Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$




Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$




Complex Fourier transform coefficients

$$c_f = \boxed{\frac{d_f}{\sqrt{2}}} \cdot e^{-i2\pi\varphi_f}$$




Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$




Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{\boxed{-}i2\pi\varphi_f}$$




Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{i2\pi\varphi_f}$$



Complex Fourier transform coefficients

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



Continuous audio signal


$$g(t)$$

Continuous audio signal

$$g(t) \quad g : \mathbb{R} \rightarrow \mathbb{R}$$

Continuous audio signal

$$g(t) \quad g : \mathbb{R} \rightarrow \mathbb{R}$$



Complex Fourier transform

$$\hat{g}(f) = c_f$$

Complex Fourier transform


$$\hat{g}(f) = c_f$$

$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$

Complex Fourier transform

$$\hat{g}(f) = c_f$$


$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$



Complex Fourier transform

$$\hat{g}(f) = c_f$$


$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$



Complex Fourier transform

$$\hat{g}(f) = c_f$$

$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$



Complex Fourier transform

$$d_f = \max_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

Complex Fourier transform


$$d_f = \max_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

Complex Fourier transform


$$d_f = \max_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left(\int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$



Complex Fourier transform

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$



Complex Fourier transform

$$\hat{g}(f) = \int \boxed{g(t)} \cdot e^{-i2\pi ft} dt$$

Complex Fourier transform

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

Complex Fourier transform

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

Complex Fourier transform

$$e^{i\gamma} = \cos(\gamma) + i \sin(\gamma)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

Complex Fourier transform

$$e^{i\gamma} = \cos(\gamma) + i \sin(\gamma)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt = \int g(t) \cdot \cos(-2\pi ft) dt + i \int g(t) \cdot \sin(-2\pi ft) dt$$

Complex Fourier transform

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt = \overset{\text{Real part}}{\int g(t) \cdot \cos(-2\pi ft) dt} + i \overset{\text{Imaginary part}}{\int g(t) \cdot \sin(-2\pi ft) dt}$$

Magnitude Fourier transform

$$|\hat{g}(f)|$$

Magnitude and phase

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

Magnitude and phase

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

Magnitude and phase

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

Magnitude and phase

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

$$\varphi_f = -\frac{\gamma_f}{2\pi}$$


Magnitude and phase

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$


$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

$$\varphi_f = -\frac{\gamma_f}{2\pi}$$

Inverse Fourier transform



IFT
←



Fourier representation

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$

Fourier representation


Pure tone of frequency f

$$g(t) = \int c_f \cdot \boxed{e^{i2\pi ft}} df$$

Fourier representation

Pure tone of frequency f


$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$



Fourier representation

Weight pure tone with magnitude and add phase


$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$



Fourier representation

Add up all (weighted) sinusoids

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$



A Fourier roundtrip

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$



