

1. What is the energy and momentum of photon of a LASER beam of wavelength 6328 Angstrom? ($h = 6.6 \times 10^{-34} \text{ J}\cdot\text{s}$, $c = 3 \times 10^8 \text{ m/s}$)

- A. $3.128 \times 10^{-19} \text{ J}$ and $1.04 \times 10^{-27} \text{ kg}\cdot\text{m/Sec}$**
- B. $31.28 \times 10^{-19} \text{ J}$ and $10.44 \times 10^{-27} \text{ kg}\cdot\text{m/Sec}$**
- C. $5.65 \times 10^{-16} \text{ J}$ and $4.454 \times 10^{-30} \text{ kg}\cdot\text{m/Sec}$**
- D. $55.128 \times 10^{-16} \text{ J}$ and $4.454 \times 10^{-30} \text{ kg}\cdot\text{m/Sec}$**

1. What is the energy and momentum of photon of a LASER beam of wavelength 6328 Angstrom? ($h = 6.6 \times 10^{-34} \text{ J s}$, $c = 3 \times 10^8 \text{ m/s}$)

A. $3.128 \times 10^{-19} \text{ J}$ and $1.04 \times 10^{-27} \text{ kg.m/Sec}$

B. $31.28 \times 10^{-19} \text{ J}$ and $10.44 \times 10^{-27} \text{ kg.m/Sec}$

C. $5.65 \times 10^{-16} \text{ J}$ and $4.454 \times 10^{-30} \text{ kg.m/Sec}$

D. $55.128 \times 10^{-16} \text{ J}$ and $4.454 \times 10^{-30} \text{ kg.m/Sec}$

Given $\lambda = 6328 \times 10^{-10} \text{ m}$, $h = 6.63 \times 10^{-34} \text{ J K sec.}$ and $c = 3 \times 10^8 \text{ m/sec.}$

Formula used $E = h\nu = \frac{hc}{\lambda}$

$$= \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{6.328 \times 10^{-7} \text{ m}} = 1.05 \times 10^{-19} \text{ Joule}$$

~~$E = 3.112 \text{ Joule}$~~

Momentum $p = \frac{E}{c} = \frac{h}{\lambda} = \frac{6.63 \times 10^{-34}}{6.328 \times 10^{-7}} = 1.05 \times 10^{-27} \text{ kg.m/sec}$

~~$p = 1.05 \text{ kg.m/sec.}$~~

2. What is the energy of a Ruby LASER pulse, if the wavelength of the emitted radiation and the number of Cr^{3+} ions are 6943 Angstrom and 2.8×10^{19} respectively?

A. 11.24 J

B. 10.69 J

C. 7.98 J

D. 5.66 J

2. What is the energy of a Ruby LASER pulse, if the wavelength of the emitted radiation and the number of Cr^{3+} ions are 6943 Angstrom and 2.8×10^{19} respectively?

A.11.24 J

B.10.69 J

C.7.98 J

D.5.66 J

Ques :- Calculate the energy of laser pulse in a ruby laser for 2.8×10^{19} Cr^{3+} ions. If the laser emits radiation of wavelength 6943 Å.

Ans :- Given: $\lambda = 6943 \times 10^{-10}$ m, $n = 2.8 \times 10^{19}$

The energy of a photon, $= h\nu$

and the total energy due to n Cr^{3+} ions is

$$E = nh\nu = n \frac{hc}{\lambda} = 2.8 \times 10^{19} \cdot \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{6.943 \times 10^{-7}}$$
$$= 8.02 \text{ J}$$

3. A three-level laser emits a light of wavelength of 5500 Å, What will be the ratio of population of upper level (E_2) to the lower energy level (E_1) if the optical pumping mechanism is shut off (Assume $T = 300$ K)?

A. 1.3×10^{-48}

B. 1.3×10^{-28}

C. 1.3×10^{-18}

D. 1.3×10^{-38}

3. A three-level laser emits a light of wavelength of 5500 Å, What will be the ratio of population of upper level (E_2) to the lower energy level (E_1) if the optical pumping mechanism is shut off (Assume $T = 300$ K)?

A. 1.3×10^{-48}

B. 1.3×10^{-28}

C. 1.3×10^{-18}

D. 1.3×10^{-38}

Given $\lambda = 5500 \text{ Å}$

Formula used is

$$\begin{aligned} E_2 - E_1 &= h\nu = \frac{hc}{\lambda} \\ &= \frac{(6.63 \times 10^{-34} \text{ J/sec}) \times (3 \times 10^8 \text{ m/sec})}{(5.5 \times 10^{-7} \text{ m}) \times (1.6 \times 10^{-19} \text{ J/eV})} \\ &= 2.26 \text{ eV} \end{aligned}$$

and kT can be calculated as

$$\begin{aligned} kT &= (8.62 \times 10^{-5} \text{ eV/K}) \times (300 \text{ K}) \\ &= 0.0259 \text{ eV} \end{aligned}$$

The ratio of upper to the lower energy levels i.e.,

$$\begin{aligned} \frac{E_2}{E_1} &= e^{(-E_2 - E_1)/kT} = e^{-2.26/0.0259} \\ &= e^{-87.3} \\ \frac{E_2}{E_1} &= 1.3 \times 10^{-38} \end{aligned}$$

4. At what temperature the ratio in the previous question (Q: 3) would be $\frac{1}{2}$?

A. 57842 K

B. 27854 K

C. 37832 K

D. 47854 K

4. At what temperature the ratio in the previous question (Q: 3) would be 1/2?

A. 57842 K

B. 27854 K

C. 37832 K

D. 47854 K

$$\frac{E_2}{E_1} = \frac{1}{2} = e^{-(E_2 - E_1)/kT}$$

or

$$e^{(E_2 - E_1)/kT} = 2 \quad \text{or} \quad \frac{E_2 - E_1}{kT} = \log_e 2$$

or

$$T = \frac{E_2 - E_1}{K \log_e 2} = \frac{2.26 \text{ eV}}{\left(8.62 \times 10^{-5} \frac{\text{eV}}{\text{K}}\right) \times (0.693)} = 37832.75 \text{ K}$$

or

$$T = 37832 \text{ K}$$

This temperature is much hotter than the sun.

5. Calculate the power per unit area delivered by a laser pulse of energy 4×10^{-3} Joule, the pulse length in time as 10^{-9} sec and when the pulse is focused on target to a very small spot of radius 1.5×10^{-5} m.

- A. $5.7 \times 10^{20} \text{ W.m}^{-2}$
- B. $5.7 \times 10^{15} \text{ W.m}^{-2}$
- C. $5.7 \times 10^{10} \text{ W.m}^{-2}$
- D. $5.7 \times 10^5 \text{ W.m}^{-2}$

5. Calculate the power per unit area delivered by a laser pulse of energy 4×10^{-3} Joule, the pulse length in time as 10^{-9} sec and when the pulse is focused on target to a very small spot of radius 1.5×10^{-5} m.

- A. $5.7 \times 10^{20} \text{ W.m}^{-2}$
- B. $5.7 \times 10^{15} \text{ W.m}^{-2}$
- C. $5.7 \times 10^{10} \text{ W.m}^{-2}$
- D. $5.7 \times 10^5 \text{ W.m}^{-2}$

Given $P = 4.0 \times 10^{-3} \text{ J}$, $r = 1.5 \times 10^{-5} \text{ m}$

Formula used for power delivered per unit area is given by

$$I = \frac{P}{A}, \text{ where } P = \frac{4.0 \times 10^{-3} \text{ J}}{10^{-9} \text{ sec.}}$$

or

$$P = 4.0 \times 10^6 \text{ W}$$



and $A = \pi r^2 = 3.14 \times (1.5 \times 10^{-5})^2 = 7.065 \times 10^{-10} \text{ m}^2$

so

$$I = \frac{P}{A} = \frac{4.0 \times 10^6 \text{ W}}{7.065 \times 10^{-10} \text{ m}^2} = 5.7 \times 10^{15} \text{ W/m}^2$$

or

$$I = 5.7 \times 10^{15} \text{ W/m}^2$$