Evolution and Reinforcement Learning in Changing Environments

Abhinav Guntaka

Abstract

We compare floor-supported ecological neuroevolution (OE), evolution strategies (ES) and proximal policy optimization (PPO) in a simulated foraging world. Each method has separately tuned fixed- and varied-environment variants, with ten training runs per variant and a shared nominal agent-step budget. We measure food acquisition before and during adaptation to thirty predefined test environments, and population persistence in a separate assay. OE performs below PPO with domain randomization: on the twenty shifted test environments, the difference in pooled interquartile mean of normalized adaptation area under the curve is -0.5592, with a nominal 95% bootstrap interval of $[-0.6088, -0.5196]$. Varied OE also scores below its fixed variant. A later ES-fixed extension finds a positive ES contrast of 0.0286, below the prespecified 0.05 margin; the predicted PPO-DR advantage over PPO-fixed is unsupported under its original rule. OE relies heavily on population-floor replacements, particularly in the resource-scarce test suite. Separate tuning prevents the fixed/varied comparisons from isolating the effect of environmental variation. Objectives and population structure also differ between methods.

1 Introduction

Milano et al. (2019) found that environmental variation across generations improved evolved controllers, with the best performance at a moderate rate of change. Ecological neuroevolution imposes a different selection pressure: agents must acquire enough food to survive and reproduce. Food acquisition also determines task performance, but selection through survival may favor different controllers from direct optimization of food intake.

We compare ecological neuroevolution (OE), evolution strategies (ES) and proximal policy optimization (PPO) in a continuous two-dimensional foraging world. OE maintains an interacting population with energy-dependent death and reproduction, supplemented by a minimum population of externally supplied replacements. ES and PPO optimize food acquisition and deploy copies of a single controller. Each method has separately tuned fixed- and varied-environment configurations. We evaluate food acquisition before and during adaptation, then measure population persistence in a separate assay without replacements.

We predicted that varied-environment OE would outperform PPO with domain randomization during adaptation. OE instead starts below PPO-DR and remains below it throughout the measured interval. Varied OE also scores below fixed OE; a later ES comparison favors the varied configuration by a smaller amount. Because the fixed and varied configurations were tuned separately, these comparisons do not isolate environmental variation.

2 Related work

Milano et al. (2019) varied the rate of environmental change during evolution. Milano and Nolfi (2024)

compared ES and PPO across tasks and treatments of environmental variation. Our experiment compares separately selected fixed- and varied-training configurations in a foraging task, using one variation schedule.

Hamon et al. (2023) studied persistent neural-agent populations with physiology-dependent reproduction and death. Nisioti and Moulin-Frier (2022) used a minimal ecological model and ablations to distinguish fitness-based selection from niche-limited competition. Our implementation also couples survival and reproduction to agent behavior, but uses feedforward controllers and externally supplied population replacements.

Nisioti et al. (2025) compared neuroevolution and reinforcement learning across curricula and found that the outcome depended on the method and task. Nisioti et al. (2026) examine adaptation and forgetting under sequential task changes. Ge and Cheng (2026) compare update and inheritance regimes through nested ecological evaluations of frozen controllers. Both studies are preprints. In our experiment, adaptation begins independently from a trained snapshot in each test environment and permits further parameter updates.

XLand and PAIRED adapt task distributions during learning (Open Ended Learning Team et al., 2021; Dennis et al., 2020); our varied-training configurations sample from a fixed distribution. DERL and Evolution Gym combine morphological and controller optimization (Gupta et al., 2021; Bhatia et al., 2021). All neural agents in the present experiment share a fixed body.

3 Experimental design

3.1 Foraging environment and controllers

Agents move in a square arena with walls and regenerating food. Their observations contain 30 values: eight directional sectors for each of food, other agents and walls; current energy; the previous two-dimensional action; and three body traits. A feedforward controller has two hidden layers of 32 tanh units and two outputs, controlling turn and forward speed. Every method uses this actor architecture. Maximum speed, size and sensing range are fixed at 1, 1 and 10, respectively. No method evolves morphology in this experiment.

Energy decreases with time, movement and sensing. A food item supplies 40 energy units; an agent starts with 100 and dies when its energy reaches zero. In ecological mode, agents can reproduce above an energy threshold, paying an energy cost for each offspring; they also die at a maximum age. Appendix A specifies these rules and the recorded settings. Episodic competence evaluation uses the same movement, sensing and food dynamics, without reproduction.

Varied training samples arena size, food supply, layout and regrowth from the ranges in Table 1. The fixed environment has side length 60, 40 food items, a uniform layout and regrowth probability 0.03. Validation uses 20 sampled training-range environments and four corner cases. Test conditions are fixed in advance of final evaluation: ten training-range environments (T1), ten smaller and resource-scarce clustered environments (T2), and ten environments with food drift and seasonal relocation (T3). T3 retains the training ranges for the remaining parameters. T2 tests resource scarcity; T3 tests changing food locations.

3.2 Training procedures

Ecological neuroevolution. OE starts with 32 agents and permits at most 128. Each agent has its own controller. Offspring inherit the parent's weights with Gaussian perturbations; natural death

Table 1: Training and fixed test suites. Each test suite contains ten environments. The canonical fixed-training setting is given in the text. Unused layout parameters have no effect in a given environment.

Parameter	Varied training / T1	T2	T3
Arena side L	40–80	30–40	40–80
Food items F	20–60	10–20	20–60
Layouts	Uniform, clusters, patches	Clusters	Uniform, clusters, patches
Cluster count k	2–5	2–3	2–5
Cluster scale σ	3–8	2–4	3–8
Patch radius	4–10	Not used	4–10
Regrowth probability	0.01–0.05	0.005–0.01	0.01–0.05
Food drift	0	0	0.05–0.3
Season period (steps)	None	None	100–300

and reproduction determine which lineages persist. In varied OE, the environmental parameters change every 2,000 world steps while the population continues. OE-fixed retains the canonical environment. During adaptation, either variant continues evolving in the assigned test environment.

Both variants include a population floor of 16. After a step, vacant slots are replenished with mutated offspring of uniformly sampled surviving, non-newborn agents; if that pool is empty, agents alive at the start of the step supply the parent pool. Replacements receive birth energy and a new position. Replacements can maintain the population when natural reproduction is insufficient. We count natural and floor births separately.

Evolution strategies. ES optimizes the shared actor weights using the `Open_ES` implementation in evosax (Salimans et al., 2017; Lange, 2023). It uses antithetic perturbations and centered-rank fitness shaping, with candidate fitness determined by episodic food acquisition. Candidates in a generation share evaluation conditions. ES-DR samples training environments; ES-fixed uses the canonical environment. Evaluation uses the strategy mean. ES-fixed was added after inspecting the core results, under a separate protocol written before its own tuning and final runs.

Each candidate is evaluated with 16 policy copies in 500-step worlds without births. Its score is food per initial agent, averaged over its configured episodes. Candidates in a generation share environment draws and reset/step random keys; ES-DR redraws the episode environments each generation.

Proximal policy optimization. PPO uses a clipped policy-gradient objective (Schulman et al., 2017), a separate value network, and a parameter-shared policy across agents in 64 parallel episodic worlds. Food acquisition supplies the reward. PPO-DR samples environment parameters at episode resets; PPO-fixed uses the canonical environment. Training actions are stochastic, whereas competence evaluation uses the actor's deterministic output. The implementation derives from PureJaxRL's continuous-action PPO, with the changes and software provenance recorded in Appendix A.

Each world contains 16 agents and restarts at 500 steps or when all agents die. An agent's death terminates its trajectory. Steps beginning with a dead agent are excluded from losses and advantage normalization; when enabled, reward-normalization statistics are collected over living agents.

OE maintains a heterogeneous population; ES and PPO deploy copies of one policy. Table 2 summarizes their deployment and transfer operations.

Table 2: Training and transfer procedures. Each row has fixed and varied variants, separately tuned. Body traits and actor architecture are shared.

Procedure	Training signal	Competence deployment	State at adaptation start
OE	Energy-dependent survival/reproduction plus floor replacements	16 sampled population members	Genomes/alive mask retained; world state and energy reset
ES	Rank-shaped episodic food score	16 copies of mean actor	Mean solution retained; fresh strategy/optimizer
PPO	Food reward and clipped policy objective	16 copies of actor	Actor/critic and reward statistics retained; fresh Adam/rollout state

3.3 Tuning, budgets and evaluation

Eight random-search trials were run separately for each variant, each with a distinct tuning seed and 5×10^7 nominal living-agent steps. The selection score was mean zero-shot food acquisition above the random-policy score across the 24 validation environments. Tuning therefore used a different endpoint from the primary adaptation AUC. The selected configuration was then trained with ten final seeds, 5000–5009, using 2×10^8 nominal living-agent steps per run. All ten final runs per variant are retained. Settings and search ranges appear in Appendix A.

One living-agent step counts an agent alive at the beginning of a simulated step. Wall time, floating-point operations, optimizer updates and checkpoint positions can differ at the same nominal agent-step budget. Updates finish in indivisible units and may cross a stopping target. We analyze both the prespecified nominal grid and the actual step counts.

Competence is total food consumed divided by the 16 initial agents in a 500-step episode, averaged across eight evaluation episodes. OE samples population members afresh for each episode, without replacement when at least 16 are alive; ES and PPO use 16 policy copies. Episode and reset seeds are shared across methods, although the random-action reference does not share every subsequent world-step random draw with learned policies.

Adaptation starts independently from the final trained snapshot in each test environment and receives 2×10^7 nominal living-agent steps. There are eleven competence measurements: the zero-shot score and ten subsequent checkpoints. OE retains genomes and the alive mask while resetting physical state and energy. PPO retains actor/critic parameters and reward-normalization statistics but resets Adam and rollout state. ES retains its mean solution with fresh optimizer and strategy state.

Persistence is evaluated separately, before adaptation. Thirty-two sampled OE agents or policy copies enter ecological mode for 5,000 world steps, with reproduction but no mutation and no population floor. We record extinction and steps survived.

3.4 Outcomes and statistical analysis

The analyses were specified in version-controlled protocols before the corresponding runs. These protocols were private during the study. The ES-fixed extension was specified after the core outcomes were known but before its own runs. Appendix E records this chronology.

Let $c_{mre}(t)$ be competence for method m, run r and environment e at normalized adaptation

checkpoint $t \in \{0, 0.1, \ldots, 1\}$. We use the prespecified transformation

$$z_{mre}(t) = \frac{c_{mre}(t) - q_e}{b_e - q_e}, \qquad A_{mre} = \frac{1}{10} \sum_{j=1}^{10} \frac{z_{mre}(t_{j-1}) + z_{mre}(t_j)}{2}, \tag{1}$$

where q_e is the random-policy score and b_e is the largest observed score across the five core methods, all final runs and all eleven checkpoints in that environment. No environment had $b_e \leq q_e$, so none was excluded. The later ES-fixed analysis uses the saved core anchors. The scores are relative to this observed reference.

The primary effect is

$$D = \text{IQM}\{A_{\text{OE},r,e} : e \in T2 \cup T3\} - \text{IQM}\{A_{\text{PPO-DR},r,e} : e \in T2 \cup T3\}, \tag{2}$$

where the interquartile mean (IQM) averages the middle half of the pooled run–environment values. We resample complete runs, keeping each run's environment outcomes together, following the principle that repeated evaluation of a trained agent does not create new independent training runs (Agarwal et al., 2021; Henderson et al., 2018). There are 50,000 bootstrap draws with statistics seed 4800. The prespecified margin is 0.05 on this reference scale. AUC measures performance over the adaptation interval, including starting competence.

The seven core secondary comparisons use two-sided Mann–Whitney tests on per-run environment means, with Holm adjustment within the family. Their companion IQM effects have a different statistical target from the rank tests. The later A2–A3 comparisons form a separate Holm family. H1 and A1 use their interval-based margin rules; A1 and A2 describe the same underlying ES AUC contrast. Appendix B reports the planned core comparisons and later addendum comparisons, including the four-method summaries H2b and A4.

All reported neural intervals are nominal percentile bootstrap intervals. The calculation freezes the observed normalization anchors and test environments; it does not propagate uncertainty in anchor selection, environment/episode sampling or selection of tuning configurations. Because the maximum anchors are selected from these same runs, formal conditional 95% coverage is not established. Final seeds are independent repetitions within each method, but matching seed streams may couple different methods. We examine paired-seed resampling and re-estimation of the anchors in post hoc sensitivity analyses. Ten runs provide limited precision, especially for equivalence claims.

4 Results

4.1 OE scores below PPO-DR during adaptation

OE performs below PPO-DR, contrary to the primary prediction. Across T2 and T3, OE has a pooled AUC IQM of 0.2905, compared with 0.8497 for PPO-DR. Their difference is -0.5592 with a nominal 95% interval of $[-0.6088, -0.5196]$, entirely below the negative margin. OE also scores below ES-DR in the prespecified S1 comparison: the IQM difference is -0.5871 and the Holm-adjusted rank-test p value is 0.00110. Figure 1 shows estimates for all six procedures. The prespecified comparisons are reported in Appendix B.

OE is already below PPO-DR before adaptation. The zero-shot IQM difference on T2/T3 is -0.4648; at the final checkpoint it is -0.6035 (S4 and S5, both Holm-adjusted $p = 0.00110$). The corresponding mean trajectories separate the initial level from subsequent change (Figure 3). In T2,




Figure 1: Performance during adaptation on T2 and T3. Points show pooled IQMs of normalized AUC; bars are nominal 95% whole-run percentile bootstrap intervals from 50,000 draws. Each procedure has ten training runs evaluated on twenty fixed environments. The empirical normalization anchors remain fixed during resampling (Section 3.4). ES-fixed, marked with an asterisk, is the later extension.

OE's mean normalized competence declines from 0.4383 to 0.2338, while PPO-DR changes from 0.7244 to 0.7478. T3 shows a smaller decline for OE, from 0.2966 to 0.2655. These mean trajectories summarize changes from different initial policies.

The deficit is also present on the in-range T1 suite. S6 compares T1 adaptation AUC and gives an OE–PPO-DR IQM difference of -0.5371 with interval $[-0.5951, -0.4799]$.

4.2 Fixed and varied training show different patterns across procedures

The separately tuned varied OE procedure scores below OE-fixed on T2/T3 AUC: difference -0.1210, interval $[-0.1799, -0.0680]$ (S2; Holm-adjusted $p = 0.00440$). For PPO, the corresponding difference is 0.0099, interval $[-0.0004, 0.0201]$. Its predicted direction is unsupported under the prespecified rank-test rule (S3; adjusted $p = 0.6232$). The analysis provides no evidence of equivalence.

The later ES-fixed extension gives an ES-DR–ES-fixed AUC difference of 0.0286, interval $[0.0203, 0.0384]$. The interval lies above zero and below the 0.05 margin, yielding a small positive contrast under the A1 rule. A2, the rank-test summary of the same contrast, has adjusted $p = 0.000365$. ES-DR also has higher zero-shot competence on T1: A3 gives 0.0645, interval $[0.0499, 0.0788]$, with adjusted $p = 0.000365$. That interval is not wholly above 0.05. A1 and A3 use different environment sets as well as different endpoints.

The fixed/varied comparisons in Figure 2 also differ in selected hyperparameters. For example, OE's mutation standard deviation is 0.03412 under varied training and 0.00507 under fixed training. The contrast consequently includes the effects of both training regime and hyperparameter selection.




Figure 2: Fixed and varied configurations. Upper panels: T2/T3 AUC, with individual run means (circles), pooled IQMs (diamonds) and nominal prespecified 95% whole-run intervals. Lower panels: T1 zero-shot run means and their mean (squares), with descriptive 95% whole-run intervals from 20,000 draws. The A3 annotation is a pooled-IQM contrast, not the difference between the mean squares. All resampling freezes environments and empirical anchors. Lower OE/PPO panels are descriptive. The endpoints differ in environment set, and all fixed/varied pairs were separately tuned. Asterisk: later ES-fixed extension.

4.3 Population persistence and support

OE has higher extinction in the separate persistence assay than PPO-DR. Across T2/T3, the S7 difference in IQMs of per-run extinction fractions is 0.3750, interval $[0.2500, 0.4833]$, against the predicted OE advantage (adjusted $p = 0.00110$). Table 3 reports the descriptive extinction frequencies for every procedure. Extinction is frequent across all six procedures in resource-scarce T2, including those with high normalized competence. High relative food acquisition can coexist with insufficient intake for population survival.

The population floor makes a substantial contribution during OE training and adaptation. The mean run-level share of training births supplied by the floor is approximately 58% for OE and 59% for OE-fixed. During T2 adaptation, the pooled shares rise to 99.53% and 99.60%, respectively (Table 4). The population is therefore almost entirely replenished through floor births in T2. The persistence assay evaluates these populations after that support is removed.

Table 3: Descriptive extinction frequencies in the separate 5,000-step no-floor assay. Each cell pools ten runs across ten fixed environments; these 100 observations are not independent training runs.

Procedure	T1	T2	T3
OE	0.48	0.94	0.76
OE-fixed	0.30	0.93	0.48
PPO-DR	0.03	0.83	0.15
PPO-fixed	0.10	0.95	0.14
ES-DR	0.03	0.89	0.22
ES-fixed*	0.11	0.87	0.21

Table 4: Share of new agents supplied by population-floor replacements during adaptation. Fractions pool event counts over ten runs and ten environments per suite.

Suite	OE	OE-fixed
T1	64.35%	59.29%
T2	99.53%	99.60%
T3	78.90%	71.87%

4.4 Recorded learning and sensitivity checks

Training-validation records begin near 20 million steps. Between that observation and 200 million steps, OE's mean validation score above random changes from 9.4818 to 10.6102; six of ten runs increase. No initialization measurement is available. Figure 3 shows the full saved training and adaptation trajectories. A later exploratory extension trained three fresh OE runs to one billion steps (Appendix C).

The primary contrast is stable across the sensitivity analyses. Re-estimating the core best anchors within each bootstrap draw gives an H1 interval of $[-0.6105, -0.5215]$; resampling corresponding seeds together gives $[-0.6101, -0.5185]$. Integrating piecewise-linear saved curves against actual steps, truncated at the common 20-million-step endpoint, changes the H1 estimate to -0.559203. The analogous ES AUC estimate is 0.028921, interval $[0.020572, 0.038833]$, still below the margin. The intervals remain approximate at this sample size.

One sensitivity changes the interpretation of the weak PPO contrast: the paired IQM interval for S3 is $[0.0020, 0.0177]$, above zero. The prespecified decision remains unsupported under its rank test on run means. The paired IQM interval uses both a different outcome summary and a different treatment of cross-method dependence.

4.5 Separate supporting trait experiment

A trait-only experiment preceded the neural comparison and examined food-versus-population search, mutation and cloning under related foraging shifts. It used four environmental arms and 200 seeds per group per arm to compare evolving populations, frozen search-selected designs, populations with standing variation, mutation freezes and cloning. The model and its complete 31-pair result record are reported in Appendix D; its agents use inherited traits and fixed behavioral rules.

In its main arm, a frozen population-search design reaches the population ceiling, while the evolving population is close to it. Their mean population difference is 0.6375 against a 2.5-agent




Figure 3: Training observations and adaptation trajectories. Upper panels show validation competence above the random reference at saved training checkpoints. Lower panels show mean normalized test competence during adaptation. Lines average ten runs; shading gives descriptive, pointwise 95% whole-run percentile intervals from 20,000 draws, not simultaneous bands. Axes use nominal living-agent steps. The first training observation is near 20 million steps; no initialization score is available. For normalized adaptation scores, resampling freezes the selected anchors and test environments. ES-fixed is the later extension.

margin, and neither group becomes extinct. The population-search procedure also substantially outperforms the food-search procedure, but it uses both a different objective and a longer scoring horizon. Cloning reduces population relative to the frozen diverse population after the move, and a larger deficit appears in a no-move control. The changes in scoring horizon and the no-move clone deficit limit explanations based solely on objective or environmental shift.

5 Discussion

The tested OE procedure scores below PPO-DR before and during adaptation, in both the training-range and shifted environments. Its populations also have poorer persistence in the separate assay without floor support.

OE selects an interacting population through energy and reproduction and supplies replacement agents when the population falls below its floor. ES and PPO optimize an episodic food score and deploy copies of one actor. Each configuration was tuned separately on zero-shot validation performance. Selection objective, population structure, replacement support and tuning could each contribute to the performance difference. Separating their effects would require experiments that change one component at a time.

The ES zero-shot result on T1 (A3) is consistent with the benefit of varied training reported by

Milano et al. (2019) in a different task. The ES adaptation contrast (A1) is smaller and concerns the shifted suites. Both results come from one variation schedule and separately tuned configurations. Our experiment cannot test the moderate-rate optimum found in their rate sweep, or attribute the contrasting OE result to selection type.

In T2, where food is scarce and regrowth is slow, extinction is frequent even among policies with relatively high normalized food scores. Those scores express performance relative to an empirical reference; survival requires sufficient food to cover the population's energy costs. Floor replacements supply additional energy during OE training and adaptation, whereas the persistence assay removes that support.

There are ten training runs per configuration and thirty fixed test environments. The bootstrap holds the empirical normalization anchors fixed and excludes variation from tuning and test-environment sampling. The primary contrast persists under alternative resampling and step-count calculations, but these checks use the same observations. Generalization to other environments and tuning searches remains untested. Training records begin near 20 million steps, so they do not establish improvement from initialization. The three exploratory longer-training runs provide little evidence about performance at the larger budget.

6 Conclusion

Under the tested configurations and nominal agent-step budgets, floor-supported ecological neuroevolution performs below PPO-DR during adaptation and has poorer persistence without floor support. The varied OE configuration scores below OE-fixed, while ES-DR scores slightly above ES-fixed. The predicted advantage of PPO-DR over PPO-fixed is unsupported under its prespecified rule. These findings compare separately tuned procedures in one foraging world; they do not identify the causes of the performance differences.

Data and code availability

The manuscript, implementation, configurations, analysis scripts and data are available at `https://github.com/Guntikiwritescode/ecological-neuroevolution`. Appendix E describes the archived records and source revisions.

Author contributions

Abhinav Guntaka designed the study, implemented the software, ran the experiments, analyzed the results and wrote the manuscript.

Funding and competing interests

The author funded this work. Total project expenditure was US$87. The author declares no competing interests.

Use of AI tools

Claude Code and Cursor Agents assisted with the research. ChatGPT/Codex assisted with code inspection, statistical reanalysis, manuscript drafting and editing. The author is responsible for the scientific content and conclusions.

References

Rishabh Agarwal, Max Schwarzer, Pablo Samuel Castro, Aaron Courville, and Marc G. Bellemare. Deep Reinforcement Learning at the Edge of the Statistical Precipice. In *Advances in Neural Information Processing Systems*, volume 34, 2021. URL `https://proceedings.neurips.cc/paper_files/paper/2021/hash/f514cec81cb148559cf475e7426eed5e-Abstract.html`.

Jagdeep Singh Bhatia, Holly Jackson, Yunsheng Tian, Jie Xu, and Wojciech Matusik. Evolution Gym: A Large-Scale Benchmark for Evolving Soft Robots. In *Advances in Neural Information Processing Systems*, volume 34, 2021. URL `https://proceedings.neurips.cc/paper_files/paper/2021/hash/118921efba23fc329e6560b27861f0c2-Abstract.html`.

Michael Dennis, Natasha Jaques, Eugene Vinitsky, Alexandre Bayen, Stuart Russell, Andrew Critch, and Sergey Levine. Emergent Complexity and Zero-shot Transfer via Unsupervised Environment Design. In *Advances in Neural Information Processing Systems*, volume 33, 2020. URL `https://proceedings.nips.cc/paper_files/paper/2020/hash/985e9a46e10005356bbaf194249f6856-Abstract.html`.

Yuxu Ge and Yifei Cheng. Neuroevolution arena: Nested ecological evaluation of update-and-inheritance regimes across neural architectures. arXiv:2608.10323v1, 2026. URL `https://arxiv.org/abs/2608.10323`. Preprint, 10 August 2026.

Agrim Gupta, Silvio Savarese, Surya Ganguli, and Li Fei-Fei. Embodied intelligence via learning and evolution. *Nature Communications*, 12(1):5721, 2021. doi: 10.1038/s41467-021-25874-z.

Gautier Hamon, Eleni Nisioti, and Clément Moulin-Frier. Eco-evolutionary Dynamics of Non-episodic Neuroevolution in Large Multi-agent Environments. In *Proceedings of the Companion Conference on Genetic and Evolutionary Computation*, pages 143–146. ACM, 2023. doi: 10.1145/3583133.3590703.

Peter Henderson, Riashat Islam, Philip Bachman, Joelle Pineau, Doina Precup, and David Meger. Deep Reinforcement Learning that Matters. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 32, 2018. URL `https://doi.org/10.1609/aaai.v32i1.11694`.

Robert Tjarko Lange. evosax: JAX-Based Evolution Strategies. In *Proceedings of the Companion Conference on Genetic and Evolutionary Computation*, pages 659–662. ACM, 2023. doi: 10.1145/3583133.3590733.

Nicola Milano and Stefano Nolfi. Qualitative differences between evolutionary strategies and reinforcement learning methods for control of autonomous agents. *Evolutionary Intelligence*, 17 (2):1185–1195, 2024. doi: 10.1007/s12065-022-00801-3.

Nicola Milano, Jônata Tyska Carvalho, and Stefano Nolfi. Moderate Environmental Variation Across Generations Promotes the Evolution of Robust Solutions. *Artificial Life*, 24(4):277–295, 2019. doi: 10.1162/artl_a_00274.

Eleni Nisioti and Clément Moulin-Frier. Plasticity and evolvability under environmental variability: the joint role of fitness-based selection and niche-limited competition. In *Proceedings of the Genetic and Evolutionary Computation Conference*, pages 113–121. ACM, 2022. doi: 10.1145/3512290.3528826.

Eleni Nisioti, Erwan Plantec, Milton Montero, Joachim Pedersen, and Sebastian Risi. When Does Neuroevolution Outcompete Reinforcement Learning in Transfer Learning Tasks? In *Proceedings of the Genetic and Evolutionary Computation Conference*, pages 48–57. Association for Computing Machinery, 2025. doi: 10.1145/3712256.3726475. Author preprint: arXiv:2505.22696.

Eleni Nisioti, Andrea Cossu, Kathrin Korte, and Sebastian Risi. Continual Reinforcement Learning with Neuroevolution. arXiv:2610.01583, 2026.

Open Ended Learning Team, Adam Stooke, Anuj Mahajan, Catarina Barros, Charlie Deck, Jakob Bauer, Jakub Sygnowski, Maja Trebacz, Max Jaderberg, Michael Mathieu, Nat McAleese, Nathalie Bradley-Schmieg, Nathaniel Wong, Nicolas Porcel, Roberta Raileanu, Steph Hughes-Fitt, Valentin Dalibard, and Wojciech Marian Czarnecki. Open-Ended Learning Leads to Generally Capable Agents. arXiv:2107.12808, 2021.

Jasper Palfree. Evolution simulator (MinuteLabs.io). `https://github.com/minutelabsio/evolution-simulator`, commit 55af956, 2021. GPL-3.0.

Tim Salimans, Jonathan Ho, Xi Chen, Szymon Sidor, and Ilya Sutskever. Evolution Strategies as a Scalable Alternative to Reinforcement Learning. arXiv:1703.03864, 2017.

John Schulman, Filip Wolski, Prafulla Dhariwal, Alec Radford, and Oleg Klimov. Proximal Policy Optimization Algorithms. arXiv:1707.06347, 2017.

A Implementation and selected settings

The actor has dimensions $30 \to 32 \to 32 \to 2$, with tanh hidden activations and 2,114 parameters. Initial weights are orthogonal, with gain $\sqrt{2}$ in hidden layers and 0.01 in the action layer; biases start at zero. The world maps the outputs through tanh to a turn bounded by $\pi/4$ per step and a speed between zero and the body's maximum. Evaluation actions are deterministic. The observation includes clipped energy divided by birth energy and scaled body traits, even though the latter remain fixed.

The per-step energy cost is $0.5 + s^3v^2 + 0.05r$, where s is size, v is commanded speed after action squashing but before position clipping at a wall, and r is sensing range. With the default body it is $1 + v^2$. Food supplies 40 energy units. Ecological reproduction requires energy at least 200, costs the parent 120 and supplies the offspring with 100. Maximum age is 1,000 steps and the offspring placement offset is 1. The ecological capacity is 128; competence evaluation has 16 agent slots. The floor rule described in Section 3.2 adds energy from outside this budget and is logged separately.

Within a step, agents move, consume reachable food, pay costs, die and reproduce; food dynamics then update. Random priority resolves contested food. Births require free slots, with eligible parents processed in random order, at most one child per parent per step, and payment only for successful births. At a varied-OE environment change, positions are clipped to the new arena as needed, while energy, age and genomes persist. Food is reinitialized under the new parameters.

For PPO, the common settings are 64 parallel worlds, 32 minibatches, GAE parameter 0.95, clipping parameter 0.2, value-loss coefficient 0.5, maximum gradient norm 0.5, initial log standard deviation zero and no learning-rate annealing. The separately selected values are in Table 5. PPO's actor parameters, critic parameters, log standard deviation and reward-normalization state are carried into adaptation; its return accumulator, optimizer, environment state, counters and random stream are reset. OE carries controller genomes and the alive mask, while resetting positions, headings, energy, age, actions and counters. ES carries its mean solution while restarting its optimizer/strategy and generation counter.

The PPO critic has the same two hidden layers as the actor and a scalar output. Action exploration is Gaussian with learned, state-independent log standard deviation. The adapted implementation starts from PureJaxRL commit `31756b1`; ES uses evosax version 0.3.2. Source notices and licenses are retained with the scientific implementation.

Table 5: Selected settings from the recorded tuning outputs, rounded to six significant digits. Exact values are retained in the accompanying tuning CSV. Fixed and varied configurations were selected separately. Asterisk: later ES-fixed extension.

Procedure	Settings
OE	Mutation SD = 0.0341186
OE-fixed	Mutation SD = 0.00507475
PPO-DR	Entropy coefficient = 0; Discount = 0.99; Learning rate = 0.000180353; Reward normalization = on; Rollout length = 64; Update epochs = 8
PPO-fixed	Entropy coefficient = 0; Discount = 0.95; Learning rate = 0.000706297; Reward normalization = off; Rollout length = 128; Update epochs = 8
ES-DR	Episodes per candidate = 1; Learning rate = 0.0073746; Population size = 64; Perturbation SD = 0.089879
ES-fixed*	Episodes per candidate = 4; Learning rate = 0.0117805; Population size = 32; Perturbation SD = 0.0441052

Table 6: Interval-rule comparisons. A1 and A2 use the same ES contrast; they are not independent confirmations. The margin is 0.05 on the empirical normalized scale.

ID	Comparison	Difference [interval]	Decision
H1	OE minus PPO-DR	-0.5592 $[-0.6088, -0.5196]$	Against predicted superiority
A1	ES-DR minus ES-fixed	$+0.0286$ $[0.0203, 0.0384]$	Positive, below margin

The tuning search for PPO sampled learning rate log-uniformly from 10^{-4} to 10^{-3}, rollout length from $\{32, 64, 128\}$, update epochs from $\{2, 4, 8\}$, discount from $\{0.95, 0.99, 0.995\}$, entropy coefficient from $\{0, 0.001, 0.01\}$ and reward normalization on or off. ES sampled perturbation scale from $[0.01, 0.1]$ and learning rate from $[0.003, 0.03]$, both log-uniformly, population size from $\{32, 64, 128\}$ and episodes per candidate from $\{1, 2, 4\}$. OE sampled weight-mutation standard deviation log-uniformly from $[0.005, 0.05]$. Population floor and change interval were design choices, not tuned parameters. Each trial had its own tuning seed; a score tie favored the lower trial index.

Validation environments used seeds 4900–4919 plus four corner cases. Final T1, T2 and T3 environments used seeds 5900–5909, 5910–5919 and 5920–5929. Competence episodes were keyed from environment-specific evaluation seeds and checkpoint indices; the same test conditions were used across methods. Final run seeds were 5000–5009. The implementation separates named random streams by purpose, although common seeds can still couple methods. The exact realized environment parameters, rather than only their generating ranges, are retained with every evaluation record.

B Complete neural comparisons

The original private protocol called H1 a prediction that OE would adapt faster than PPO-DR. Its specified statistic was the AUC in Equation 1. AUC includes the starting score and measures performance during adaptation, not adaptation speed. H1 is against the prediction because its interval lies below -0.05. The analogous later A1 contrast is positive but wholly below 0.05 (Table 6). The margin is a fraction of an empirical reference gap; it has no established interpretation outside this normalization scale.

The H1/A1 rule checks the interval in this order. An interval wholly above zero supports the predicted direction, qualified as below the margin if it also lies below $+0.05$. An interval wholly below zero is against that direction, qualified as below the margin if it also lies above -0.05. If neither condition holds, an interval contained in $[-0.05, +0.05]$ yields the protocol's no-meaningful-difference label; any other interval is inconclusive. Thus directional support is not the same as exceeding the practical margin. These labels inherit the interval-calibration limitations in Section 3.4.

Table 7: Complete core S1–S7 and later A2–A3 families. Differences are pooled IQMs except S7, which uses IQMs of per-run extinction fractions. Intervals are nominal 95% whole-run intervals. Holm correction is separate for the two families. The rank-test p values apply to per-run summaries, not directly to the IQM effects.

ID	Comparison	Outcome	Difference [interval]	Adjusted p
S1	OE minus ES-DR	T2/T3 AUC	-0.5871 $[-0.6364, -0.5475]$	0.0011
S2	OE minus OE-fixed	T2/T3 AUC	-0.1210 $[-0.1799, -0.0680]$	0.0044
S3	PPO-DR minus PPO-fixed	T2/T3 AUC	$+0.0099$ $[-0.0004, 0.0201]$	0.6232
S4	OE minus PPO-DR	T2/T3 zero-shot	-0.4648 $[-0.5570, -0.4073]$	0.0011
S5	OE minus PPO-DR	T2/T3 final	-0.6035 $[-0.6486, -0.5643]$	0.0011
S6	OE minus PPO-DR	T1 AUC	-0.5371 $[-0.5951, -0.4799]$	0.0011
S7	OE minus PPO-DR	T2/T3 extinction	$+0.3750$ $[0.2500, 0.4833]$	0.0011
A2	ES-DR minus ES-fixed*	T2/T3 AUC	$+0.0286$ $[0.0203, 0.0384]$	3.65×10^{-4}
A3	ES-DR minus ES-fixed*	T1 zero-shot	$+0.0645$ $[0.0499, 0.0788]$	3.65×10^{-4}

S1, S2, S5 and S7 are against their predicted OE directions. S3 does not support its predicted PPO direction. S4 and S6 were nondirectional and favor PPO-DR. A2 and A3 support the predicted ES-DR direction under the separate addendum family. The Mann–Whitney probability summary is $U/(n_X n_Y)$, with ties contributing one half; the test is not generally a test of arithmetic mean equality. S7 first reduces each run to its extinction fraction across T2/T3, then compares the distributions of those fractions. It does not treat individual environments as independent training replications.

For H2b, write $q_{ov}, q_{of}, q_{pv}, q_{pf}$ for pooled AUC IQMs of OE, OE-fixed, PPO-DR and PPO-fixed. The prespecified summaries are

$$V = \tfrac{1}{2}[(q_{ov} - q_{of}) + (q_{pv} - q_{pf})], \tag{3}$$

$$M = \tfrac{1}{2}[(q_{ov} - q_{pv}) + (q_{of} - q_{pf})], \tag{4}$$

$$I = (q_{ov} - q_{of}) - (q_{pv} - q_{pf}), \qquad \Delta = |V| - |M|. \tag{5}$$

Table 8 reports the estimates. The original rule labeled the negative Δ interval "method matters more." That label describes these four configurations on this scale. Their interaction is nonzero, and tuning differs, so it is not a general or causal main-effect statement. Absolute-value contrasts are also unstable when a component is close to zero.

A4 compares the ES varied-minus-fixed contrast with the corresponding OE contrast. Its difference is 0.149628, interval $[0.095763, 0.209123]$. A4 is descriptive: the OE outcome was known before the ES-fixed extension. Its component intervals come from the four-method resampling routine and need not exactly equal the intervals from separate two-method routines.

Table 8: Prespecified four-method summaries. These are contrasts among selected configurations, not identified causal main effects. A4 is a later descriptive comparison whose OE component was already known.

Analysis	Quantity	Estimate	Nominal 95% interval
H2b	delta	-0.438162	[-0.474258, -0.405598]
H2b	interaction	-0.130959	[-0.190704, -0.076383]
H2b	method	-0.493721	[-0.524014, -0.467100]
H2b	variety	-0.055559	[-0.085238, -0.028180]
A4	interaction	+0.149628	[0.095763, 0.209123]
A4	variety_es	+0.028590	[0.020276, 0.038461]
A4	variety_oe	-0.121038	[-0.179733, -0.067839]

Table 9: All-method descriptive means. These average environments within runs and then runs; they are distinct from pooled IQMs.

Procedure	Suite	Zero-shot	Final	AUC
OE	T1	0.4052	0.3946	0.3960
OE	T2	0.4383	0.2338	0.3205
OE	T3	0.2966	0.2655	0.2759
OE-fixed	T1	0.5923	0.5088	0.5303
OE-fixed	T2	0.5285	0.2897	0.3860
OE-fixed	T3	0.5441	0.4076	0.4510
PPO-DR	T1	0.9236	0.9235	0.9295
PPO-DR	T2	0.7244	0.7478	0.7611
PPO-DR	T3	0.8981	0.8840	0.8861
PPO-fixed	T1	0.9161	0.9196	0.9216
PPO-fixed	T2	0.7181	0.7123	0.7290
PPO-fixed	T3	0.9202	0.9147	0.9089
ES-DR	T1	0.9017	0.9270	0.9231
ES-DR	T2	0.7114	0.8833	0.8743
ES-DR	T3	0.8543	0.8772	0.8686
ES-fixed*	T1	0.8223	0.9067	0.8930
ES-fixed*	T2	0.6770	0.8656	0.8377
ES-fixed*	T3	0.8041	0.8530	0.8470

C Sensitivity analyses and descriptive diagnostics

Post hoc re-estimation of the core maximum anchor inside each of 50,000 bootstrap draws gives intervals of $[-0.61045, -0.52146]$ for H1, $[-0.17997, -0.06776]$ for S2, $[0.02042, 0.03851]$ for A1 and $[0.04965, 0.07903]$ for A3. Paired resampling of corresponding seeds gives $[-0.61012, -0.51847]$, $[-0.18100, -0.06919]$ and $[0.02039, 0.03964]$ for H1, S2 and A1. S3's paired IQM interval is positive, as reported in the main text. None of these analyses replaces a prespecified test or establishes exact coverage for a selected maximum.

Realized adaptation counts differ between methods. The mean final counts per environment are approximately 20,000,010 for either OE variant, 20,017,188 for PPO-DR, 20,039,595 for PPO-fixed,




Figure 4: Descriptive performance summaries. Left: empirical fraction above each AUC threshold among ten runs across twenty fixed environments; no confidence bands are shown. Right: fraction of the 100 cross-run comparisons in which the row method's mean exceeds the column method's mean, with ties counting one half. These are descriptive effect summaries. ES-fixed is the later extension.

20,144,256 for ES-DR and 20,288,091 for ES-fixed. ES-fixed can overshoot its first two-million-step checkpoint by 49.5%; the largest checkpoint excess is 4.98% of the complete adaptation budget, and the largest final excess is 4.49%. Actual-step sensitivity integrates the piecewise-linear saved observations and truncates at 20 million steps. Scores between saved checkpoints are linearly interpolated.

The primary deficit persists when individual runs or environments are omitted and when raw scores replace normalized scores. Figure 4 shows empirical performance profiles and pairwise run-mean probability summaries.

The time-to-threshold diagnostic compares raw competence against 90% of each run's mean T1 zero-shot score. It therefore depends on the food supply and attainable raw score in the target environment. In the core study, OE and OE-fixed each reach that threshold in 43 of 200 T2/T3 run–environment pairs; PPO-DR, PPO-fixed and ES-DR each reach it in 40. These counts are not independent trials and non-reaching cases cannot be dropped when interpreting adaptation. The threshold is not a general recovery criterion.

Population diversity is descriptive. Weight diversity is mean pairwise Euclidean distance among sixteen sampled neural parameter vectors. Behavior diversity uses action distances on 256 fixed validation probes. Body-trait diversity is zero by design. The final saved OE weight and behavior summaries exceed OE-fixed's, but no diversity intervention in this neural experiment identifies the effect of that difference (Figure 5).

Three fresh OE runs were trained to one billion steps after the core results. Gain compares the final two saved validation checkpoints with the first two. None exceeded a prewritten 4.3-point threshold, whose motivating noise estimate was subsequently found unreliable; this is not a calibrated test. Figure 6 shows their saved trajectories.




Figure 5: Population diagnostics and separate persistence. Weight and behavioral diversity summarize sixteen sampled agents per saved training population, averaged across ten runs. No uncertainty bands are assigned. The right panel shows extinction frequencies in the separate no-floor persistence assay, not extinction during floor-supported training. Body traits remain fixed.




Figure 6: Exploratory longer training. Three fresh OE runs trained to one billion nominal steps, with the ten original 200-million-step runs in gray. Gain is the mean of the final two saved checkpoints minus the mean of the first two. No inferential interval or significance test is assigned to these three runs. Same-seed trajectories need not agree across machines.

D Supporting trait-only experiment

D.1 Design and analysis

The trait model combines inherited speed, size, sensing range and reach with fixed behavioral rules. Populations spend 100 generations in environment A and 50 in B. Each final run seed defines its own searches and deployments. There are 200 seeds per primary group and arm (2000–2199). Shared food, behavior and initial-position streams couple groups with the same seed. The main B has 25 food items; three additional arms permit agent consumption in B, enlarge the B map to side 750, or reduce B's energy budget per generation to 25. The larger-map and reduced-energy settings were selected during recorded pilot and amendment rounds before the corresponding final runs.

The generation rules derive from the MinuteLabs Evolution Simulator at commit `55af956` (Palfree, 2021). A has side length 500 and 50 uniformly placed food items. Main B has side length 300 and 25 items in five clusters with standard deviation 15; centers and food are redrawn each generation. A generation lasts at most 200 ticks, with an energy allowance of 500 per agent except in the reduced-energy B arm. Agents start at edge home positions, search using fixed steering rules and return home according to feeding and remaining energy. Exhaustion and, in the consumption arm, predation can kill agents; agents with no food at generation end also die. Survivors carry into the next generation, and each survivor that consumed at least two food units produces exactly one offspring. Food units count both food items and prey. Effective movement speed is $u = \text{speed} \times \text{size}/10$, and each active movement tick costs $10^{-4}(\text{size}^3 u^2 + \text{sense})$. Food supports survival and reproduction without replenishing this generation-level energy allowance. There is no externally supported population floor.

Initial populations have 50 agents. ET starts from speed/size/sense/reach values 10/10/20/1. Mutating offspring receive independent Gaussian trait perturbations with standard deviations 1/1/1/0.25; sensing range is clipped at zero and the other traits at the smallest positive normalized float64 value. ET and M-pop mutate in both phases. F-food, F-pop, SV and Oracle-B have mutation disabled throughout. ET-freeze and ET-clone are derived from ET after A and have mutation disabled in B. SV samples each trait independently and log-uniformly over the search ranges: speed and size 2.5–40, sense 5–80 and raw reach 0.25–4.

In the consumption arm, a predator can consume only an agent whose size is at most 80% of its own. Identical clones therefore cannot consume one another. Comparisons involving clones and heterogeneous populations also differ in access to this additional food source; differences among heterogeneous groups depend on their size distributions. Across all 50 B generations, mean prey counts per run are 150.8 for ET, 153.2 for M-pop, 18.4 for ET-freeze and 3.4 for SV, with zero for the four homogeneous frozen groups.

The A search begins with eight logarithmically spaced values for each trait. Merging designs with the same effective reach, $\max(\text{reach}, \text{size}/4)$, leaves 1,472 distinct designs. Stage one evaluates 50 clones of each design for 30 generations on shared streams. Stage two evaluates the top 32 designs per objective on eight fresh sets of streams. Population search scores mean population over generations 10–99; food search scores food per agent over generations 10–29, even when a candidate was simulated for longer. The comparison therefore changes both objective and scoring horizon. Search-selected mutating agents inherit effective reach.

The group identifiers used in the tables are: ET, the evolving population; F-food and F-pop, frozen copies of the food- and population-search winners; M-pop, mutating copies of the population-search winner; SV, a frozen initial population with standing variation; ET-freeze, the end-of-A evolved population with mutation disabled in B; ET-clone, copies of a randomly sampled end-of-A

member with mutation disabled; and Oracle-B, the reference design selected directly in B. Oracle-B is not part of the confirmatory contrast family. Cloning changes the realized traits along with heterogeneity.

P1 is population at the start of a generation, averaged over B generations 40–49 with zeros retained. P2 indicates that no agent would start B generation 50. Each arm has eight planned comparison pairs, except arm 2 omits the population-search versus food-search pair because those data repeat the main arm by construction: 31 pairs in total. P1 uses a two-sided 10,000-permutation test of mean differences and a studentized bootstrap interval with 20,000 draws. P2 uses a two-sided Fisher test and a Newcombe interval for the difference in extinction proportions. Holm adjustment is within each arm's 16-test family, or 14 tests for arm 2. Shared streams make independent-group test calibration an approximation; paired sensitivity is retained separately in the supporting analysis records.

The practical margins are 2.5 agents for P1 and 0.10 for P2. The original "meaningfully different" label requires adjusted $p < 0.05$ and an absolute point estimate at least as large as the margin; it does not require the entire interval to exceed it. Equivalence uses an interval at level $1 - 2(0.05)/k$, where k is the arm's family size, wholly inside the margins. These equivalence intervals are wider than the 95% intervals printed below. Boundary labels distinguish ceiling, one-sided ceiling and floor cases. Equivalence near a population ceiling does not establish interchangeable adaptability.

D.2 Full primary outcome family and diagnostic controls

Table 10 reports all 31 pairs and both outcomes. Table 11 gives every primary group's mean population, extinction count, pre-shift losses and boundary frequencies. Full text labels, including ceiling qualifications and A-phase-loss flags, are also retained in `data/trait_complete_contrasts.csv`. Figure 7 provides the group-level display; its intervals do not replace the contrast rules.

Table 10: All 31 trait-study comparison pairs; all differences are X minus Y, with comparisons written X / Y. P1 is mean late-B population and P2 is terminal extinction probability. Entries are differences with 95% intervals. MD: significant with point estimate at or beyond the margin; EQ: equivalence under the specified interval rule; DU: detected difference with size unresolved; IN: inconclusive. Boundary qualifications are in Table 11; full labels are retained in the accompanying CSV.

Arm-row	X / Y	P1 difference [CI]	P2 difference [CI]	Adj. p_1 / p_2	P1/P2
main-1	F-food / ET	-20.93 $[-21.66, -20.11]$	$+0.71$ $[0.64, 0.77]$	0.0013 / 9.54×10^{-60}	MD/MD
main-2	F-pop / ET	$+0.64$ $[0.59, 0.68]$	$+0.00$ $[-0.02, 0.02]$	0.0013 / 1	EQ/EQ
main-3	M-pop / ET	-0.02 $[-0.08, 0.04]$	$+0.00$ $[-0.02, 0.02]$	1 / 1	EQ/EQ
main-4	SV / ET	-0.16 $[-0.64, 0.10]$	$+0.00$ $[-0.02, 0.02]$	1 / 1	EQ/EQ
main-5	ET-freeze / ET	-0.23 $[-0.41, -0.08]$	$+0.00$ $[-0.02, 0.02]$	0.04 / 1	EQ/EQ
main-6	ET-clone / ET-freeze	-3.11 $[-4.36, -2.11]$	$+0.10$ $[0.06, 0.15]$	0.0013 / 1.62×10^{-5}	MD/MD
main-7	F-pop / F-food	$+21.57$ $[20.76, 22.28]$	-0.71 $[-0.77, -0.64]$	0.0013 / 9.54×10^{-60}	MD/MD

Arm-row	X / Y	P1 difference [CI]	P2 difference [CI]	Adj. p_1 / p_2	P1/P2
main-8	M-pop / F-pop	-0.66 $[-0.70, -0.62]$	$+0.00$ $[-0.02, 0.02]$	0.0013 / 1	EQ/EQ
A2-1	F-food / ET	-19.16 $[-20.02, -18.17]$	$+0.70$ $[0.63, 0.76]$	0.0012 / 5.18×10^{-56}	MD/MD
A2-2	F-pop / ET	$+2.41$ $[1.97, 3.02]$	-0.01 $[-0.04, 0.01]$	0.0012 / 1	DU/EQ
A2-3	M-pop / ET	$+1.68$ $[1.05, 2.34]$	-0.01 $[-0.03, 0.02]$	0.0012 / 1	DU/EQ
A2-4	SV / ET	$+1.40$ $[0.75, 2.05]$	-0.01 $[-0.03, 0.02]$	0.0012 / 1	EQ/EQ
A2-5	ET-freeze / ET	$+1.24$ $[0.76, 1.84]$	-0.01 $[-0.04, 0.01]$	0.0012 / 1	EQ/EQ
A2-6	ET-clone / ET-freeze	-2.80 $[-4.03, -1.81]$	$+0.10$ $[0.06, 0.15]$	0.0012 / 1.50×10^{-5}	MD/MD
A2-8	M-pop / F-pop	-0.73 $[-1.26, -0.38]$	$+0.01$ $[-0.01, 0.03]$	0.0018 / 1	EQ/EQ
A3-1	F-food / ET	-18.10 $[-18.83, -17.18]$	$+0.94$ $[0.89, 0.96]$	0.0012 / 5.98×10^{-98}	MD/MD
A3-2	F-pop / ET	$+5.74$ $[5.00, 6.66]$	-0.07 $[-0.11, -0.03]$	0.0012 / 0.0012	MD/DU
A3-3	M-pop / ET	$+5.53$ $[4.80, 6.43]$	-0.07 $[-0.11, -0.03]$	0.0012 / 0.0012	MD/DU
A3-4	SV / ET	-6.86 $[-8.56, -5.13]$	$+0.39$ $[0.31, 0.46]$	0.0012 / 1.48×10^{-18}	MD/MD
A3-5	ET-freeze / ET	-0.50 $[-1.87, 0.81]$	$+0.07$ $[0.02, 0.14]$	0.9333 / 0.07	EQ/IN
A3-6	ET-clone / ET-freeze	-14.16 $[-15.54, -12.58]$	$+0.66$ $[0.58, 0.72]$	0.0012 / 1.44×10^{-41}	MD/MD
A3-7	F-pop / F-food	$+23.84$ $[23.70, 23.97]$	-1.00 $[-1.00, -0.97]$	0.0012 / 3.11×10^{-118}	MD/MD
A3-8	M-pop / F-pop	-0.22 $[-0.39, -0.04]$	$+0.00$ $[-0.02, 0.02]$	0.07 / 1	EQ/EQ
A4-1	F-food / ET	-13.43 $[-15.02, -11.81]$	$+0.57$ $[0.50, 0.64]$	0.0012 / 3.99×10^{-44}	MD/MD
A4-2	F-pop / ET	$+0.12$ $[-1.98, 2.27]$	-0.14 $[-0.23, -0.04]$	1 / 0.0329	IN/MD
A4-3	M-pop / ET	-0.76 $[-3.04, 1.49]$	$+0.04$ $[-0.06, 0.13]$	1 / 1	IN/IN
A4-4	SV / ET	-10.58 $[-12.43, -8.63]$	$+0.45$ $[0.36, 0.53]$	0.0012 / 1.01×10^{-20}	MD/MD
A4-5	ET-freeze / ET	-3.93 $[-6.12, -1.66]$	$+0.14$ $[0.05, 0.24]$	0.0032 / 0.0303	MD/MD
A4-6	ET-clone / ET-freeze	-8.97 $[-10.63, -7.33]$	$+0.41$ $[0.33, 0.48]$	0.0012 / 1.39×10^{-23}	MD/MD
A4-7	F-pop / F-food	$+13.55$ $[12.15, 14.93]$	-0.71 $[-0.77, -0.64]$	0.0012 / 9.54×10^{-60}	MD/MD
A4-8	M-pop / F-pop	-0.88 $[-3.02, 1.24]$	$+0.17$ $[0.08, 0.26]$	1 / 0.0044	IN/MD




Figure 7: Supporting trait experiment. All eight primary groups in four arms, with 200 seeds per group. Upper panels: mean P1 and descriptive whole-seed percentile 95% intervals (20,000 draws). Lower panels: P2 proportions with Wilson 95% intervals. These differ from the studentized P1 and Newcombe P2 intervals for contrasts. Shared seeds induce dependence across groups. Population and food searches differ in objective and scoring horizon; no-shift controls constrain diversity interpretations.

Table 11: Trait-study group outcomes and boundary diagnostics. All groups have 200 seeds. High/low are fractions with P1 at least 22.5 / at most 2.5. In arm 2, 25 is not a hard population ceiling. A-dead identifies losses preceding the environmental shift.

Arm	Group	Mean P1	Extinct	A-dead	High	Low
main	F-food	3.43	142/200	62	0.00	0.69
main	ET	24.36	0/200	0	1.00	0.00
main	F-pop	25.00	0/200	0	1.00	0.00
main	M-pop	24.34	0/200	0	1.00	0.00
main	SV	24.20	0/200	0	0.92	0.00
main	ET-freeze	24.13	0/200	0	0.90	0.00
main	ET-clone	21.02	20/200	0	0.73	0.10
main	Oracle-B	25.00	0/200	0	1.00	0.00
A2	F-food	3.43	142/200	62	0.00	0.69
A2	ET	22.59	2/200	0	0.61	0.01
A2	F-pop	25.00	0/200	0	1.00	0.00
A2	M-pop	24.27	1/200	0	0.85	0.01
A2	SV	23.99	1/200	0	0.90	0.01
A2	ET-freeze	23.82	0/200	0	0.86	0.00
A2	ET-clone	21.02	20/200	0	0.73	0.10
A2	Oracle-B	25.00	0/200	0	1.00	0.00
A3	F-food	0.00	200/200	62	0.00	1.00

Arm	Group	Mean P1	Extinct	A-dead	High	Low
A3	ET	18.10	13/200	0	0.10	0.07
A3	F-pop	23.84	0/200	0	0.93	0.00
A3	M-pop	23.63	0/200	0	0.94	0.00
A3	SV	11.24	90/200	0	0.24	0.45
A3	ET-freeze	17.60	28/200	0	0.33	0.14
A3	ET-clone	3.44	160/200	0	0.04	0.79
A3	Oracle-B	24.76	0/200	0	1.00	0.00
A4	F-food	0.00	200/200	62	0.00	1.00
A4	ET	13.43	85/200	0	0.50	0.42
A4	F-pop	13.55	58/200	0	0.34	0.28
A4	M-pop	12.67	92/200	0	0.54	0.46
A4	SV	2.85	175/200	0	0.09	0.88
A4	ET-freeze	9.50	114/200	0	0.28	0.56
A4	ET-clone	0.54	195/200	0	0.01	0.97
A4	Oracle-B	25.00	0/200	0	1.00	0.00

In the main arm, the food-search group's deficit includes 62 of 200 runs lost during A. Restricting analysis to groups that both reach B selects survivors and can bias the comparison; such conditional outcomes are retained as diagnostics rather than replacing unconditional results. The no-move control continues ET's end-of-A population for fifty further generations in A. Mean late population is 48.0675 with mutation, 47.9895 after freezing, and 40.9730 after cloning. The freeze-minus-mutation difference is -0.0780, interval $[-0.2720, 0.1140]$, which includes zero. The clone-minus-freeze difference is -7.0165, interval $[-9.1271, -5.0470]$. The clone deficit therefore also occurs without an environmental shift. No formal B-minus-A interaction test follows from comparing whether separate intervals include zero.

The predefined diagnostic that B might not require a different design is triggered in the main, consumption and larger-map arms, but not the reduced-energy arm. The positive control that B separates at least some groups holds in all four. The majority-single-design warning applies to the standing-variation group in all four arms. The search-reliability diagnostic U3 is not triggered. The U1 boundary frequencies and U4 pre-shift loss counts are shown in Table 11; they restrict how the apparent ties should be interpreted. No group exceeds the U4 threshold of losing more than half its runs during A.

Exploratory analyses use subsets of the retained seeds and do not change primary decisions. They cover alternative map sizes and food/tick settings, landscape ranks, half/double mutation sizes, longer B deployment, stage-one-only search winners, populations started directly in B, trait-change summaries, first-generation deaths, energy use and no-move controls. Their complete numerical outputs and sample sizes are preserved in `data/trait_exploratory.json`. For example, on the 100-seed exploratory subset, starting ET directly in B instead of transferring from A changes P1 by -0.22 in the main arm, -17.62 in the larger-map arm and -12.85 in the reduced-energy arm. These initialization comparisons concern the trait model.

E Chronology and reproducibility

Pilot and tuning stages preceded the corresponding final runs. The private protocols specified five neural procedures and seven secondary comparisons. The pilot-based precision estimate used three seeds per method. Ten final runs were chosen per procedure, retaining the 0.05 margin with limited precision for equivalence. ES-fixed was specified after inspection of the core results and before its own tuning and ten final runs. The one-billion-step OE extension was exploratory. Planned lifetime-learning and neural mechanism ablations were not run.

The neural core training and evaluation records identify revision `16e8657`; the ES-fixed extension identifies `06ff7b2`. Trait generation records identify `a1e1cdc` and the confirmatory analysis identifies `561bcd4`. The source inspection used revision `ea7792af7f66`. Full identifiers and configuration hashes are provided with the code.

Core neural records report NVIDIA RTX 4090 hardware, JAX 0.11.2 and Python 3.13.18. The statistical reanalysis used CPU execution with Python 3.12.14 and separately recorded dependencies. Two same-seed ES-DR check runs under the addendum configuration remained within the specified tolerance around the core range. Cross-machine runs need not be numerically identical. The test-access logs cover the recorded evaluations; access outside those logs cannot be assessed.

The archive contains all sixty final neural evaluations and 19,800 adaptation checkpoint observations, trait generation records, final trained policies, training snapshots and tuning-trial records. Core tuning policy files were not retained. A correction to detection of completely tied Mann–Whitney samples changed no reported effect, p value or decision.

The repository includes source code, configurations, data, analysis scripts and the manuscript build files. Its README gives the commands for reproducing the tables, figures and statistical checks.