Table of Contents

S.no	Title	Population Set
1	Table 02: Baseline Demographic and Clinical Characteristics [FDA STF-IG]	Safety Population 2
2	Table 3.1.1: Study Disposition [JPMA]	Safety Population 2
3	Table 3.1.2: Summary of Reason for Discontinuation [JPMA]	Safety Population 2
4	Table 3.2: Summary of Protocol Deviations [JPMA]	Safety Population 2
5	Table 3.3: Summary of Data Sets Analyzed [JPMA]	Safety Population 2
6	Table 3.5.1.2: Subgroup Analyses of <efficacy continuous variables> [JPMA]	Efficacy Analysis Set
7	Table 3.7: Time to <efficacy event(s)> Analyses [JPMA]	Efficacy Analysis Set
8	Table 3.8: Response Rate of <efficacy categorical variables> [JPMA]	Safety Population 2
9	Table 3.9: Compliance with Investigational Product [JPMA]	Safety Population 2
10	Table 3.10: Summary of Subject Incidence of Treatment-Emergent Adverse Events [JPMA]	Safety Population 2
11	Table 3.11.1: Treatment-Emergent Adverse Events [JPMA]	Safety Population 2
12	Table 3.12: Summary of Laboratory Tests by Visit [JPMA]	Safety Population 2
13	Table 3.13.1: Shells in Site Reference Value from Baseline for Laboratory Tests [JPMA]	Safety Population 2
14	Table 3.13.2: Summary of Abnormal Change for Laboratory Tests [JPMA]	Safety Population 2
15	Table 3.14.1: Summary of Vital Signs by Visit [JPMA]	Safety Population 2
16	Table 3.14.2: Summary of ECG Findings [JPMA]	Safety Population 2
17	Table 06: Duration of Treatment Exposure [FDA STF-IG]	Safety Population 2
18	Table 07: Overview of Adverse Events [FDA STF-IG]	Safety Population 2
19	Table 08: Deaths [FDA STF-IG]	Safety Population 2
20	Table 12: Subjects With Adverse Events Leading to Treatment Discontinuation by System Organ Class and Preferred Term [FDA	Safety Population 2

	STF-IG]	
21	Table 13: Subjects With Adverse Events Leading to Treatment Discontinuation by Organ System and OCMQ (Narrow) [FDA STF-IG]	Safety Population 2
22	Table 15: Subjects With Common Adverse Events Occurring at ≥X% Frequency by Preferred Term	Safety Population 2
23	Table 17: Subjects With Adverse Events by Organ System and OCMQ [FDA STF-IG]	Safety Population 2
24	Table 20: Subjects With Kidney Function Analyte Values Exceeding Specified Levels [FDA STF-IG]	Safety Population 2
25	Table 44: Subjects With Adverse Events by Male-Specific OCMQ (Broad) and Preferred Term [FDA STF-IG]	Male Safety Population

Table 02

Baseline Demographic and Clinical Characteristics [FDA STF-IG]

Safety Population 2

Pooled Analyses (or Trial X)

Characteristic	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Total Population N = XXX**
Sex, n (%)	X (Y)	X (Y)	X (Y)
Male	X (Y)	X (Y)	X (Y)
Female	X (Y)	X (Y)	X (Y)
Age, Years			
Mean (SD)	X (Y.YY)	X (Y.YY)	X (Y.YY)
Median (min, max)	X (Y, Z)	X (Y, Z)	X (Y, Z)
Age groups (years), n (%)	X (Y)	X (Y)	X (Y)
Group X	X (Y)	X (Y)	X (Y)
Race, n (%)	X (Y)	X (Y)	X (Y)
American Indian or Alaska Native	X (Y)	X (Y)	X (Y)
Asian	X (Y)	X (Y)	X (Y)
Black or African American	X (Y)	X (Y)	X (Y)
Native Hawaiian or Other Pacific Islander	X (Y)	X (Y)	X (Y)
White	X (Y)	X (Y)	X (Y)
Not reported	X (Y)	X (Y)	X (Y)
Other	X (Y)	X (Y)	X (Y)
Unknown	X (Y)	X (Y)	X (Y)

Ethnicity, n (%)	X (Y)	X (Y)	X (Y)
Hispanic	X (Y)	X (Y)	X (Y)
Not Hispanic or Latino	X (Y)	X (Y)	X (Y)
Unknown	X (Y)	X (Y)	X (Y)
Country of participation, n (%)	X (Y)	X (Y)	X (Y)
Country X	X (Y)	X (Y)	X (Y)
Clinical baseline characteristics, n (%)	X (Y)	X (Y)	X (Y)
Characteristic X	X (Y)	X (Y)	X (Y)

Source: [include Applicant source, datasets and/or software tools used].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

Abbreviations: n, number of subjects with given characteristic; N, number of subjects in treatment arm; SD, standard deviation.

Table 3.1.1

Study Disposition [JPMA]

Safety Population 2

	Treatment A **N=XX** **n (%)**	**Treatment B** **N=XX** **n (%)**
Randomized	XX (XX.X)	XX (XX.X)
Randomized, but not treated	XX (XX.X)	XX (XX.X)
Enrolled	XX (XX.X)	XX (XX.X)
Enrolled, but not treated	XX (XX.X)	XX (XX.X)
Treated investigational product	XX (XX.X)	XX (XX.X)
Completed study	XX (XX.X)	XX (XX.X)
Completed treatment	XX (XX.X)	XX (XX.X)
Discontinued study	XX (XX.X)	XX (XX.X)
Discontinued treatment	XX (XX.X)	XX (XX.X)

Table 3.1.2

Summary of Reason for Discontinuation [JPMA]

Safety Population 2

	Treatment A **N=XX** **n (%)**	**Treatment B** **N=XX** **n (%)**
Randomized	XX (XX.X)	XX (XX.X)
Enrolled	XX (XX.X)	XX (XX.X)
Completed study	XX (XX.X)	XX (XX.X)
Discontinued study	XX (XX.X)	XX (XX.X)
Reason for Discontinuation from Study X	XX (XX.X)	XX (XX.X)
Completed investigational product	XX (XX.X)	XX (XX.X)
Discontinued investigational product	XX (XX.X)	XX (XX.X)
Reason for Discontinuation of Treatment X	XX (XX.X)	XX (XX.X)

Table 3.2

Summary of Protocol Deviations [JPMA]

Safety Population 2

	n (%)
All Center	
Treatment A (N=XX)	
Number of protocol deviations	XX (XX.X)
Protocol deviation X	XX (XX.X)
Treatment B (N=XX)	
Number of protocol deviations	XX (XX.X)
Protocol deviation X	XX (XX.X)
Center X	
Treatment A (N=XX)	
Number of protocol deviations	XX (XX.X)
Protocol deviation X	XX (XX.X)
Treatment B (N=XX)	
Number of protocol deviations	XX (XX.X)
Protocol deviation X	XX (XX.X)

Table 3.3

Summary of Data Sets Analyzed [JPMA]

Safety Population 2

	Treatment A **N=XX** **n (%)**	**Treatment B** **N=XX** **n (%)**
ITT	XX (XX.X)	XX (XX.X)
Safety	XX (XX.X)	XX (XX.X)
FAS	XX (XX.X)	XX (XX.X)
PPS	XX (XX.X)	XX (XX.X)

Table 3.5.1.2

Subgroup Analyses of <efficacy continuous variables> [JPMA]

Efficacy Analysis Set

<Category> (N=XX)

	Treatment A N=XX	**Treatment B N=XX**
<Parameter>	XX (XX.X)	XX (XX.X)
<Visit>	XX (XX.X)	XX (XX.X)
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX

Table 3.7

Time to <efficacy event(s)> Analyses [JPMA]

Efficacy Analysis Set

	Treatment A N=XX	**Treatment B N=XX**
Subjects with event	XX (XX.X %)	XX (XX.X %)
Subjects without event	XX (XX.X %)	XX (XX.X %)
Time to event (days)		
Median (95% CI)	XX.X (XXX.X, XXX.X)	XX.X (XXX.X, XXX.X)
Hazard Ratio (95% CI)	X.XXX (XXX.X, XXX.X)	X.XXX (XXX.X, XXX.X)
p-value	XXX.X	XXX.X

EXPORT-STUDY-01

Table 3.8

Response Rate of <efficacy categorical variables> [JPMA]

Safety Population 2

	Treatment A N=XX	**Treatment B N=XX**
Responders	XX (XX.X %)	XX (XX.X %)
Non-Responders	XX (XX.X %)	XX (XX.X %)
95% CI for Response Rates	XX.X, XXX.X	XX.X, XXX.X
Difference in Response Rates	XX.X	XX.X
95% CI		XX.X, XX.X
Odds Ratio	XXX.X	XXX.X
95% CI		X
<Category> (%)	XX (XX.X %)	XX (XX.X %)
95% CI	XX.X, XX.X	XX.X, XX.X

Table 3.9

Compliance with Investigational Product [JPMA]

Safety Population 2

	Treatment A N=XX	**Treatment B N=XX**
Duration of treatment (day)		
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
Compliance Rate (%)		
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
Patients with < X %		
n (%)	XX (XX.X %)	XX (XX.X %)

Table 3.10

Summary of Subject Incidence of Treatment-Emergent Adverse Events [JPMA]

Safety Population 2

	Treatment A **N=XX** **n (%)**	**Treatment B** **N=XX** **n (%)**
TEAE	XX (XX.X)	XX (XX.X)
AE related to study treatment	XX (XX.X)	XX (XX.X)
Serious TEAE	XX (XX.X)	XX (XX.X)
Serious TEAE related to study treatment	XX (XX.X)	XX (XX.X)
TEAE leading to death	XX (XX.X)	XX (XX.X)
AE related to study treatment leading to death	XX (XX.X)	XX (XX.X)
Other important TEAE	XX (XX.X)	XX (XX.X)
Other important AE related to study treatment	XX (XX.X)	XX (XX.X)

TEAE: Treatment emergent adverse events

Table 3.11.1

Treatment-Emergent Adverse Events [JPMA]

Safety Population 2

	Treatment A **N=XX** **n (%)**	**Treatment B** **N=XX** **n (%)**
Number of subjects reporting at least one TEAE	XX (XX.X)	XX (XX.X)
System Organ Class X	XX (XX.X)	XX (XX.X)
Preferred Term X	XX (XX.X)	XX (XX.X)

TEAE: Treatment emergent adverse events

MedDRA ver. xx.x

Table 3.12

Summary of Laboratory Tests by Visit [JPMA]

Safety Population 2

Characteristics	**Treatment A N=XX**	**Treatment B N=XX**
<Parameter>		
<Visit>		
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX

Table 3.13.1

Shells in Site Reference Value from Baseline for Laboratory Tests [JPMA]

Safety Population 2

	Baseline			
	Normal **n (%)**	**Low** **n (%)**	**High** **n (%)**	**Total** **n (%)**
<Parameter>				
<Visit>	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Normal	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Low	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
High	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Total	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)

Table 3.13.2

Summary of Abnormal Change for Laboratory Tests [JPMA]

Safety Population 2

	Treatment A N=XX	**Treatment B N=XX**
<Parameter>	XX (XX.X)	XX (XX.X)
<Visit>	XX (XX.X)	XX (XX.X)
n	XX	XX
High n (%)	XX (XX.X)	XX (XX.X)
Low n (%)	XX (XX.X)	XX (XX.X)

Table 3.14.1

Summary of Vital Signs by Visit [JPMA]

Safety Population 2

Characteristics	**Treatment A N=XX**	**Treatment B N=XX**
<Parameter>		
<Visit>	XX (XX.XX)	XX (XX.XX)
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX

Table 3.14.2

Summary of ECG Findings [JPMA]

Safety Population 2

Characteristics	**Treatment A N=XX**	**Treatment B N=XX**
<Visit>		
n	XX	XX
Normal (%)	XX (XX.X)	XX (XX.X)
Abnormal, not clinically significant (%)	XX (XX.X)	XX (XX.X)
Abnormal, clinically significant (%)	XX (XX.X)	XX (XX.X)
Unable to evaluate (%)	XX (XX.X)	XX (XX.X)

EXPORT-STUDY-01

Table 06

Duration of Treatment Exposure [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

Parameter	**Drug Name Dosage A N = XXX**	**Control N = XXX**
Duration of treatment, weeks (or months, or days, or cycles)		
Mean (SD)	X (Y)	X (Y)
Median (Min, Max)	X (Y, Z)	X (Y, Z)
Interquartile range	X - Y	X - Y
Total exposure (person years)	X (Y)	X (Y)
Patients treated, by duration, n (%)		
≥ 1 day	X (Y)	X (Y)
≥1 month	X (Y)	X (Y)
≥3 month	X (Y)	X (Y)
≥6 month	X (Y)	X (Y)
≥12 month	X (Y)	X (Y)

Abbreviations: n, number of subjects with given treatment duration; N, number of subjects in treatment arm; SD, standard deviation.

Source: [include Applicant source, datasets and/or software tools used].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

Table 07

Overview of Adverse Events [FDA STF-IG]

Safety Population 2

Pooled Analyses (or Trial X)

Adverse Event Category	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI)** [1]
SAE [2]			X (Y, Z)
Deaths	X (Y)	X (Y)	X (Y, Z)
Life-threatening	X (Y)	X (Y)	X (Y, Z)
Initial or prolonged hospitalization	X (Y)	X (Y)	X (Y, Z)
Disability or permanent damage	X (Y)	X (Y)	X (Y, Z)
Congenital anomaly or birth defect	X (Y)	X (Y)	X (Y, Z)
Other	X (Y)	X (Y)	X (Y, Z)
AE leading to permanent discontinuation of treatment	X (Y)	X (Y)	X (Y, Z)
AE leading to action taken of treatment [3]	X (Y)	X (Y)	X (Y, Z)
Action X	X (Y)	X (Y)	X (Y, Z)
Any AE [4]	X (Y)	X (Y)	X (Y, Z)
Severity X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with at least one event; N, number of subjects in treatment arm; SAE, serious adverse event.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

[2] SAEs classified by Applicant as [insert Applicant’s definition of SAE].

[3] Subjects may be counted more than once.

[4] Severity scale as defined by the protocol.

Table 08

Deaths [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

Deaths	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [1]**
Death Details	X (Y)	X (Y)	X (Y, Z)
Cause of death X	X (Y)	X (Y)	X (Y, Z)
Adverse events[2] with an outcome of death	X (Y)	X (Y)	X (Y, Z)
Cause of death X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with adverse event; N, number of subjects in treatment arm; PT, preferred term.

Source: [include Applicant source, datasets and/or software tools used].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

1 Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

2 Treatment-emergent AE defined as [definition]. MedDRA version X.

Table 12

Subjects With Adverse Events Leading to Treatment Discontinuation by System Organ Class and Preferred Term [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

System Organ Class Preferred Term	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [1]**
Patients with at least one AE leading to discontinuation	X (Y)	X (Y)	X (Y, Z)
SOC X	X (Y)	X (Y)	X (Y, Z)
PT X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: CI, confidence interval; MedDRA, Medical Dictionary for Regulatory Activities; N, number of patients in treatment arm; n, number of patients with at least one event; PT, preferred term; SOC, System Organ Class

Source: [include Applicant source, datasets and/or software tools used].
[1] Treatment-emergent adverse event defined as [definition]. MedDRA version X.
[2] Duration = [e.g., X week double-blind treatment period or median and a range indicating pooled trial durations].
[3] Difference is shown between [treatment arms] (e.g., difference is shown between Drug Name dosage X vs. placebo).

Table 13

Subjects With Adverse Events Leading to Treatment Discontinuation by Organ System and OCMQ (Narrow) [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

Organ System[1] OCMQ (Narrow)	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [2]**
Organ System X			X (Y, Z)
OCMQ X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with adverse event; N, number of subjects in treatment arm; PT, preferred term; SOC, system organ class.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

1 Each OCMQ is aligned to a single organ system based on clinical judgment. Please be aware that some OCMQs may contain PTs from more than one MedDRA SOC.

2 Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 15

Subjects With Common Adverse Events Occurring at ≥X% Frequency by Preferred Term

Safety Population 2

Pooled Analysis (or Trial X)

Preferred Term[1]	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [2], [3]**
PT X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with adverse event; N, number of subjects in treatment arm; PT, preferred term.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Coded as MedDRA preferred terms.

[2] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

[3] Table display is ordered by the risk difference.

Table 17

Subjects With Adverse Events by Organ System and OCMQ [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

Organ System [2] OCMQ	**Narrow OCMQs**			**Broad OCMQs[1]**		
	Drug Name Dosage A N = XXX n (%)	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [3]**	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference**
Organ System X			X (Y, Z)			X (Y, Z)
OCMQ X	X (Y)	X (Y)	X (Y, Z)	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with at least one event; N, number of subjects in treatment arm. PT, preferred term; SOC, system organ class.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Broad OCMQ analysis incorporates narrow OCMQ preferred terms to maximize sensitivity.

[2] Each OCMQ is aligned to a single organ system based on clinical judgment. Please be aware that some OCMQs may contain PTs from more than one MedDRA SOC.

[3] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 20

Subjects With Kidney Function Analyte Values Exceeding Specified Levels [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

	Drug Name Dosage A N = XXX n (%)	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [1]**
Creatinine, high (mg/dL)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 1 (≥1.5 x baseline)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 2 (≥2.0 x baseline)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 3 (≥3.0 x baseline)	X/Y (Z)	X/Y (Z)	X (Y, Z)
eGFR, low (mL/min/1.73 m2)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 1 (≥25% decrease)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 2 (≥50% decrease)	X/Y (Z)	X/Y (Z)	X (Y, Z)
Level 3 (≥75% decrease)	X/Y (Z)	X/Y (Z)	X (Y, Z)

Abbreviations: CI, confidence interval; eGFR, estimated glomerular filtration rate; n, number of subjects meeting the specified laboratory criteria; N, number of subjects in treatment arm; Ns, total number of subjects with data available for the laboratory test of interest.

Source: [include Applicant source, datasets and/or software tools used].

Note: Threshold Levels 1, 2, and 3 are defined by Table 56. Abnormality Level Criteria for Chemistry Laboratory Results.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Note: Subject counts are cumulative for each abnormality threshold.

Table 44

Subjects With Adverse Events by Male-Specific OCMQ (Broad) and Preferred Term [FDA STF-IG]

Male Safety Population

Pooled Analysis (or Trial X)

OCMQ (Broad) Preferred Term [1]	**Drug Name Dosage A N = XXX n/N_s (%)**	**Control N = XXX n/N_s (%)**	**Risk Difference % (95% CI) [2,3]**
Erectile Dysfunction	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Gynecomastia	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom Medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of male subjects with indicated OCMQ; N, number of subjects in treatment arm; Ns, number of male subjects with adverse event data.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Broad OCMQ analysis incorporates narrow OCMQ preferred terms to maximize sensitivity.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] OCMQs include AEs that are not MedDRA PTs.

[2] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

[3] Table displays categories by risk difference.