Table of Contents

S.no	Title	Population Set
1	Table 3.4: Demographic and other baseline characteristics [JPMA]	Safety Population 2
2	Table 3.5.1.1: Summary of <efficacy continuous variables> [JPMA]	Efficacy Analysis Set
3	Table 3.5.2.1: Summary of <efficacy categorical variables> [JPMA]	Efficacy Analysis Set
4	Table 3.5.2.2: Subgroup Analyses of <efficacy categorical variables> [JPMA]	Efficacy Analysis Set
5	Table 3.6: Covariate Analyses of <efficacy continuous variables> [JPMA]	Efficacy Analysis Set
6	Table 3.11.2: Treatment-Emergent Adverse Events by Severity [JPMA]	Safety Population 2
7	Table 04: Subject Populations [FDA STF-IG]	Randomized Population
8	Table 10: Patients With Serious Adverse Events by System Organ Class and Preferred Term [FDA STF-IG]	Safety Population 2
9	Table 11: Subjects With Serious Adverse Events by Organ System and OCMQ (Narrow) [FDA STF-IG]	Safety Population 2
10	Table 18: Summary Assessment of [Insert AE of Interest] [FDA STF-IG]	Safety Population 2
11	Table 33: Subjects With Adverse Events by Male-Specific OCMQ (Narrow) and Preferred Term [FDA STF-IG]	Male Safety Population
12	Table 34: Subjects With Adverse Events by Female-Specific OCMQ (Narrow) and Preferred Term [FDA STF-IG]	Female Safety Population
13	Table 45: Subjects With Adverse Events by Female-Specific OCMQ (Broad) and Preferred Term [FDA STF-IG]	Female Safety Population
14	Figure 14.1.1: Distribution of Race	Safety Population 4
15	Figure 14.x.y: Bar Chart	Safety Population 3
16	Figure 14.x.y: Box Plot	Safety Population 3

17	Figure 14.1.2: Box Plot of Age Distribution	Safety Population 4
18	Figure 14.1.3: The Effect of Age on Weight by Gender	Safety Population 4
19	Figure 14.x.y: Line Chart	Safety Population 3
20	Figure 14.1.4: Distribution of Patients by Age Group and Ethnicity	Safety Population 4
21	Figure 14.x.y: Stacked Bar Chart	Safety Population 3
22	Figure 14.2.1.1 [ONC]: Kaplan-Meier Chart for Overall Survival	Efficacy Population
23	Figure 14.x.y: Kaplan-Meier Chart	Safety Population 3
24	Figure 14.2.1.2 [ONC]: Kaplan-Meier Chart for Progression Free Survival	Efficacy Population
25	Figure 14.2.2.1 [ONC]: Swimmer Chart - Patient Disposition and RECIST 1.1 Response	Efficacy Population

Table 3.4

Demographic and other baseline characteristics [JPMA]

Safety Population 2

	Treatment A N=XX	**Treatment B N=XX**
Sex		
n	XX	XX
<Sex>	XX (XX.X %)	XX (XX.X %)
Age (year)		
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
Age (year)		
n	XX	XX
<Age category>	XX (XX.X %)	XX (XX.X %)
Weight (kg)		
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
Weight (kg)		
n	XX	XX
<Weight category>	XX (XX.X %)	XX (XX.X %)
Height (cm)		
n	XX	XX

Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX
Race		
n	XX	XX
<Race>	XX (XX.X %)	XX (XX.X %)

Table 3.5.1.1

Summary of <efficacy continuous variables> [JPMA]

Efficacy Analysis Set

	Treatment A N=XX	**Treatment B N=XX**
<Parameter>	XX (XX.X)	XX (XX.X)
<Visit>	XX (XX.X)	XX (XX.X)
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min - Max	XX - XX	XX - XX

Table 3.5.2.1

Summary of <efficacy categorical variables> [JPMA]

Efficacy Analysis Set

	Treatment A **N=XX**	**Treatment B** **N=XX**
< Parameter >		
< Visit >		
< Category >	XX (XX.X %)	XX (XX.X %)

Table 3.5.2.2

Subgroup Analyses of <efficacy categorical variables> [JPMA]

Efficacy Analysis Set

<Category> (N=XX)

	Treatment A **N=XX**	**Treatment B** **N=XX**
< Parameter >		
< Visit >		
< Category >	XX (XX.X %)	XX (XX.X %)

Table 3.6

Covariate Analyses of <efficacy continuous variables> [JPMA]

Efficacy Analysis Set

Characteristics	**Treatment A N=XX**	**Treatment B N=XX**
Parameter		
n	XX	XX
LS Mean	XXX.X	XXX.X
Difference vs. Treatment A		XXX.X
95% CI for difference		XXX.X, XXX.X
p-value		X.XXX

Table 3.11.2

Treatment-Emergent Adverse Events by Severity [JPMA]

Safety Population 2

	Treatment A (N=XX)			
	Any **n (%)**	**Mild** **n (%)**	**Moderate** **n (%)**	**Severe** **n (%)**
Number of subjects reporting at least one TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
System Organ Class X	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term X	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)

TEAE: Treatment emergent adverse events

Table 04

Subject Populations [FDA STF-IG]

Randomized Population

Pooled Analysis (or Trial X)

Population	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**
Subjects randomized	X (Y)	X (Y)
ITT population	X (Y)	X (Y)
mITT population	X (Y)	X (Y)
Safety population	X (Y)	X (Y)
Per-protocol population	X (Y)	X (Y)

Abbreviations: ITT, intent-to-treat; mITT, modified intent-to-treat; n, number of subjects in specified subset of the treatment arm; N, number of subjects in treatment arm.

Source: [include Applicant source, datasets and/or software tools used].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

Table 10

Patients With Serious Adverse Events by System Organ Class and Preferred Term [FDA STF-IG]

Safety Population 2

Pooled Analyses (or Trial X)

System Organ Class Preferred Term	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [1]**
Any SAE	X (Y)	X (Y)	X (Y, Z)
SOC X	X (Y)	X (Y)	X (Y, Z)
PT X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: CI, confidence interval; n, number of subjects with adverse event; N, number of subjects in treatment arm; PT, preferred term; SAE, serious adverse event; SOC, system organ class.

Source: [include Applicant source, datasets and/or software tools used].

Note: SAEs classified by Applicant as [insert Applicant’s definition of SAE].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 11

Subjects With Serious Adverse Events by Organ System and OCMQ (Narrow) [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

Organ System[1] OCMQ (Narrow)	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI) [2]**
Organ System X			X (Y, Z)
OCMQ X	X (Y)	X (Y)	X (Y, Z)

Abbreviations: CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of subjects with adverse event; N, number of subjects in treatment arm; PT, preferred term; SAE, serious adverse event; SOC, system organ class.

Source: [include Applicant source, datasets and/or software tools used].

Note: SAEs classified by Applicant as [insert Applicant’s definition of SAE].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

1 Each OCMQ is aligned to a single organ system based on clinical judgment. Please be aware that some OCMQs may contain PTs from more than one MedDRA SOC.

2 Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 18

Summary Assessment of [Insert AE of Interest] [FDA STF-IG]

Safety Population 2

Pooled Analysis (or Trial X)

AE of Interest Assessment	**Drug Name Dosage A N = XXX n (%)**	**Control N = XXX n (%)**	**Risk Difference % (95% CI)** [1]
[Insert AE of Interest] [2]	X (Y)	X (Y)	X (Y, Z)
PT X	X (Y)	X (Y)	X (Y, Z)
Maximum severity [3]	X (Y)	X (Y)	X (Y, Z)
Severity X	X (Y)	X (Y)	X (Y, Z)
SAE [4]	X (Y)	X (Y)	X (Y, Z)
Death	X (Y)	X (Y)	X (Y, Z)
Life-threatening	X (Y)	X (Y)	X (Y, Z)
Initial or prolonged hospitalization	X (Y)	X (Y)	X (Y, Z)
Disability or permanent damage	X (Y)	X (Y)	X (Y, Z)
Congenital anomaly or birth defect	X (Y)	X (Y)	X (Y, Z)
Other	X (Y)	X (Y)	X (Y, Z)
Resulting in treatment discontinuation	X (Y)	X (Y)	X (Y, Z)

Abbreviations: AE, adverse event; AESI, adverse event of special interest; CI, confidence interval; OCMQ,

OND custom medical query; n, number of subjects with at least one event; N, number of subjects in treatment arm; PT, preferred term; SAE, serious adverse event.

Source: [include Applicant source, datasets and/or software tools used].

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

[2] Use OCMQ grouping if appropriate.

[3] Severity scale as defined by the protocol.

[4] SAEs classified by Applicant as [insert Applicant’s definition of SAE].

Table 33

Subjects With Adverse Events by Male-Specific OCMQ (Narrow) and Preferred Term [FDA STF-IG]

Male Safety Population

Pooled Analysis (or Trial X)

OCMQ (Narrow) Preferred Term [1]	**Drug Name Dosage A N = XXX n/N_s (%)**	**Control N = XXX n/N_s (%)**	**Risk Difference % (95% CI) [2]**
Erectile Dysfunction	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Gynecomastia	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of male subjects with indicated OCMQ; N, number of subjects in treatment arm; Ns, number of male subjects with adverse event data.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] OCMQs include AEs that are not MedDRA PTs.

[2] Risk Difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 34

Subjects With Adverse Events by Female-Specific OCMQ (Narrow) and Preferred Term [FDA STF-IG]

Female Safety Population

Pooled Analyses (or Trial X)

OCMQ (Narrow) Preferred Term [1]	**Drug Name Dosage A N = XXX n/N_s (%)**	**Control N = XXX n/N_s (%)**	**Risk Difference % (95% CI) [2]**
Abnormal uterine bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Amenorrhea	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Bacterial vaginosis	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Decreased menstrual bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Excessive menstrual bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of female subjects with indicated OCMQ; N, number of subjects in treatment arm; Ns, number of female subjects with adverse event data.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial durations].

[1] OCMQs include AEs that are not MedDRA PTs.

[2] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

Table 45

Subjects With Adverse Events by Female-Specific OCMQ (Broad) and Preferred Term [FDA STF-IG]

Female Safety Population

Pooled Analyses (or Trial X)

OCMQ (Narrow) Preferred Term [1]	**Drug Name Dosage A N = XXX n/N_s (%)**	**Control N = XXX n/N_s (%)**	**Risk Difference % (95% CI) [2,3]**
Abnormal uterine bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Amenorrhea	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Bacterial vaginosis	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Decreased menstrual bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)
Excessive menstrual bleeding	X/Y (Z)	X/Y (Z)	X (Y, Z)
PT X	X/Y (Z)	X/Y (Z)	X (Y, Z)

Abbreviations: AE, adverse event; CI, confidence interval; OCMQ, OND custom medical query; MedDRA, Medical Dictionary for Regulatory Activities; n, number of female subjects with indicated OCMQ; N, number of subjects in treatment arm; Ns, number of female subjects with adverse event data.

Source: [include Applicant source, datasets and/or software tools used].

Note: Treatment-emergent AE defined as [definition]. MedDRA version X.

Note: Broad OCMQ analysis incorporates narrow OCMQ preferred terms to maximize sensitivity.

Note: Duration = [e.g., X-week double-blind treatment period or median and a range of pooled trial

durations].

[1] OCMQs include AEs that are not MedDRA PTs.

[2] Risk difference is shown between [treatment arms] (e.g., difference is shown between Drug Name Dosage A vs. Control).

[3] Table displays categories by risk difference.

Figure 14.1.1

Distribution of Race

Safety Population 4



Source: adxx; Program name; Program run date/time

Figure 14.x.y

Bar Chart

Safety Population 3



Figure 14.x.y

Box Plot

Safety Population 3



Figure 14.1.2

Box Plot of Age Distribution

Safety Population 4



Source: adxx; Program name; Program run date/time

Figure 14.1.3

The Effect of Age on Weight by Gender

Safety Population 4



Source: adxx; Program name; Program run date/time

Figure 14.x.y

Line Chart

Safety Population 3



Figure 14.1.4

Distribution of Patients by Age Group and Ethnicity

Safety Population 4



Source: adxx; Program name; Program run date/time

EXPORT-STUDY-01

Figure 14.x.y

Stacked Bar Chart

Safety Population 3



Figure 14.2.1.1 [ONC]

Kaplan-Meier Chart for Overall Survival

Efficacy Population



Figure 14.x.y

Kaplan-Meier Chart

Safety Population 3



Figure 14.2.1.2 [ONC]

Kaplan-Meier Chart for Progression Free Survival

Efficacy Population



Figure 14.2.2.1 [ONC]

Swimmer Chart - Patient Disposition and RECIST 1.1 Response

Efficacy Population