Table of Contents

S.no	Title	Population Set
1	Table 14.1.1.1: Summary of Analysis Populations	All Enrolled Population
2	Table 14.1.1.3: Summary of Subject Disposition	Safety Population 2
3	Table 14.1.1.4.1: Summary of Eligibility Criteria Deviations	All Enrolled Population
4	Table 14.1.1.4.2: Summary of Key Protocol Deviations	Safety Population 2
5	Table 14.1.1.5.2: Summary of Other Baseline Characteristics	Safety Population 2
6	Table 14.1.2.1: Summary of Medical History by System Organ Class and Preferred Term	Safety Population 2
7	Table 14.1.5.1 [PK]: Summary of [Analyte] in Plasma PK Parameters	PK Population
8	Table 14.2.1.1 [ONC]: Overall Survival	Efficacy Population
9	Table 14.2.1.2 [CNS]: Analysis of Change from Baseline in Parameter by Visit using MMRM	Efficacy Population
10	Table 14.2.1.2 [ONC]: Progression Free Survival by RECIST 1.1 [or irRC or imRECIST]	Efficacy Population
11	Table 14.2.1.3 [CNS]: Subgroup Analysis of Parameter at Visit by Subgroup using MMRM	Efficacy Population
12	Table 14.2.1.5 [CNS]: Analysis of Responder Endpoint at Visit	Efficacy Population
13	Table 14.3.1.1.1: Overall Summary of Treatment-Emergent Adverse Events	Safety Population 2
14	Table 14.3.1.1.2: Overall Summary of Treatment-Emergent Adverse Events of Interest	Safety Population 2
15	Table 14.3.1.2.1: Summary of Treatment-Emergent Adverse Events by System Organ Class and Preferred Term	Safety Population 2
16	Table 14.3.1.4.1: Summary of Treatment-Emergent Serious Adverse Events by System Organ Class and Preferred Term	Safety Population 2
17	Table 14.3.1.6: Summary of Treatment-Emergent Adverse Events Leading to Death by System Organ Class and Preferred Term	Safety Population 2
18	Table 14.3.2.1: Summary of Observed and Change from Baseline	Safety Population 2

	by Scheduled Visits - Hematology Laboratory Test	
19	Table 14.3.3.1: Shift from Baseline to Worst Post-Baseline Grade - Hematology Laboratory Test	Safety Population 2
20	Table 14.3.3.2: Shift from Baseline to Worst Post-Baseline Grade - Chemistry Laboratory Test	Safety Population 2
21	Table 14.3.3.3: Shift from Baseline to Worst Post-Baseline Grade - Coagulation Laboratory Test	Safety Population 2
22	Table 14.3.5.1: Summary of Observed and Change from Baseline by Scheduled Visits - Electrocardiogram	Safety Population 2
23	Table 14.7.1 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Screening (Lifetime Assessment)	Efficacy Population
24	Table 14.7.3 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Baseline	Efficacy Population
25	Table 14.7.4 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at any Postbaseline Assessment	Efficacy Population

Table 14.1.1.1

Summary of Analysis Populations

All Enrolled Population

Analysis Population, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Safety	XX (XX.X)	XX (XX.X)	XX (XX.X)
Intent-To-Treat (ITT)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Full Analysis Set	XX (XX.X)	XX (XX.X)	XX (XX.X)
Per-Protocol	XX (XX.X)	XX (XX.X)	XX (XX.X)
Completed Study	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Enrolled Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). N in column headers represents number of subjects entered in study (i.e., signed informed consent). The ITT population includes all subjects randomized. The Safety population includes all randomized subjects known to have taken at least one dose of randomized study drug. The Full Analysis Set population includes all subjects in the safety population who also have at least one post-baseline ADAS-Cog and CIBIC+assessment.

Source: adxx; Program name; Program run date/time

Table 14.1.1.3

Summary of Subject Disposition

Safety Population 2

Disposition Information, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Completed Study Treatment	XX (XX.X)	XX (XX.X)	XX (XX.X)
Discontinued Study Treatment			
Adverse Event	XX (XX.X)	XX (XX.X)	XX (XX.X)
Pregnancy	XX (XX.X)	XX (XX.X)	XX (XX.X)
Progressive Disease	XX (XX.X)	XX (XX.X)	XX (XX.X)
Protocol Deviation	XX (XX.X)	XX (XX.X)	XX (XX.X)
Study Terminated by Sponsor	XX (XX.X)	XX (XX.X)	XX (XX.X)
Site Terminated by Sponsor	XX (XX.X)	XX (XX.X)	XX (XX.X)
Withdrawal by Subject	XX (XX.X)	XX (XX.X)	XX (XX.X)
Other	XX (XX.X)	XX (XX.X)	XX (XX.X)
Completed Study	XX (XX.X)	XX (XX.X)	XX (XX.X)
Discontinued Study			
Death	XX (XX.X)	XX (XX.X)	XX (XX.X)
Adverse Event	XX (XX.X)	XX (XX.X)	XX (XX.X)
Lost to Follow-up	XX (XX.X)	XX (XX.X)	XX (XX.X)
Pregnancy	XX (XX.X)	XX (XX.X)	XX (XX.X)
Progressive Disease	XX (XX.X)	XX (XX.X)	XX (XX.X)
Protocol Deviation	XX (XX.X)	XX (XX.X)	XX (XX.X)
Screen Failure	XX (XX.X)	XX (XX.X)	XX (XX.X)

Study Terminated by Sponsor	XX (XX.X)	XX (XX.X)	XX (XX.X)
Site Terminated by Sponsor	XX (XX.X)	XX (XX.X)	XX (XX.X)
Withdrawal by Subject	XX (XX.X)	XX (XX.X)	XX (XX.X)
Other	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Note: Percentages were calculated as 100 × (n/N).

Source: adxx; Program name; Program run date/time

Table 14.1.1.4.1

Summary of Eligibility Criteria Deviations

All Enrolled Population

Category, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number of Subjects who did not meet at least one Eligibility Criterion	XX (XX.X)	XX (XX.X)	XX (XX.X)
Number of Subjects who did not meet at least one Inclusion Criteria	XX (XX.X)	XX (XX.X)	XX (XX.X)
<Criterion # / Short Description>	XX (XX.X)	XX (XX.X)	XX (XX.X)
Number of Subjects who met at least one Exclusion Criteria	XX (XX.X)	XX (XX.X)	XX (XX.X)
<Criterion # / Short Description>	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Enrolled Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Each subject is counted once per category, regardless of the number of criteria not met/met.

Source: adxx; Program name; Program run date/time

Table 14.1.1.4.2

Summary of Key Protocol Deviations

Safety Population 2

Category, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number of Subjects with at least one Protocol Deviation	XX (XX.X)	XX (XX.X)	XX (XX.X)
Number of Subjects with at least one Major Protocol Deviation [a]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Type of Major Protocol Deviation			
Deviations Category #1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Deviations Category #2	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Note: Percentages were calculated as 100 × (n/N).

[a] A major protocol deviation is defined as any departure from the approved protocol that impacted the safety, rights, and/or welfare of the patient; negatively impacted the quality or completeness of the data; or made the informed consent document/form inaccurate.

Source: adxx; Program name; Program run date/time

Table 14.1.1.5.2

Summary of Other Baseline Characteristics

Safety Population 2

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**	**Total (N=xxx)**	**Treatment A vs Treatment B P Value**
{Other Characteristic 1}				0.XXXX
n	XX	XX	XX	
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)	
Median	XX.X	XX.X	XX.X	
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	
{Other Characteristic 2}				0.XXXX
n	XX	XX	XX	
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)	
Median	XX.X	XX.X	XX.X	
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	
{Other Characteristic 3}				0.XXXX
n	XX	XX	XX	
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)	
Median	XX.X	XX.X	XX.X	
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	

Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X	

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.
Note: Percentages were calculated as 100 × (n/N).
Source: adxx; Program name; Program run date/time

Table 14.1.2.1

Summary of Medical History by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term [a], n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number of Subjects with at least one Medical History	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term.

[a] All investigators adverse events were coded using MedDRA version xx.x.

Source: adxx; Program name; Program run date/time

Table 14.1.5.1 [PK]

Summary of [Analyte] in Plasma PK Parameters

PK Population

PK Parameter	**Treatment Group A (N=XXX)**	**Treatment Group B (N=XXX)**	**Total (N=xxx)**
Global Cmax ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
Early Cmax ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
Late Cmax ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
Global Tmax (h) ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)

Early Tmax (h) ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
Late Tmax (h) ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
AUC 0-last ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
AUC 0-∞ ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
CL/F ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)
Vd/F ([units])			
n	XX	XX	XX
Mean (SD)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)
GM (CV%)	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.XXX)

Median (Min, Max)	X.XX (X.X, X.X)	X.XX (X.X, X.X)	X.XX (X.X, X.X)

Note: N=the number of subjects in each treatment group in the PK Population.

Abbreviations: AUC 0-∞=Area under the concentration-time curve from time zero to infinity. AUC 0-last=Area under the concentration-time curve from time zero to the last quantifiable concentration. C_{max}=Maximum observed concentration. CL/F=Apparent oral clearance. CV%=coefficient of variation. GM=geometric mean. Max=maximum. Min=minimum. n=number of participants. PK=pharmacokinetic(s). SD=standard deviation. T_{max}=Time of maximum observed concentration. Vd/F=Apparent volume of distribution during the terminal phase.

Table 14.2.1.1 [ONC]

Overall Survival

Efficacy Population

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Number (%) of Participants with Events (Death)	XX (XX.X)	XX (XX.X)
Number (%) of Participants Censored [a]	XX (XX.X)	XX (XX.X)
Ongoing	XX (XX.X)	XX (XX.X)
Discontinued Study	XX (XX.X)	XX (XX.X)
Kaplan-Meier Estimate of OS (Months) (95% CI) [b]		
Q1	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
Median	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
Q3	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
Kaplan-Meier Estimate of OS Rate (95% CI)		
At {xx Months}	0.XXX (0.XXX, 0.XXX)	0.XXX (0.XXX, 0.XXX)
At {xx Months}	0.XXX (0.XXX, 0.XXX)	0.XXX (0.XXX, 0.XXX)
{Stratified} Hazard Ratio {of A vs B} (95% CI) [c]	0.XX (0.XX, NEst)	0.XX (0.XX, NEst)
P value (Non-Inferiority Test)	0.XXX	0.XXX
P value	0.XXX	0.XXX

P value (Non-Inferiority Test) [d]	0.XXX	
P value [e]	0.XXX	
Duration of OS Follow-up (Months) [f]		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
>= *{xx Months}*	XX (XX.X)	XX (XX.X)
>= *{xx Months}*	XX (XX.X)	XX (XX.X)
>= *{xx Months}*	XX (XX.X)	XX (XX.X)

[a] {Definition of “Ongoing” and “Discontinued Study” categories, and censoring rules.}

[b] OS (months) = (date of death or censoring – date of {randomization/first dose} + 1)/30.4375, and the 95% CI is derived using log-log transformation.

[c] Hazard ratio and 95% CIs are calculated using {the Cox proportional hazards model / statistical method}{, adjusted for randomization stratification factors (xxx) / without any adjustment}.

[d] P value for the non-inferiority test is based on {statistical method}, with hypothesis {H0: HR >= xx vs. H1: HR < xx}.

[e] P value for comparing the overall survival functions from the two treatment groups is from {stratified log-rank test, adjusted for randomization stratification factors (xxx) / unstratified log-rank test}.

[f] Duration of OS follow-up (months) = (date of death or censoring – date of {randomization/first dose} + 1)/30.4375.

Table 14.2.1.2 [CNS]

Analysis of Change from Baseline in Parameter by Visit using MMRM

Efficacy Population

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Number of Participants in the Analysis	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Change from Baseline at <<Visit>>		
LS Mean (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
LS Mean Difference from Placebo (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
P-value	0.XXXX	0.XXXX
Change from Baseline at <<Visit>>		
LS Mean (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
LS Mean Difference from Placebo (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
P-value	0.XXXX	0.XXXX

Abbreviations: CI = confidence interval; LS = least squares; MMRM = mixed-effect model for repeated measures; SD = standard deviation; SE = standard error; <<add other abbreviations/acronyms>>.

Note: LS means are estimated from an MMRM which includes <<add definition from SAP>>.

Table 14.2.1.2 [ONC]

Progression Free Survival by RECIST 1.1 [or irRC or imRECIST]

Efficacy Population

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Progression Free Survival (PFS) [a]		
Number of death events, n (%)	XX (XX.XX)	XX (XX.XX)
RECIST 1.1 Progression	XX (XX.XX)	XX (XX.XX)
Death	XX (XX.XX)	XX (XX.XX)
Median	XX.X	XX.X
Min, Max	XX, XX	XX, XX
K-M Median (95% CI) [b]	XX.X (X.X, X.X)	XX.X (X.X, X.X)
25%, 75%	XX, XX	XX, XX

Note: The efficacy analysis set consists of all subjects who receive at least 1 infusion of any amount of XXX.

[a]: Overall Survival is defined as time from treatment start until death from any reason. Subjects who terminate the study without documented death will be censored on the date of the last visit/contact.

[b]: The 95% CIs for K-M estimate were calculated using Brookmeyer Crowley method.

Table 14.2.1.3 [CNS]

Subgroup Analysis of Parameter at Visit by Subgroup using MMRM

Efficacy Population

Subgroup Statistic	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
<<Name of Subgroup 1 (e.g., Female)>>		
Number of Participants in the Analysis	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Change from Baseline at <<Visit>>		
LS Mean (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
LS Mean Difference from Placebo (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
P-value	0.XXXX	0.XXXX
<<Name of Subgroup 2 (e.g., Male)>>		
Number of Participants in the Analysis	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
n	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Median	XX.X	XX.X
Min, Max	XX, XX	XX, XX
Change from Baseline at <<Visit>>		
LS Mean (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)

LS Mean Difference from Placebo (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
P-value	0.XXXX	0.XXXX

Abbreviations: CI = confidence interval; LS = least squares; MMRM = mixed-effect model for repeated measures; SD = standard deviation; SE = standard error; <<add other abbreviations/acronyms>>.

Note: <<Add footnote describing the subgroup analysis>>.

Table 14.2.1.5 [CNS]

Analysis of Responder Endpoint at Visit

Efficacy Population

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Number of Participants in the Analysis	XX	XX
<<Response>> [a], n (%)	XX (XX.X)	XX (XX.X)
<<Non-response>>, n (%)	XX (XX.X)	XX (XX.X)
P-value compared to TRT1 [b]		0.XXXX
P-value	0.XXXX	0.XXXX

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Note: Percentages are based on the number of participants in the <Analysis Set>.

[a] [add definition of responder/non-responder, and approaches to handle certain intercurrent events or missing data as defined in the SAP].

[b] P-value is based on <<test>>.

Table 14.3.1.1.1

Overall Summary of Treatment-Emergent Adverse Events

Safety Population 2

Categories, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}			
TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
Serious TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related Serious TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Death	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related TEAE Leading to Death	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Dose Modification [a]	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Treatment Discontinuation	XX (XX.X)	XX (XX.X)	XX (XX.X)

[a] Dose Modification includes Dose Reduced; Drug Interrupted in the AE action taken with study treatment.

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.1.2

Overall Summary of Treatment-Emergent Adverse Events of Interest

Safety Population 2

Categories, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}			
TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE with Grade 3 or [Higher][4]	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE with Grade 2 or [Higher][, 3, or 4]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related TEAE with Grade 3 or [Higher][4]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related TEAE with Grade 2 or [Higher][, 3 or 4]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Serious TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
Related Serious TEAE	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to [Premature] Discontinuation of Study Drug	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Dose Interruption of Study Drug	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Dose Modification of Study Drug	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Dose	XX (XX.X)	XX (XX.X)	XX (XX.X)

Modification or Interruption of Study Drug			
TEAE Leading to Dose Reduction of Study Drug	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to [Premature] Discontinuation of Study	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE with Dose-Limiting Toxicity	XX (XX.X)	XX (XX.X)	XX (XX.X)
TEAE Leading to Death	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}. Death includes any death that occurred during the study. Percentages for TEAE Dose Limiting Toxicity was calculated based on the number of subjects in the DLT Evaluable Set.

Source: adxx; Program name; Program run date/time

Table 14.3.1.2.1

Summary of Treatment-Emergent Adverse Events by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.4.1

Summary of Treatment-Emergent Serious Adverse Events by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.6

Summary of Treatment-Emergent Adverse Events Leading to Death by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

EXPORT-STUDY-01

Table 14.3.2.1

Summary of Observed and Change from Baseline by Scheduled Visits - Hematology Laboratory Test

Safety Population 2

Parameter (Units) **Visit**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Parameter [results terminology]			
Baseline			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n Change from Baseline>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: Max=Maximum; Min=Minimum; N=Number of patients in the Safety Set; n=Number of patients in the specific category; Q1=Quartile 1 (the first quartile); Q3=Quartile 3 (the third quartile); SD=Standard Deviation.

Notes: Percentages were calculated as 100 × (n/N). Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. Change from baseline is defined as post-baseline value minus baseline value.

Source: adxx; Program name; Program run date/time

Programming Notes: Repeat for all analysis parameters by each analysis visits as per the results terminology provided.

Table 14.3.3.1

Shift from Baseline to Worst Post-Baseline Grade - Hematology Laboratory Test

Safety Population 2

Parameter (unit) Grade at Baseline, n (%)	**Treatment A**			**Treatment B**		
	Worst Grade Post-Baseline			**Worst Grade Post-Baseline**		
	Normal	**Abnormal (NCS)**	**Abnormal (CS)**	**Normal**	**Abnormal (NCS)**	**Abnormal (CS)**
Parameter [results terminology]	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Normal	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (NCS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (CS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)

Abbreviations: CS=clinically significant; n=Number of subjects in the specific category; NCS=not clinically significant.

Notes: Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. The denominator for calculating percentages will be based on the number of Safety subjects with measurements on both occasions (baseline and post-baseline) for that parameter.

Source: adxx; Program name; Program run date/time

Programming Notes: A shift table from baseline to worst post-baseline grade according to normal range criteria (for quantitative measurements and categorical measurements) will be summarized by visit for each laboratory parameter. The denominator for calculating percentages will be based on the number of safety population with measurements on both occasions (baseline and post-baseline) for that parameter.

Table 14.3.3.2

Shift from Baseline to Worst Post-Baseline Grade - Chemistry Laboratory Test

Safety Population 2

Parameter (unit) Grade at Baseline, n (%)	**Treatment A**			**Treatment B**		
	Worst Grade Post-Baseline			**Worst Grade Post-Baseline**		
	Normal	**Abnormal (NCS)**	**Abnormal (CS)**	**Normal**	**Abnormal (NCS)**	**Abnormal (CS)**
Parameter [results terminology]						
Normal	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (NCS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (CS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)

Abbreviations: CS=clinically significant; n=Number of subjects in the specific category; NCS=not clinically significant.

Notes: Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. The denominator for calculating percentages will be based on the number of Safety subjects with measurements on both occasions (baseline and post-baseline) for that parameter.

Source: adxx; Program name; Program run date/time

Programming Notes: A shift table from baseline to worst post-baseline grade according to normal range criteria (for quantitative measurements and categorical measurements) will be summarized by visit for each laboratory parameter. The denominator for calculating percentages will be based on the number of safety participants with measurements on both occasions (baseline and post-baseline) for that parameter.

Table 14.3.3.3

Shift from Baseline to Worst Post-Baseline Grade - Coagulation Laboratory Test

Safety Population 2

Parameter (unit) Grade at Baseline, n (%)	**Treatment A**			**Treatment B**		
	Worst Grade Post-Baseline			**Worst Grade Post-Baseline**		
	Normal	**Abnormal (NCS)**	**Abnormal (CS)**	**Normal**	**Abnormal (NCS)**	**Abnormal (CS)**
Parameter [results terminology]						
Normal	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (NCS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (CS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)

Abbreviations: CS=clinically significant; n=Number of subjects in the specific category; NCS=not clinically significant.

Notes: Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. The denominator for calculating percentages will be based on the number of Safety subjects with measurements on both occasions (baseline and post-baseline) for that parameter.

Source: adxx; Program name; Program run date/time

Programming Notes: A shift table from baseline to worst post-baseline grade according to normal range criteria (for quantitative measurements and categorical measurements) will be summarized by visit for each laboratory parameter. The denominator for calculating percentages will be based on the number of safety participants with measurements on both occasions (baseline and post-baseline) for that parameter.

Table 14.3.5.1

Summary of Observed and Change from Baseline by Scheduled Visits - Electrocardiogram

Safety Population 2

Parameter (Units) **Visit**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Parameter [results terminology]			
Baseline			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n Change from Baseline>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: Max=Maximum; Min=Minimum; N=Number of patients in the Safety Set; n=Number of patients in the specific category; Q1=Quartile 1 (the first quartile); Q3=Quartile 3 (the third quartile); SD=Standard Deviation.

Notes: Percentages were calculated as 100 × (n/N). Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. Change from baseline is defined as post-baseline value minus baseline value. All 12-lead ECG parameters are at the averages of triplicate readings.

Source: adxx; Program name; Program run date/time

Programming Notes: Repeat for all analysis parameters by each analysis visits as per the results terminology provided.

Table 14.7.1 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Screening (Lifetime Assessment)

Efficacy Population

Category/Item	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
Number of Participants with Data	XX	XX
Suicidal Ideation [a], n (%)	XX (XX.X)	XX (XX.X)
1 - Wish to be dead	XX (XX.X)	XX (XX.X)
2 - Non-specific active suicidal thoughts	XX (XX.X)	XX (XX.X)
3 - Active suicidal ideation with any methods (not plan) without intent to act	XX (XX.X)	XX (XX.X)
4 - Active suicidal ideation with some intent to act, without specific plan	XX (XX.X)	XX (XX.X)
5 - Active suicidal ideation with specific plan and intent	XX (XX.X)	XX (XX.X)
Suicidal Behavior [b], n (%)	XX (XX.X)	XX (XX.X)
6 - Preparatory acts or behavior	XX (XX.X)	XX (XX.X)
7 - Aborted attempt	XX (XX.X)	XX (XX.X)
8 - Interrupted attempt	XX (XX.X)	XX (XX.X)
9 - Non-fatal suicide attempt	XX (XX.X)	XX (XX.X)
Suicidal Ideation or Behavior [c], n (%)	XX (XX.X)	XX (XX.X)

Notes: Data shown (n) are number of participants with a “Yes” response to the specified item or category. Percentages are based on the number of participants in the <<analysis set>> with data.

[a] A "Yes" response to any of items 1 - 5.

[b] A "Yes" response to any of items 6 - 9.

[c] A "Yes" response to any of items 1 - 9.

Table 14.7.3 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Baseline

Efficacy Population

Category/Item	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
Number of Participants with Data	XX	XX
Suicidal Ideation [a], n (%)	XX (XX.X)	XX (XX.X)
1 - Wish to be dead	XX (XX.X)	XX (XX.X)
2 - Non-specific active suicidal thoughts	XX (XX.X)	XX (XX.X)
3 - Active suicidal ideation with any methods (not plan) without intent to act	XX (XX.X)	XX (XX.X)
4 - Active suicidal ideation with some intent to act, without specific plan	XX (XX.X)	XX (XX.X)
5 - Active suicidal ideation with specific plan and intent	XX (XX.X)	XX (XX.X)
Suicidal Behavior [b], n (%)	XX (XX.X)	XX (XX.X)
6 - Preparatory acts or behavior	XX (XX.X)	XX (XX.X)
7 - Aborted attempt	XX (XX.X)	XX (XX.X)
8 - Interrupted attempt	XX (XX.X)	XX (XX.X)
9 - Non-fatal suicide attempt	XX (XX.X)	XX (XX.X)
10 - Completed suicide	XX (XX.X)	XX (XX.X)
Suicidal Ideation or Behavior [c], n (%)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Notes: Data shown (n) are number of participants with a “Yes” response to the specified item or category based on "Since Last Visit" questionnaire.

Percentages are based on the number of participants in the <<analysis set>> with data.

[a] A "Yes" response to any of items 1 - 5.

[b] A "Yes" response to any of items 6 - 10.

[c] A "Yes" response to any of items 1 - 10.

Table 14.7.4 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at any Postbaseline Assessment

Efficacy Population

Category/Item	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
Number of Participants with Data	XX	XX
Suicidal Ideation [a], n (%)	XX (XX.X)	XX (XX.X)
1 - Wish to be dead	XX (XX.X)	XX (XX.X)
2 - Non-specific active suicidal thoughts	XX (XX.X)	XX (XX.X)
3 - Active suicidal ideation with any methods (not plan) without intent to act	XX (XX.X)	XX (XX.X)
4 - Active suicidal ideation with some intent to act, without specific plan	XX (XX.X)	XX (XX.X)
5 - Active suicidal ideation with specific plan and intent	XX (XX.X)	XX (XX.X)
Suicidal Behavior [b], n (%)	XX (XX.X)	XX (XX.X)
6 - Preparatory acts or behavior	XX (XX.X)	XX (XX.X)
7 - Aborted attempt	XX (XX.X)	XX (XX.X)
8 - Interrupted attempt	XX (XX.X)	XX (XX.X)
9 - Non-fatal suicide attempt	XX (XX.X)	XX (XX.X)
10 - Completed suicide	XX (XX.X)	XX (XX.X)
Suicidal Ideation or Behavior [c], n (%)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Notes: Data shown (n) are number of participants with a “Yes” response to the specified item or category at any postbaseline visit.

Percentages are based on the number of participants in the <<analysis set>> with data.

[a] A "Yes" response to any of items 1 - 5.

[b] A "Yes" response to any of items 6 - 10.

[c] A "Yes" response to any of items 1 - 10.