Table of Contents

S.no	Title	Population Set
1	Table 14.1.5.2 [PK]: Summary of [Analyte] in Plasma PK Parameters (Horizontal)	PK Population
2	Table 14.3.2.2: Summary of Observed and Change from Baseline by Scheduled Visits - Chemistry Laboratory Test	Safety Population 2
3	Table 14.3.1.5.2: Summary of Treatment-Emergent Drug-Related Adverse Events by System Organ Class, Preferred Term, and Severity	Safety Population 2
4	Table 14.3.1.5.1: Summary of Treatment-Emergent Drug-Related Adverse Events by System Organ Class and Preferred Term	Safety Population 2
5	Table 14.3.1.3.3: Summary of Grade 3 or [Higher] Treatment-Emergent Adverse Events by System Organ Class and Preferred Term	Safety Population 2
6	Table 14.3.6.1: Shift from Baseline to Worst Post-Baseline Grade - Overall ECG Interpretation	Safety Population 2
7	Table 14.3.1.3.2: Summary of Treatment-Emergent Adverse Events by Preferred Term and Severity	Safety Population 2
8	Table 14.1.3.2: Summary of Concomitant Medications by Anatomic Classification and Preferred Term	Safety Population 2
9	Table 14.1.1.2: Summary of Enrollment by Country and Investigator	All Enrolled Population
10	Table 14.7.6 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS) Suicidal Ideation Score: Shift from Baseline Category to Maximum Postbaseline Category	Efficacy Population
11	Table 14.7.2 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Screening (Past <<x>> Months)	Efficacy Population
12	Table 14.2.1.4 [CNS]: Summary of Responder Endpoint by Visit	Efficacy Population
13	Table 14.2.1.1 [CNS]: Analysis of Change from Baseline in Parameter at Visit using MMRM	Efficacy Population
14	Table 14.2.2.1 [ONC]: Observed Tumor Burden, Change from	Efficacy Population

	Baseline, and Percent Change from Baseline by Scheduled Visits - RECIST 1.1	
15	Table 14.3.4.1: Summary of Observed and Change from Baseline by Scheduled Visits - Vital Signs	Safety Population 2
16	Table 14.3.2.3: Summary of Observed and Change from Baseline by Scheduled Visits - Coagulation Laboratory Test	Safety Population 2
17	Table 14.3.1.7: Summary Treatment-Emergent Adverse Events Leading to [Premature] Discontinuation of Study Drug by System Organ Class and Preferred Term	Safety Population 2
18	Table 14.3.1.5.3: Summary of Treatment-Emergent Drug-Related Serious Adverse Events by System Organ Class and Preferred Term	Safety Population 2
19	Table 14.3.1.4.2: Summary of Treatment-Emergent Serious Adverse Events by Preferred Term	Safety Population 2
20	Table 14.3.1.3.1: Summary of Treatment-Emergent Adverse Events by System Organ Class, Preferred Term, and Severity	Safety Population 2
21	Table 14.3.1.2.2: Summary of Treatment-Emergent Adverse Events by Preferred Term	Safety Population 2
22	Table 14.1.4.1: Extent of Exposure to Study Drug	Safety Population 2
23	Table 14.1.3.1: Summary of Prior Medications by Anatomic Classification and Preferred Term	Safety Population 2
24	Table 14.1.1.5.1: Summary of Demographic and Baseline Characteristics	Safety Population 2
25	Table 14.7.5 [CNS]: Columbia-Suicide Severity Rating Scale (C-SSRS) Suicidal Ideation Score: Shift from Baseline to Maximum Postbaseline Value	Efficacy Population

Table 14.1.5.2 [PK]

Summary of [Analyte] in Plasma PK Parameters (Horizontal)

PK Population

Characteristics	Treatment Group A (N=XXX)				Treatment Group A (N=XXX)			
	n	Mean (SD)	GM (CV%)	Median (Min, Max)	n	Mean (SD)	GM (CV%)	Median (Min, Max)
Parameter X	XX	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.X, X.X)	XX	X.XX (X.XXX)	X.XX (X.XXX)	X.XX (X.X, X.X)

Abbreviations: CV%=coefficient of variation. GM=geometric mean. Max=maximum. Min=minimum. n=number of participants. PK=pharmacokinetic(s). SD=standard deviation.

Note: N=the number of subjects in each treatment group in the PK Population.

Table 14.3.2.2

Summary of Observed and Change from Baseline by Scheduled Visits - Chemistry Laboratory Test

Safety Population 2

Parameter (Units) **Visit**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Parameter [results terminology]			
Baseline			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n Change from Baseline>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: Max=Maximum; Min=Minimum; N=Number of patients in the Safety Set; n=Number of patients in the specific category; Q1=Quartile 1 (the first quartile); Q3=Quartile 3 (the third quartile); SD=Standard Deviation.

Notes: Percentages were calculated as 100 × (n/N). Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. Change from baseline is defined as post-baseline value minus baseline value.

Source: adxx; Program name; Program run date/time

Programming Notes: Repeat for all analysis parameters by each analysis visits as per the results terminology provided.

Table 14.3.1.5.2

Summary of Treatment-Emergent Drug-Related Adverse Events by System Organ Class, Preferred Term, and Severity

Safety Population 2

System Organ Class Preferred Term Grade, n(%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)

Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: Grade 1 = Mild; Grade 2 = Moderate; Grade 3 = Severe; Grade 4 = Life-Threatening; Grade 5 = Death; N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once for the highest severity grade within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Programming Notes: Present SOCs alphabetically and PTs within SOC present by alphabetical order.

Table 14.3.1.5.1

Summary of Treatment-Emergent Drug-Related Adverse Events by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.3.3

Summary of Grade 3 or [Higher] Treatment-Emergent Adverse Events by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: Grade 3 = Severe; Grade 4 = Life-Threatening; Grade 5 = Death; N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once for the highest severity grade within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Programming Notes: Present SOCs alphabetically and PTs within SOC present by alphabetical order.

Table 14.3.6.1

Shift from Baseline to Worst Post-Baseline Grade - Overall ECG Interpretation

Safety Population 2

Parameter (unit) Grade at Baseline, n (%)	**Treatment A**			**Treatment B**		
	Worst Grade Post-Baseline			**Worst Grade Post-Baseline**		
	Normal	**Abnormal (NCS)**	**Abnormal (CS)**	**Normal**	**Abnormal (NCS)**	**Abnormal (CS)**
Parameter [results terminology]						
Normal	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (NCS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)
Abnormal (CS)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)	XX (XX.XX)

Abbreviations: CS=clinically significant; ECG=electrocardiogram; n=Number of subjects in the specific category; NCS=not clinically significant.

Notes: Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. The denominator for calculating percentages will be based on the number of Safety subjects with measurements on both occasions (baseline and post-baseline) for that parameter.

Source: adxx; Program name; Program run date/time

Programming Notes: A shift table from baseline to worst post-baseline grade according to normal range criteria (for quantitative measurements and categorical measurements) will be summarized by visit for each laboratory parameter. The denominator for calculating percentages will be based on the number of safety participants with measurements on both occasions (baseline and post-baseline) for that parameter.

Table 14.3.1.3.2

Summary of Treatment-Emergent Adverse Events by Preferred Term and Severity

Safety Population 2

Preferred Term Severity, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, descending order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: Grade 1 = Mild; Grade 2 = Moderate; Grade 3 = Severe; Grade 4 = Life-Threatening; Grade 5 = Death; N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once for the highest severity grade within each preferred term. Treatment-emergent events began on or after the study drug start date up

to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Programming Notes: Present PTs by descending order of the [total]{or specify} frequencies.

Table 14.1.3.2

Summary of Concomitant Medications by Anatomic Classification and Preferred Term

Safety Population 2

Anatomic Class **Preferred Term [a], n (%)**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Number of Subjects with at least one Concomitant Medications	XX (XX.X)	XX (XX.X)	XX (XX.X)
Anatomic Class [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each ATC class. A medication can be counted in multiple ATC classes. Concomitant medications are defined as any medication that was taken on or after the date of first dose of study drug.

[a] Concomitant Medications are coded using WHODrug dictionary version YYYY-XX.

Source: adxx; Program name; Program run date/time

Table 14.1.1.2

Summary of Enrollment by Country and Investigator

All Enrolled Population

Country **Investigator Name (Investigator ID), n (%)**	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Country <n>	XX (XX.X)	XX (XX.X)	XX (XX.X)
Investigator <n> (Investigator ID <n>)	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Investigators who enrolled participants were included. Investigator at the time of [randomization][enrollment] was summarized for participants who changed sites after enrollment.

Source: adxx; Program name; Program run date/time

Table 14.7.6 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS) Suicidal Ideation Score: Shift from Baseline Category to Maximum Postbaseline Category

Efficacy Population

Treatment	Baseline Category	**Treatment Category**			**Total**
		No suicidal ideation or behavior n (%)	**Suicidal ideation n (%)**	**Suicidal behavior n (%)**	**n (%)**
Treatment A (N=XXX)	No suicidal ideation or behavior	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Suicidal ideation	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Suicidal behavior	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Total	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Treatment B (N=XXX)	No suicidal ideation or behavior	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Suicidal ideation	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Suicidal behavior	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Total	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Notes: Suicidal Ideation includes any one of the five suicidal ideation events (Categories 1-5). Suicidal

behavior includes any one of the five suicidal behavior events (Categories 6-10).

Each participant is counted in one cell only. Participants with both Suicidal Ideation and Suicidal Behavior are included in the Suicidal Behavior category.

Percentages are based on the number of participants in the <<analysis set>> with data at baseline and at least one postbaseline assessment.

Table 14.7.2 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS): Reports of Suicidal Ideation or Behavior at Screening (Past <<x>> Months)

Efficacy Population

Category/Item	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
Number of Participants with Data	XX	XX
Suicidal Ideation [a], n (%)	XX (XX.X)	XX (XX.X)
1 - Wish to be dead	XX (XX.X)	XX (XX.X)
2 - Non-specific active suicidal thoughts	XX (XX.X)	XX (XX.X)
3 - Active suicidal ideation with any methods (not plan) without intent to act	XX (XX.X)	XX (XX.X)
4 - Active suicidal ideation with some intent to act, without specific plan	XX (XX.X)	XX (XX.X)
5 - Active suicidal ideation with specific plan and intent	XX (XX.X)	XX (XX.X)
Suicidal Behavior [b], n (%)	XX (XX.X)	XX (XX.X)
6 - Preparatory acts or behavior	XX (XX.X)	XX (XX.X)
7 - Aborted attempt	XX (XX.X)	XX (XX.X)
8 - Interrupted attempt	XX (XX.X)	XX (XX.X)
9 - Non-fatal suicide attempt	XX (XX.X)	XX (XX.X)
Suicidal Ideation or Behavior [c], n (%)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Notes: Data shown (n) are number of participants with a “Yes” response to the specified item or category.

Percentages are based on the number of participants in the <<analysis set>> with data.

The lookback period is the past <<xx>> months.

[a] A "Yes" response to any of items 1 - 5.

[b] A "Yes" response to any of items 6 - 9.

[c] A "Yes" response to any of items 1 - 9.

Table 14.2.1.4 [CNS]

Summary of Responder Endpoint by Visit

Efficacy Population

Visit Statistic	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
<PBL #1>		
<<Response>>, n (%)	XX (XX.X)	XX (XX.X)
<<Non-response>>, n (%)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. PBL = Postbaseline)>>

Note: Percentages are based on the number of participants in the <Analysis Set>.

Table 14.2.1.1 [CNS]

Analysis of Change from Baseline in Parameter at Visit using MMRM

Efficacy Population

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Number of Participants in the Analysis	XX	XX
Mean (SD)	XX.X (XX.X)	XX.X (XX.X)
Change from Baseline at <<Visit>>		
LS Mean (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
LS Mean Difference from Placebo (SE)	XX.X (XX.X)	XX.X (XX.X)
95% CI	(XX.X, XX.X)	(XX.X, XX.X)
P-value	0.XXXX	0.XXXX

Abbreviations: CI = confidence interval; LS = least squares; MMRM = mixed-effect model for repeated measures; SD = standard deviation; SE = standard error; <<add other abbreviations/acronyms>>.

Note: LS means are estimated from an MMRM which includes <<add definition from SAP>>.

Table 14.2.2.1 [ONC]

Observed Tumor Burden, Change from Baseline, and Percent Change from Baseline by Scheduled Visits - RECIST 1.1

Efficacy Population

Measurement Statistics	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**
Baseline [a]		
n	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
<Cycle X Day Y>		
n	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
Change from baseline to <Cycle X Day Y>		
n	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
Percent change from baseline to <Cycle X Day Y>		

n	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X

Notes: [Definition of analysis set].

[a] Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product.

Table 14.3.4.1

Summary of Observed and Change from Baseline by Scheduled Visits - Vital Signs

Safety Population 2

Parameter (Units) **Visit**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Parameter [results terminology]			
Baseline			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n Change from Baseline>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: Max=Maximum; Min=Minimum; N=Number of patients in the Safety Set; n=Number of patients in the specific category; Q1=Quartile 1 (the first quartile); Q3=Quartile 3 (the third quartile); SD=Standard Deviation.

Notes: Percentages were calculated as 100 × (n/N). Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. Change from baseline is defined as post-baseline value minus baseline value.

Source: adxx; Program name; Program run date/time

Programming Notes: Repeat for all analysis parameters by each analysis visits as per the results terminology provided.

Table 14.3.2.3

Summary of Observed and Change from Baseline by Scheduled Visits - Coagulation Laboratory Test

Safety Population 2

Parameter (Units) **Visit**	**Treatment Group A** **(N=xxx)**	**Treatment Group B** **(N=xxx)**	**Total** **(N=xxx)**
Parameter [results terminology]			
Baseline			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
<Visit n Change from Baseline>			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: Max=Maximum; Min=Minimum; N=Number of patients in the Safety Set; n=Number of patients in the specific category; Q1=Quartile 1 (the first quartile); Q3=Quartile 3 (the third quartile); SD=Standard Deviation.

Notes: Percentages were calculated as 100 × (n/N). Baseline is defined as the last assessment that is non-missing prior to first dose of investigational product. Change from baseline is defined as post-baseline value minus baseline value.

Source: adxx; Program name; Program run date/time

Programming Notes: Repeat for all analysis parameters by each analysis visits as per the results terminology provided.

Table 14.3.1.7

Summary Treatment-Emergent Adverse Events Leading to [Premature] Discontinuation of Study Drug by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.5.3

Summary of Treatment-Emergent Drug-Related Serious Adverse Events by System Organ Class and Preferred Term

Safety Population 2

System Organ Class Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.3.1.4.2

Summary of Treatment-Emergent Serious Adverse Events by Preferred Term

Safety Population 2

Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, descending order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Programming Notes: Present PTs by descending order of the [total]{or specify} frequencies.

Table 14.3.1.3.1

Summary of Treatment-Emergent Adverse Events by System Organ Class, Preferred Term, and Severity

Safety Population 2

System Organ Class Preferred Term Grade, n(%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)
SOC [value, alphabetical order]			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]			
Grade 1	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 2	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 3	XX (XX.X)	XX (XX.X)	XX (XX.X)

Grade 4	XX (XX.X)	XX (XX.X)	XX (XX.X)
Grade 5	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: Grade 1 = Mild; Grade 2 = Moderate; Grade 3 = Severe; Grade 4 = Life-Threatening; Grade 5 = Death; N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once for the highest severity grade within each system organ class and preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Programming Notes: Present SOCs alphabetically and PTs within SOC present by alphabetical order.

Table 14.3.1.2.2

Summary of Treatment-Emergent Adverse Events by Preferred Term

Safety Population 2

Preferred Term, n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number (%) of Participants with Any {type of adverse events summarized in table title}	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, descending order of the total frequencies]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category; TEAE=Treatment-Emergent Adverse Events.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each preferred term. Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach}. PTs are presented by descending order of the total frequencies.

Adverse events were coded according to MedDRA Version {XX.X}.

Source: adxx; Program name; Program run date/time

Table 14.1.4.1

Extent of Exposure to Study Drug

Safety Population 2

	Treatment Group A (N=xxx)	**Treatment Group B (N=xxx)**
Total Duration of Exposure to Study Drug ([days][weeks][months])		
n	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
Cumulative N (%) of Subjects Exposed to Study Drug		
{>= xx [days][weeks][months]}	XX (XX.X)	XX (XX.X)
{>= yy [days][weeks][months]}	XX (XX.X)	XX (XX.X)
{>= zz [days][weeks][months]}	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Total duration of exposure to study drug = last dosing date - first dosing date + {1 for daily dose, 7 for weekly dose, 14 for bi-weekly dose}.

Source: adxx; Program name; Program run date/time

Table 14.1.3.1

Summary of Prior Medications by Anatomic Classification and Preferred Term

Safety Population 2

Anatomic Class Preferred Term [a], n (%)	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Number of Subjects with at least one Prior Medications	XX (XX.X)	XX (XX.X)	XX (XX.X)
Anatomic Class [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)
Preferred Term [value, alphabetical order]	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Notes: Percentages were calculated as 100 × (n/N). Subjects are counted once within each ATC class. A medication can be counted in multiple ATC classes. Prior medications are defined as medications that stopped before the date and time of the first dose.

[a] Prior Medications are coded using WHODrug dictionary version YYYY-XX.

Source: adxx; Program name; Program run date/time

Table 14.1.1.5.1

Summary of Demographic and Baseline Characteristics

Safety Population 2

Characteristics	**Treatment Group A (N=xxx)**	**Treatment Group B (N=xxx)**	**Total (N=xxx)**
Age			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Age Group			
< 65 yrs	XX (XX.X)	XX (XX.X)	XX (XX.X)
65 to 80 yrs	XX (XX.X)	XX (XX.X)	XX (XX.X)
> 80 yrs	XX (XX.X)	XX (XX.X)	XX (XX.X)
Gender			
Female	XX (XX.X)	XX (XX.X)	XX (XX.X)
Male	XX (XX.X)	XX (XX.X)	XX (XX.X)
Primary Race			
American Indian or Alaska Native	XX (XX.X)	XX (XX.X)	XX (XX.X)
Asian	XX (XX.X)	XX (XX.X)	XX (XX.X)
Black or African American	XX (XX.X)	XX (XX.X)	XX (XX.X)
Native Hawaiian or Other Pacific Islander	XX (XX.X)	XX (XX.X)	XX (XX.X)
Not Reported	XX (XX.X)	XX (XX.X)	XX (XX.X)

Other	XX (XX.X)	XX (XX.X)	XX (XX.X)
Unknown	XX (XX.X)	XX (XX.X)	XX (XX.X)
Ethnicity			
Hispanic or Latino	XX (XX.X)	XX (XX.X)	XX (XX.X)
Not Hispanic or Latino	XX (XX.X)	XX (XX.X)	XX (XX.X)
Baseline Height (cm)			XX (XX.XX)
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Baseline Weight (lbs)			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Baseline BMI (kg/m2)			
n	XX	XX	XX
Mean (SD)	XX.X (XX.XX)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X	XX.X, XX.X

Abbreviations: N=Number of subjects in the Safety Population; n=Number of subjects in a specific category.

Note: Percentages were calculated as 100 × (n/N).

Source: adxx; Program name; Program run date/time

Table 14.7.5 [CNS]

Columbia-Suicide Severity Rating Scale (C-SSRS) Suicidal Ideation Score: Shift from Baseline to Maximum Postbaseline Value

Efficacy Population

Treatment	Baseline Score	**Maximum Postbaseline Suicidal Ideation Score**						
		0 **n (%)**	**1** **n (%)**	**2** **n (%)**	**3** **n (%)**	**4** **n (%)**	**5** **0n (%)**	**Total** **n (%)**
Treatment A (N=XXX)	0	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	1	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	2	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	3	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	4	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	5	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Total	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
Treatment B (N=XXX)	0	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	1	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	2	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	3	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	4	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	5	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)
	Total	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)	XX (XX.X)

Abbreviations: <<add abbreviations/acronyms (e.g. CFB = change from baseline)>>

Notes: Suicidal ideation score: 0 = No suicidal ideation; 1 = Wish to be dead; 2 = Non-specific active suicidal thoughts; 3 = Active suicidal ideation with any methods (not plan) without intent to act; 4 = Active suicidal ideation with some intent to act, without specific plan; 5 = Active suicidal ideation with specific plan and intent.

Percentages are based on the number of participants in the <<analysis set>> with data at baseline and at least one postbaseline assessment.