


4DM50: Dynamics and Control of Cooperation

Network synchronization II

Erik Steur
e.steur@tue.nl



TU/e

Technische Universiteit
Eindhoven
University of Technology

Recap

Partial synchronization

Partial synchronization with delays

Role of network topology


Systems of the form

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

with $CB > 0$ and diffusive coupling functions

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij}(y_j - y_i)$$

where diffusion coefficients $\gamma_{ij} = \gamma_{ji} > 0$ and $j \in \mathcal{N}_i \Leftrightarrow i \in \mathcal{N}_j$



Diffusive coupling matrix

$$\Gamma = \begin{pmatrix} \sum_{j=2}^N \gamma_{1j} & -\gamma_{12} & \cdots & -\gamma_{1N} \\ -\gamma_{21} & \sum_{j=1, j \neq 2}^N \gamma_{2j} & \ddots & \vdots \\ \vdots & \ddots & \ddots & -\gamma_{(N-1)N} \\ -\gamma_{N1} & \cdots & -\gamma_{N(N-1)} & \sum_{j=1}^{N-1} \gamma_{Nj} \end{pmatrix}$$

with $\gamma_{ij} = 0$ if (and only if) $j \notin \mathcal{N}_i$

- ▶ Γ is singular
- ▶ $\Gamma = \Gamma^T$ is positive semi-definite
- ▶ Eigenvalues of Γ (if and only if the network is [connected](#)):

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

The systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

are **strictly semi-passive** with a radially unbounded storage function S

The systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

are **strictly semi-passive** with a radially unbounded storage function S

That is: there exists $S : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that the dissipation inequality

$$\dot{S} \leq y_i^T u_i$$

is strict for all x_i outside some ball $\mathcal{B} \subset \mathbb{R}^n$

The systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

are **strictly semi-passive** with a radially unbounded storage function S

That is: there exists $S : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that the dissipation inequality

$$\dot{S} \leq y_i^T u_i$$

is strict for all x_i outside some ball $\mathcal{B} \subset \mathbb{R}^n$

Result: networks of diffusively coupled strictly semi-passive systems have uniformly ultimately bounded solutions

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

with CB (similar to) a positive definite matrix

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

with CB (similar to) a positive definite matrix in new coordinates

$$\begin{cases} \dot{z}_i = q(z_i, y_i) \\ \dot{y}_i = a(z_i, y_i) + CBu_i \end{cases}$$

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

with CB (similar to) a positive definite matrix in new coordinates

$$\begin{cases} \dot{z}_i = q(z_i, y_i) \\ \dot{y}_i = a(z_i, y_i) + CBu_i \end{cases}$$

Assumption: There exists a $(n - m) \times (n - m)$ matrix $P = P^T > 0$ such that the eigenvalues of

$$P \left(\frac{\partial q}{\partial z}(z, w) \right) + \left(\frac{\partial q}{\partial z}(z, w) \right)^T P$$

are negative and separated away from zero for all $(z, w) \in \mathbb{R}^{n-m} \times \mathcal{W}$

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases}$$

with CB (similar to) a positive definite matrix in new coordinates

$$\begin{cases} \dot{z}_i = q(z_i, y_i) \\ \dot{y}_i = a(z_i, y_i) + CBu_i \end{cases}$$

Assumption: There exists a $(n - m) \times (n - m)$ matrix $P = P^T > 0$ such that the eigenvalues of

$$P \left(\frac{\partial q}{\partial z}(z, w) \right) + \left(\frac{\partial q}{\partial z}(z, w) \right)^T P$$

are negative and separated away from zero for all $(z, w) \in \mathbb{R}^{n-m} \times \mathcal{W}$
 \Rightarrow subsystem $\dot{z}_i = q(z_i, y_i)$ is **exponentially convergent** w.r.t. input y_i

Given a network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (1)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (2)$$

Given a network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (1)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (2)$$

Assumptions:

- ▶ system (8) is strictly semi-passive with a radially unbounded storage function S
- ▶ the subsystem

$$\dot{z}_i = q(z_i, w(t))$$

associated to (8) satisfies the test for exponential convergence

Given a network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (1)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (2)$$

Result:

- ▶ solutions of the diffusively coupled systems are uniformly ultimately bounded
- ▶ there exists a positive constant $\bar{\lambda}$ such that for

$$\lambda_2 \geq \bar{\lambda} \quad (\lambda_2 \text{ is the smallest non-zero eigenvalue of } \Gamma)$$

Then the set $\mathcal{M} := \{x \in \mathbb{R}^{Nn} \mid x_1 = x_2 = \dots = x_N\}$ contains a globally asymptotically stable subset, i.e. the systems synchronize

Extensions to

- ▶ Directed networks

Extensions to

- ▶ Directed networks
- ▶ Time-delay coupling


$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t))$$

and

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t - \tau))$$

Extensions to

- ▶ Directed networks
- ▶ Time-delay coupling



Recap

Partial synchronization

Partial synchronization with delays

Role of network topology

What if a network does not fully synchronize?

What if a network does not fully synchronize?

- ▶ no synchronization of any pair of systems

What if a network does not fully synchronize?

- ▶ no synchronization of any pair of systems
- ▶ spatiotemporal patterns (e.g. traveling/rotating waves)


What if a network does not fully synchronize?

- ▶ no synchronization of any pair of systems
- ▶ spatiotemporal patterns (e.g. traveling/rotating waves)
- ▶ some form of incomplete network synchronization

What if a network does not fully synchronize?

- ▶ no synchronization of any pair of systems
- ▶ spatiotemporal patterns (e.g. traveling/rotating waves)
- ▶ some form of incomplete network synchronization

Partial synchronization = asymptotic match of the states of some, but not all systems



Partial synchronization =
existence of partial synchronization manifold \mathcal{P}

Partial synchronization =
existence of partial synchronization manifold \mathcal{P}
+ asymptotic stability of \mathcal{P}

Partial synchronization =
existence of partial synchronization manifold \mathcal{P}
+ asymptotic stability of \mathcal{P}

Questions:

- ▶ How to find \mathcal{P} ?


Partial synchronization =
existence of partial synchronization manifold \mathcal{P}
+ asymptotic stability of \mathcal{P}

Questions:

- ▶ How to find \mathcal{P} ?
- ▶ When is \mathcal{P} stable?


Look for **symmetries**

- ▶ in the coupling configuration (network)
- ▶ in the system dynamics




Look for **symmetries**


- ▶ in the coupling configuration (network)
- ▶ in the system dynamics



Rearrangement of some nodes of the network leave the network unchanged




Original network




Rearranged network

Rearrangement of some nodes of the network leave the network unchanged



Original network



Rearranged network

Permutation $\pi : \{1, \dots, N\} \rightarrow \{1, \dots, N\}$

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ \pi(1) = 2 & \pi(2) = 1 & \pi(3) = 4 & \pi(4) = 3 \end{pmatrix}$$

Permutation matrix associated to π

$$\Pi = (\Pi_{ij})$$

with

$$\Pi_{ij} = \begin{cases} 1 & \text{if } \pi(i) = j \\ 0 & \text{otherwise} \end{cases}$$

Permutation matrix associated to π

$$\Pi = (\Pi_{ij})$$

with

$$\Pi_{ij} = \begin{cases} 1 & \text{if } \pi(i) = j \\ 0 & \text{otherwise} \end{cases}$$

Result: if the permutation matrix Π and the coupling matrix Γ commute,

$$\Pi\Gamma = \Gamma\Pi,$$

then the set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$

is **invariant** w.r.t. dynamics of the coupled systems

- ▶ The set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$

is described by equations of the form

$$x_i = x_j \quad (\text{for } \pi(i) = j)$$

- ▶ The set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$

is described by equations of the form

$$x_i = x_j \quad (\text{for } \pi(i) = j)$$

- ▶ A sufficient and **necessary** condition for the set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$

to be invariant is that there exists a matrix X that solves the matrix equation

$$(I_N - \Pi)\Gamma = X(I_N - \Pi)$$

- ▶ The set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$

is described by equations of the form

$$x_i = x_j \quad (\text{for } \pi(i) = j)$$


- ▶ A sufficient and **necessary** condition for the set

$$\ker(I_{Nn} - \Pi \otimes I_n)$$


to be invariant is that there exists a matrix X that solves the matrix equation

$$(I_N - \Pi)\Gamma = X(I_N - \Pi)$$


Π and L commute $\Rightarrow X = \Gamma$ is a solution




Original network



Rearranged network




Original network




Rearranged network

Permutation $\pi : \{1, \dots, 4\} \rightarrow \{1, \dots, 4\}$

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ \pi(1) = 2 & \pi(2) = 1 & \pi(3) = 4 & \pi(4) = 3 \end{pmatrix} \Rightarrow \Pi = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$




Original network




Rearranged network

$$\Pi \Gamma = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$




Original network




Rearranged network

$$= \begin{pmatrix} -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \end{pmatrix} =$$



Original network



Rearranged network

$$\begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} = \Gamma \Pi$$

Other permutation matrices that commute with

$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$

are:

$$\Pi_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

Other permutation matrices that commute with

$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$

are:


$$\Pi_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

and

$$\Pi_3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

Look for **symmetries**

- ▶ in the coupling configuration (network)
- ▶ in the system dynamics



Not only the network may contain symmetries

Not only the network may contain symmetries

Suppose

- ▶ there is a matrix J such that

$$Jf(x_i) = f(Jx_i)$$

Not only the network may contain symmetries

Suppose

- ▶ there is a matrix J such that

$$Jf(x_i) = f(Jx_i)$$

- ▶ J commutes with BC : $J(BC) = (BC)J$

Not only the network may contain symmetries

Suppose

- ▶ there is a matrix J such that

$$Jf(x_i) = f(Jx_i)$$

- ▶ J commutes with BC : $J(BC) = (BC)J$
- ▶ there is a permutation matrix Π that commutes with Γ :

$$\Pi\Gamma = \Gamma\Pi$$

Not only the network may contain symmetries

Suppose

- ▶ there is a matrix J such that

$$Jf(x_i) = f(Jx_i)$$

- ▶ J commutes with BC : $J(BC) = (BC)J$
- ▶ there is a permutation matrix Π that commutes with Γ :

$$\Pi\Gamma = \Gamma\Pi$$

Then the set

$$\ker(I_{Nn} - \Pi \otimes J)$$

is invariant w.r.t. dynamics of the coupled systems

Example: Lorenz system with input u and output $y = x_1$:

$$\begin{cases} \dot{x}_1 = \sigma(x_2 - x_1) + u \\ \dot{x}_2 = x_1(\rho - x_3) - x_2 \\ \dot{x}_3 = -\gamma x_3 + x_1 x_2 \end{cases}$$

Example: Lorenz system with input u and output $y = x_1$:

$$\begin{cases} \dot{x}_1 = \sigma(x_2 - x_1) + u \\ \dot{x}_2 = x_1(\rho - x_3) - x_2 \\ \dot{x}_3 = -\gamma x_3 + x_1 x_2 \end{cases}$$

has symmetry defined by

$$J = \begin{pmatrix} -1 & & \\ & -1 & \\ & & 1 \end{pmatrix}$$

Example: Lorenz system with input u and output $y = x_1$:

$$\begin{cases} \dot{x}_1 = \sigma(x_2 - x_1) + u \\ \dot{x}_2 = x_1(\rho - x_3) - x_2 \\ \dot{x}_3 = -\gamma x_3 + x_1 x_2 \end{cases}$$

has symmetry defined by

$$J = \begin{pmatrix} -1 & & \\ & -1 & \\ & & 1 \end{pmatrix}$$

The set $\ker(I_{Nn} - \Pi \otimes J)$ is invariant for coupled Lorenz systems as

$$J(BC) = \begin{pmatrix} -1 & & \\ & -1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & & \\ & 0 & \\ & & 0 \end{pmatrix} = (BC)J$$

Network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (3)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (4)$$

Network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (3)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (4)$$

Assumptions:

- ▶ system (8) is strictly semi-passive with a radially unbounded storage function S
- ▶ the subsystem

$$\dot{z}_i = q(z_i, w(t))$$

associated to (8) satisfies the test for exponential convergence

- ▶ there is a permutation matrix Π that commutes with Γ

Network of diffusively coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (3)$$

$$u_i = \sum_{j \in \mathcal{N}_i} \gamma_{ij} (y_j - y_i) \quad (4)$$


Result:

- ▶ solutions of the diffusively coupled systems are uniformly ultimately bounded
- ▶ there exists a positive constant $\bar{\lambda}$ such that for


$$\lambda' \geq \bar{\lambda} \quad (\lambda' \text{ is the smallest eigenvalue of } \Gamma \text{ with eigenvector in } \text{range}(I_N - \Pi))$$

Then $\ker(I_{Nn} - \Pi \otimes I_n)$ contains a globally asymptotically stable subset

$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$




$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$



Recall: The following permutation matrices commute with Γ

$$\Pi_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \Pi_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad \Pi_3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$


$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$



Eigenvalues and eigenvectors of Γ

$$\begin{array}{llll} \lambda_1 = 0 & \lambda_2 = 2\gamma_1 & \lambda_3 = 2\gamma_2 & \lambda_4 = 2\gamma_1 + 2\gamma_2 \\ v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} & v_2 = \begin{pmatrix} 1 \\ -1 \\ -1 \\ 1 \end{pmatrix} & v_3 = \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} & v_4 = \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} \end{array}$$

$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$




- ▶ $v_2 \in \text{range}(I_n - \Pi_1), \text{range}(I_n - \Pi_3)$
- ▶ $v_3 \in \text{range}(I_n - \Pi_2), \text{range}(I_n - \Pi_3)$
- ▶ $v_4 \in \text{range}(I_n - \Pi_1), \text{range}(I_n - \Pi_2)$

hence

- ▶ $\Pi_1: \lambda' = 2\gamma_1$
- ▶ $\Pi_2: \lambda' = 2\gamma_2$
- ▶ $\Pi_3: \lambda' = \min(2\gamma_1, 2\gamma_2)$


$$\Gamma = \begin{pmatrix} \gamma_1 + \gamma_2 & -\gamma_1 & 0 & -\gamma_2 \\ -\gamma_1 & \gamma_1 + \gamma_2 & -\gamma_2 & 0 \\ 0 & -\gamma_2 & \gamma_1 + \gamma_2 & -\gamma_1 \\ -\gamma_2 & 0 & -\gamma_1 & \gamma_1 + \gamma_2 \end{pmatrix}$$



- ▶ $v_2 \in \text{range}(I_n - \Pi_1), \text{range}(I_n - \Pi_3)$
- ▶ $v_3 \in \text{range}(I_n - \Pi_2), \text{range}(I_n - \Pi_3)$
- ▶ $v_4 \in \text{range}(I_n - \Pi_1), \text{range}(I_n - \Pi_2)$

hence

- ▶ $\Pi_1: \lambda' = 2\gamma_1 \Rightarrow x_1 = x_2 \text{ and } x_3 = x_4 \text{ if } \gamma_1 > \gamma_2 \text{ is suff. large}$
- ▶ $\Pi_2: \lambda' = 2\gamma_2 \Rightarrow x_1 = x_2 \text{ and } x_3 = x_4 \text{ if } \gamma_2 > \gamma_1 \text{ is suff. large}$
- ▶ $\Pi_3: \lambda' = \min(2\gamma_1, 2\gamma_2) \Rightarrow \text{full sync}$



Recap

Partial synchronization

Partial synchronization with delays

Role of network topology

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (5)$$

with diffusive time-delay coupling (transmission delay)

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t)) \quad (6)$$

or (sensor/actuator delay)

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t)(t - \tau)) \quad (7)$$

Systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (5)$$

with diffusive time-delay coupling (transmission delay)

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t)) \quad (6)$$

or (sensor/actuator delay)

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t)(t - \tau)) \quad (7)$$

For now: Coupled systems (5), (7) with $a_{ij} = a_{ji}$ and $j \in \mathcal{N}_i \Leftrightarrow i \in \mathcal{N}_j$
(similar results are obtained for directed networks and/or coupled systems (5)-(6))

Coupling matrix

$$L = \begin{pmatrix} \sum_{j=2}^N a_{1j} & -a_{12} & \cdots & -a_{1N} \\ -a_{21} & \sum_{j=1, j \neq 2}^N a_{2j} & \ddots & \vdots \\ \vdots & \ddots & \ddots & -a_{(N-1)N} \\ -a_{N1} & \cdots & -a_{N(N-1)} & \sum_{j=1}^{N-1} a_{Nj} \end{pmatrix}$$

with $a_{ij} = 0$ if (and only if) $j \notin \mathcal{N}_i$

Coupling matrix

$$L = \begin{pmatrix} \sum_{j=2}^N a_{1j} & -a_{12} & \cdots & -a_{1N} \\ -a_{21} & \sum_{j=1, j \neq 2}^N a_{2j} & \ddots & \vdots \\ \vdots & \ddots & \ddots & -a_{(N-1)N} \\ -a_{N1} & \cdots & -a_{N(N-1)} & \sum_{j=1}^{N-1} a_{Nj} \end{pmatrix}$$

with $a_{ij} = 0$ if (and only if) $j \notin \mathcal{N}_i$

Note that

$$u(t) = -\sigma(L \otimes I_m)y(t - \tau)$$

(compare with the delay-free case!)

Finding modes of partial synchronization

27/35

Look for **symmetries** in the coupling matrix L

Look for **symmetries** in the coupling matrix L

Result: if a permutation matrix Π and the coupling matrix L commute,

$$\Pi L = L \Pi,$$

then the set

$$\{\phi \in \mathcal{C} \mid \phi(\theta) \in \ker(I_{Nn} - \Pi \otimes I_n), \quad -\tau \leq \theta \leq 0\}$$

is **forward invariant** w.r.t. dynamics of the coupled systems

Eigenvalues of L :

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

Eigenvalues of L :

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

Given a permutation matrix Π , denote:

- ▶ λ' the **smallest** eigenvalue of L with eigenvector in $\text{range}(I_N - \Pi)$

Eigenvalues of L :

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

Given a permutation matrix Π , denote:

- ▶ λ' the **smallest** eigenvalue of L with eigenvector in $\text{range}(I_N - \Pi)$
- ▶ λ^* the **largest** eigenvalue of L with eigenvector in $\text{range}(I_N - \Pi)$

Eigenvalues of L :


$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

Given a permutation matrix Π , denote:


- ▶ λ' the **smallest** eigenvalue of L with eigenvector in $\text{range}(I_N - \Pi)$
- ▶ λ^* the **largest** eigenvalue of L with eigenvector in $\text{range}(I_N - \Pi)$

Assumptions:


- ▶ system (5) is strictly semi-passive with a radially unbounded storage function S
- ▶ the subsystem $\dot{z}_i = q(z_i, w(t))$ associated to (5) satisfies the test for exponential convergence
- ▶ the permutation matrix Π that commutes with L




$$\lambda' = \lambda_2 \text{ and } \lambda^* = \lambda_N$$



$$\lambda' > \lambda_2 \text{ and } \lambda^* = \lambda_N$$




$$\lambda' = \lambda_2 \text{ and } \lambda^* < \lambda_N$$




$$\lambda' > \lambda_2 \text{ and } \lambda^* < \lambda_N$$

$$L = \begin{pmatrix} a_1 + a_2 & -a_1 & 0 & -a_2 \\ -a_1 & a_1 + a_2 & -a_2 & 0 \\ 0 & -a_2 & a_1 + a_2 & -a_1 \\ -a_2 & 0 & -a_1 & a_1 + a_2 \end{pmatrix}$$




$$L = \begin{pmatrix} a_1 + a_2 & -a_1 & 0 & -a_2 \\ -a_1 & a_1 + a_2 & -a_2 & 0 \\ 0 & -a_2 & a_1 + a_2 & -a_1 \\ -a_2 & 0 & -a_1 & a_1 + a_2 \end{pmatrix}$$



Eigenvalues and eigenvectors of L if $a_1 \leq a_2$

$$\begin{array}{llll} \lambda_1 = 0 & \lambda_2 = 2a_1 & \lambda_3 = 2a_2 & \lambda_4 = 2a_1 + 2a_2 \\ v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} & v_2 = \begin{pmatrix} 1 \\ -1 \\ -1 \\ 1 \end{pmatrix} & v_3 = \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} & v_4 = \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} \end{array}$$

$$L = \begin{pmatrix} a_1 + a_2 & -a_1 & 0 & -a_2 \\ -a_1 & a_1 + a_2 & -a_2 & 0 \\ 0 & -a_2 & a_1 + a_2 & -a_1 \\ -a_2 & 0 & -a_1 & a_1 + a_2 \end{pmatrix}$$




Recall: The following permutation matrix commutes with Γ

$$\Pi_3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

and $v_2, v_3 \in \text{range}(I_n - \Pi_3)$

$$L = \begin{pmatrix} a_1 + a_2 & -a_1 & 0 & -a_2 \\ -a_1 & a_1 + a_2 & -a_2 & 0 \\ 0 & -a_2 & a_1 + a_2 & -a_1 \\ -a_2 & 0 & -a_1 & a_1 + a_2 \end{pmatrix}$$



Recall: The following permutation matrix commutes with Γ

$$\Pi_3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

and $v_2, v_3 \in \text{range}(I_n - \Pi_3) \Rightarrow \lambda' = 2a_1$ and $\lambda^* = 2a_2 < \lambda_4 = 2a_1 + 2a_2$

Recap

Partial synchronization

Partial synchronization with delays

Role of network topology

Network of diffusively time-delay coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (8)$$

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij} (y_j(t - \tau) - y_i(t - \tau)) \quad (9)$$

with $a_{ij} = a_{ji}$ and $j \in \mathcal{N}_i \Leftrightarrow i \in \mathcal{N}_j$

Network of diffusively time-delay coupled systems

$$\begin{cases} \dot{x}_i = f(x_i) + Bu_i \\ y_i = Cx_i \end{cases} \quad (8)$$

$$u_i(t) = \sigma \sum_{j \in \mathcal{N}_i} a_{ij}(y_j(t - \tau) - y_i(t - \tau)) \quad (9)$$


with $a_{ij} = a_{ji}$ and $j \in \mathcal{N}_i \Leftrightarrow i \in \mathcal{N}_j$

Assumptions:

- ▶ system (8) is strictly semi-passive with a radially unbounded storage function S
- ▶ the subsystem

$$\dot{z}_i = q(z_i, w(t))$$

associated to (8) satisfies the test for exponential convergence



Recall:

$$L = \begin{pmatrix} \sum_{j=2}^N a_{1j} & -a_{12} & \cdots & -a_{1N} \\ -a_{21} & \sum_{j=1, j \neq 2}^N a_{2j} & \ddots & \vdots \\ \vdots & \ddots & \ddots & -a_{(N-1)N} \\ -a_{N1} & \cdots & -a_{N(N-1)} & \sum_{j=1}^{N-1} a_{Nj} \end{pmatrix}$$

with $a_{ij} = 0$ if (and only if) $j \notin \mathcal{N}_i$

Eigenvalues of L :

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$$

- ▶ Suppose two coupled systems synchronize for

$$(\sigma, \tau) \in \mathcal{S}$$

- ▶ Suppose two coupled systems synchronize for

$$(\sigma, \tau) \in \mathcal{S}$$

- ▶ Determine **scaled copies** of \mathcal{S} :

$$\mathcal{S}_j^* := \left\{ (\sigma, \tau) \in \mathbb{R}_+ \times \overline{\mathbb{R}}_+ \mid \left(\frac{\lambda_j}{2} \sigma, \tau \right) \in \mathcal{S} \right\}$$

with scaling factors the eigenvalues of L

- ▶ Suppose two coupled systems synchronize for

$$(\sigma, \tau) \in \mathcal{S}$$

- ▶ Determine **scaled copies** of \mathcal{S} :

$$\mathcal{S}_j^* := \left\{ (\sigma, \tau) \in \mathbb{R}_+ \times \overline{\mathbb{R}}_+ \mid \left(\frac{\lambda_j}{2} \sigma, \tau \right) \in \mathcal{S} \right\}$$

with scaling factors the eigenvalues of L

- ▶ **Result:** Network synchronizes if

$$(\sigma, \tau) \in \mathcal{S}_2^* \cap \mathcal{S}_N^*$$

