



Biopesticide

Mode of Action Unknown or Uncertain

Contact, Ingestion and Inhalation



TECHNICAL SHEET – Plant Protection

Exemption from registration under EPA FIFRA 25(b)

Classification: Botanical Insecticide and Miticide.
Active Ingredient: Garlic oil 2.00%, Cinnamon oil 1.00%, Clove oil 1.00%, and Thyme oil 1.00% wt.
Inert Ingredient: Water, Soap, Lecithins, and Xanthan gum 95.00% wt.
Formulation: Oil in Water Emulsion.

Instructions for use

Authorized for use in Safe Programs

Crop	Pest	Rate (fl.oz./Acre) *
Avocado (NL)	Thrips (<i>Frankliniella occidentalis</i>)	19 - 67 ¹
Garlic, Onion, Chives, Asparagus (NL)	Thrips (<i>Thrips tabaci</i>)	32 - 64 ¹
Celery, Chard, Spinach, Lettuce (NL)	Aphid (<i>Myzus persicae</i>)	29 - 111 ¹
Blueberry, Raspberry, Strawberry, Currant, Blackberry (NL)	Leaf roller (<i>Argyrotaenia montezumae</i>)	18 - 43 ¹
	Green Peach Aphid (<i>Myzus persicae</i>)	32 - 64 ¹
	Spotted Wing Vinegar Fly (<i>Drosophila suzukii</i>)	32 - 64 ¹
	Thrips (<i>Frankliniella occidentalis</i>)	47 - 226 ¹
	Red berry mite (<i>Acalitus essigi</i>)	21 - 49 ¹
Eggplant, Chili Pepper, Habanero	White fly (<i>Bemisia tabaci</i>)	23 - 36 ¹
Pepper, Potato, Bell Pepper, Tomatillo (NL)	Colorado Potato Beetle (<i>Leptinotarsa decemlineata</i>)	32 - 64 ¹
	Eastern Thrips (<i>Thrips palmi</i>)	16 - 53 ¹
Potato (NL)	Nematodes (<i>Meloidogyne chitwoodi</i> , <i>Pratylenchus</i> sp., <i>Tylenchorhynchus</i> sp.)	64 - 128 ⁴
	Potato Aphid (<i>Myzus persicae</i>)	64 - 128 ⁴
Broccoli, Cabbage, Brussel Sprouts, Cauliflower, Kohlrabi, Rape (NL)	Aphid (<i>Brevicoryne brassicae</i>)	28 - 110 ¹
Summer Squash, Squash, Chayote, Melon, Cucumber, Watermelon (NL)	White fly (<i>Bemisia tabaci</i>)	30 - 47 ¹
	Eastern Thrips (<i>Thrips palmi</i>)	94 - 225 ¹
Mango (NL)	Thrips (<i>Frankliniella</i> sp.)	32 - 63 ¹
Pineapple (NL)	Black weevil (<i>Metamasius callizona</i>)	39 - 116 ³
Grape (NL)	Thrips (<i>Frankliniella occidentalis</i>)	71 - 145 ⁴
Citrus: Lime, Lemon, Mandarin, Orange, Tangerine, Grapefruit (NL)	Two-Spotted Mite (<i>Tetranychus urticae</i>)	52 ¹
Almond, Pistachio (NL)	Navel Orangeworm (<i>Amyelois transitella</i>)	80 ⁵
Pome Fruit: Apple, Pear (NL)	Aphid (<i>Aphis pomi</i>)	32 - 64 ¹
	Two-Spotted Mite (<i>Tetranychus urticae</i>)	32 - 64 ¹
	Thrips (<i>Frankliniella occidentalis</i>)	32 - 64 ¹
	Psylla (<i>Cacopsylla bidens</i>)	32 - 64 ¹
	Whitefly (<i>Bemisia tabaci</i>)	32 - 64 ¹
Stone Fruit: Apricot, Cherry, Nectarine, Peach, Plum (NL)	Aphid (<i>Aphis pomi</i>)	32 - 64 ¹
	Two-Spotted Mite (<i>Tetranychus urticae</i>)	32 - 64 ¹
	Thrips (<i>Frankliniella occidentalis</i>)	32 - 64 ¹
	Psylla (<i>Cacopsylla bidens</i>)	32 - 64 ¹
	Whitefly (<i>Bemisia tabaci</i>)	32 - 64 ¹

* Volume of application 18 - 100 gallons/acre.

¹ Make 3 foliar applications, at 7-day intervals.

² Make 5 foliar applications at 7-day intervals, beginning at start of flowering.

³ Make 5 foliar applications at drip line at intervals of 7, 10, 11 and 8 days, starting 50 days after floral induction.

⁴ Make 3 foliar applications at drip line at 7-day intervals.

⁵ Make 3 applications to foliage at 7 to 14-day intervals.

All ingredients in this product are exempt from the requirement of a tolerance under 40 CFR 180.

Its components act as an insecticide and a repellent against insects

Re-Entry Intervals (REI): Four (4) hours after application.

Preharvest Interval (PHI): 0 days, No Limit (NL).




Interval between applications: 7, 8, 10, and 11 days.

Method to prepare the product: Shake product before using. Condition the water to between 5.5 and 6.0. Pour the required quantity of product directly into the water. If mixing with other products always in the proper order of addition until the mixture is homogenized.


Application of the product: Apply the spray solution on the same day it was prepared. Perform applications when the first ovipositions are observed or when the first infestation is observed, preferably the first nymphal instars which are the most susceptible. Ensure that the product contacts the insects and that the size of the drop and the pressure allow coverage of the areas where the pest is located. Apply higher doses to serious infestations.

Incompatibility: It is not compatible with oxidizing substances and/or alkaline substances. When mixing with other products, a compatibility test must be performed prior to application. Do not mix with products whose compatibility is not completely tested.


Contraindications: Do not apply to plants that are stressed by inadequate irrigation, in extreme temperatures, during hours of intense sunshine, when it is raining or when wind could cause drift.



- Do not apply during high temperatures and low relative humidity.
- Perform applications before 10 a.m. and after 4 p.m.



- Alternative to prevent or delay risk of resistance.
- Does not share a common action point with other compounds and can be freely applied.
- Effective in sequences, alternations or mixtures with insecticides of different modes of action.



The information contained in this document is solely for your consultation, without legal value or guarantee. For more information related to the product, contact Tel. (777) 3211475 / 01 800 1127 500 to 502. ptigrupoultraquimia.com



For more information call
 US Customer Service Toll Free Number:
 1(800) 801-9657 in Mexico y (777) 319 35 45
 55 7980 2287

Created:
 Updated:
 Version:
 Page:

June 16, 2023.
 March 21, 2024.
 01.
 1 of 1