

Biopesticide

Mode of Action Unknown or Uncertain
Contact, Ingestion and Inhalation



TECHNICAL SHEET – Plant Protection

Exemption from registration under EPA FIFRA 25(b)

Classification: Botanical Insecticide and Miticide.
Active Ingredient: Neem oil 55.00%, and Cinnamon oil 5.00% wt.
Inert Ingredient : 40.00% wt.
Formulation: Emulsifiable Concentrate

Instructions for use Authorized for use in Safe Programs

Crop	Pest	Rate (gal/Acre) *
Pepper, Habanero Pepper, Tomato, Sweet Pepper, Tomatillo, Eggplant, Potato (NL)	Whitefly (<i>Bemisia tabaci</i>)	0.16 - 0.21 ¹
Tomato, Eggplant, Pepper, Sweet Pepper, Potato, Tomatillo (NL)	Red Spider Mite (<i>Tetranychus urticae</i>) Tomato Russet Mite (<i>Aculops lycopersici</i>) Colorado Potato Beetle (<i>Leptinotarsa decemlineata</i>)	0.11 - 0.32 ¹
Garlic, Onion, Chives, Asparagus (NL)	Thrips (<i>Thrips tabaci</i>) European Asparagus Aphid (<i>Brachycorynella asparagi</i>)	0.11 - 0.21 ²
Zucchini, Pumpkin, Chayote, Melon, Cucumber, Watermelon (NL)	Western Flower Thrips (<i>Frankliniella occidentalis</i>)	0.16 - 0.21 ¹
Cranberry, Strawberry, Raspberry, Blackberry, Redcurrant, Dewberry, Smilax (NL)	Red Spider Mite (<i>Tetranychus urticae</i>)	0.05 - 0.29 ¹
Vine (NL)	Thrips (<i>Frankliniella occidentalis</i>)	0.11 - 0.32 ¹
Pome Fruit: Apple, Pear, Quince Tejocotel Stone Fruit: Apricot, Cherry, Nectarine, Peach, Plum (NL)	Aphid (<i>Eriosoma lanigerum</i> , <i>Aphis pomi</i>) Two-Spotted Mite (<i>Tetranychus sp.</i>) Thrips (<i>Frankliniella occidentalis</i>) Psylla (<i>Cacopsylla bidens</i>) Whitefly (<i>Bemisia tabaci</i>)	0.19 - 0.22 ¹

* Volume of application 18 – 100 gallons/acre.

¹ Make 3 foliar applications, at 7-day intervals, when the first infestations are detected.

² Make 2 foliar applications, at 8-day intervals, when the first infestations are detected.

All ingredients in this product are exempt from the requirement of a tolerance under 40 CFR 180. Its components act as an insecticide and a repellent against insects

Re-Entry Intervals (REI): Four (4) hours after application, unless wearing the appropriate PPE.

Preharvest Interval (PHI): 0 days, No Limit (NL).


Interval between applications: 7 and 8 days.

Method to prepare the product: Shake product before using. Adjust the pH of the water to between 5 and 6. Pour the required quantity of product directly into the water. If mixing with other products always in the proper order of addition until the mixture is homogenized.


Application of the product: Apply the spray solution on the same day it was prepared. Perform applications when the first ovipositions are observed or when the first infestation is observed, preferably the first nymphal instars which are the most susceptible. Ensure that the product contacts the insects and that the size of the drop and the pressure allow coverage of the areas where the pest is located. Apply higher doses to serious infestations and when the foliage is dense. It can be applied throughout the entire crop cycle.

Incompatibility: It is not compatible with sodium hydroxide. When mixing with other products, a compatibility test must be performed before any application.


Contraindications: Do not apply in the middle of the day when the sun is strong, when it is raining or when wind could cause drift. Do not apply to plants that are stressed from lack of water, recent transplants, in germination beds, cuttings or plants with sensitive growth.



Order for Mixing the Product




Conditions for Application




- Do not apply during high temperatures and low relative humidity.
- Perform applications before 10 a.m. and after 4 p.m.

Application Sequence



- Alternative to prevent or delay risk of resistance.
- Does not share a common action point with other compounds and can be freely applied.
- Effective in sequences, alternations or mixtures with insecticides of different modes of action.

Coverage Density



- Use cone nozzles.
- Fine to medium droplet size of 136 µm at 218 µm (VMD).
- Ensure coverage density from 50 to 70 droplets/cm².
- Assure coverage on the tissues where the pest is located.



The information contained in this document is solely for your consultation, without legal value or guarantee. For more information related to the product, contact Tel. (777) 3211475 / 01 800 1127 500 to 502. ptigrupoultraquimia.com



For more information call
US Customer Service Toll Free Number:
1(800) 801-9657 in Mexico y (777) 319 35 45
55 7980 2287

Created:
Updated:
Version:
Page:

November 27, 2025.
November 27, 2025.
01.
1 of 1