



Biopesticide

Mode of Action Unknown or Uncertain

Contact, Ingestion and Inhalation



TECHNICAL SHEET – Plant Protection

Exemption from registration under EPA FIFRA 25(b)

Classification: Botanical Insecticide and Miticide.
Active Ingredient: Cinnamon Oil 4.00% wt.
Inert Ingredient : Water, Soap, and Xanthan gum 96.00% wt.
Formulation: Oil in Water Emulsion.

Instructions for use

Authorized for use in Safe Programs

Crop	Pest	Rate (fl.oz./Acre) *
Avocado (NL)	Red spider mite (<i>Oligonychus punicae</i>)	13 - 26 ²
Cotton (NL)	Aphid (<i>Aphis gossypii</i>) Red spider mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Whitefly (<i>Bemisia tabaci</i>)	26 - 51 ² 26 - 51 ² 26 - 51 ² 26 - 51 ²
Garlic, Onion, Chives, Asparagus (NL)	European Asparagus Aphid (<i>Brachycorynella asparagi</i>) Thrips (<i>Thrips tabaci</i>)	50 ¹ 32 - 64 ¹
Blueberry, Cranberry, Strawberry, Current, Blackberry (NL)	Red spider mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Spotted Wing Vinegar Fly (<i>Drosophila suzukii</i>) Green Peach Aphid (<i>Myzus persicae</i>) Mealybug (<i>Dysmicoccus vaccinii</i>)	43 - 64 ² 43 - 64 ² 32 - 64 ² 32 - 64 ² 14 - 30 ²
Eggplant, Chili, Pepper, Tomato, Potato, Bell pepper, Tobacco, Tomatillo (NL)	Aphids (<i>Aphis sp.</i> , <i>Macrosiphum sp.</i> , <i>Myzus persicae</i>) Mites (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Whitefly (<i>Bemisia tabaci</i>) Colorado Potato Beetle (<i>Leptinotarsa decemlineata</i>)	40 ³ 40 ³ 40 ³ 40 ³ 32 - 64 ²
Summer squash, Winter Squash, Chayote, Melon, Cucumber, Watermelon (NL)	Whitefly (<i>Bemisia tabaci</i>)	30 - 47 ²
Lime, Lemon, Mandarin, Orange, Tangerine, Grapefruit (NL)	Asian citrus psyllid (<i>Diaphorina citri</i>) Broad mite (<i>Polyphagotarsonemus latus</i>) Citrus mealybug (<i>Planococcus citri</i>)	26 - 77 ⁵ 26 - 77 ⁵ 26 - 77 ⁵
Grape (NL)	Aphid (<i>Aphis illinoisensis</i>)	22 - 78 ²
Almond, Pistachio (NL)	Navel Orangeworm (<i>Amyelois transitella</i>)	80 ⁴
Pome Fruit: Apple, Pear (NL)	Aphid (<i>Aphis pomi</i>) Two-Spotted Mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Psylla (<i>Cacopsylla bidens</i>) Whitefly (<i>Bemisia tabaci</i>)	32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴
Stone Fruit: Apricot, Cherry, Nectarine, Peach, Plum (NL)	Aphid (<i>Aphis pomi</i>) Two-Spotted Mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Psylla (<i>Cacopsylla bidens</i>) Whitefly (<i>Bemisia tabaci</i>)	32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴ 32 - 64 ⁴
Hops, Lentils (NL)	Aphids (<i>Aphis craccivora</i> , <i>Acyrtosiphon pisum</i>) Mites (<i>Tetranychus urticae</i>)	21 - 43 ⁴ 21 - 43 ⁴

* Volume of application 20 - 100 gallons/acre.

¹ Make 2 foliar applications, at 8-day intervals.

² Make 3 foliar applications at 7-day intervals.

Make 3 foliar applications at 7-day intervals, beginning 20 days after transplanting.

Make 3 applications to foliage at 7 to 14-day intervals. Use higher rate to higher infestation and when foliage is dense.

Make 5 foliar applications at 21-day intervals.

All ingredients in this product are exempt from the requirement of a tolerance under 40 CFR 180.

Its components act as an insecticide and a repellent against insects.

Re-entry Time: Four (4) hours after application.

Preharvest Interval: 0 days, No Limit (NL).


Interval between applications: 7, 8, and 21 days.

Method to prepare the product: Shake product before using. Condition the pH of the water to between 5.5 and 6.0. Pour the required quantity of product directly into the water. If mixing with other products always in the proper order of addition, until the mixture is homogenized.


Application of the product: Apply the solution the same day it is prepared. It is recommended that applications be performed when the first ovipositions are observed or when the first infestation is observed, preferably the first nymphal instars which are the most susceptible. Ensure that the product contacts the insects and that the size of the drop and the pressure allow coverage of the areas where the pest is located. Apply higher doses to serious infestations.

Incompatibility: It is not compatible with alkaline or oxidizing substances. When mixing with other products, a compatibility test must be performed prior to application. Do not mix with products whose compatibility is not completely tested.

Contraindications: Do not apply to plants that are stressed by inadequate irrigation, in extreme temperatures, when the sun is strong, when it is raining or when it is windy and can cause drift.




Order for Mixing the Product




- Make the mixture at time of application.
- Apply the spray solution the same day it is prepared.

Conditions for Application




- Do not apply during high temperatures and low relative humidity.
- Perform applications before 10 a.m. and after 4 p.m.

Application Sequence



- Alternative to prevent or delay risk of resistance.
- Does not share a common action point with other compounds and can be freely applied.
- Effective in sequences, alternations or mixtures with insecticides of different modes of action.

Coverage Density



- Use cone nozzles.
- Fine to medium droplet size of 136 µm at 218 µm (VMD).
- Ensure coverage density from 50 to 70 droplets/cm².
- Assure coverage on the tissues where the pest is located.



The information contained in this document is solely for your consultation, without legal value or guarantee. For more information related to the product, contact Tel. (777) 3211475 / 01 800 1127 500 to 502. ptigrupoultraquimia.com



For more information call
US Customer Service Toll Free Number:
1(800) 801-9657 in Mexico y (777) 319 35 45
55 7980 2287

Created:
Updated:
Version:
Page:

September 24, 2021.
March 21, 2024.
03.
1 of 1