

# Meromorphic Func.

## (1) Singularities:

① Def: i) Removable: If  $z_0$  is removable  
Then  $f(z)$  is bounded on  $U(z_0)$

ii) pole:  $z_0$  is a pole if  $\lim_{z \rightarrow z_0} |f(z)| = \infty$

iii) Essential:  $\lim_{z \rightarrow z_0} f(z)$  doesn't exist.

Then  $z_0$  is essential singularity.

## ② Properties:

i) Thm  $f \in \mathcal{O}(\mathcal{N}/\{z_0\})$ .  $f$  can be extended  
to  $\mathcal{N}$ .  $\Leftrightarrow \lim_{z \rightarrow z_0} (z - z_0) f(z) = 0$ .

Pf: ( $\Leftarrow$ ) Def  $g(z) = \begin{cases} (z - z_0) f(z), & z \in \mathcal{N}/\{z_0\} \\ 0, & z = z_0. \end{cases}$

Check  $g \in \mathcal{O}(\mathcal{N})$ .  $g'(z_0) = 0$ .

$\therefore g(z) = \sum_{n=2}^{\infty} (z - z_0)^n a_n$ . Expand at  $z_0$ .

Def  $\phi(z) = a_2 + a_3(z - z_0) + \dots$ .  $\phi|_{\mathcal{N}/\{z_0\}} = f$

( $\Rightarrow$ )  $\exists \phi \in \mathcal{O}(\mathcal{N})$ .  $\phi|_{\mathcal{N}/\{z_0\}} = f$ .

$\therefore \lim_{z \rightarrow z_0} f(z) = \phi(z_0) < \infty$ .



ii) (Weierstrass Thm)

$f \in \mathcal{O}(D(z_0, r) \setminus \{z_0\})$ , where  $z_0$  is essential singularity of  $f$ . Then  $f(D(z_0, r) \setminus \{z_0\})$  is dense.

Pf: By contradiction:

$$\exists m_0 \in \mathcal{K}(m_0), f(D(z_0, r) \setminus \{z_0\}) \cap \mathcal{K}(m_0) = \emptyset.$$

$$\text{Let } g(z) = \frac{1}{f - m_0} \in \mathcal{O}(D(z_0, r) \setminus \{z_0\}).$$

(2) Laurant Series:

Def:  $f$  is meromorphic on  $\mathcal{A}$  if  $f$  is holomorphic except several poles on  $\mathcal{A}$ .

Lemma. The poles of  $f$  are isolated.

Pf: If not,  $\exists \{z_k\} \rightarrow z_0$ .  $z_0$  is a pole.

Then  $\frac{1}{f}$  has an accumulation zero  $z_0$ .

Since  $\{z_k\} \cup \{z_0\} \subseteq \text{int}\{f \neq 0\}$ ,  $\therefore \frac{1}{f} \equiv 0$  or  $\infty$ ,  $\forall z \in \text{int}\{f \neq 0\}$ .

Since  $\frac{1}{f} \in \mathcal{O}(\text{int}\{f \neq 0\})$ , Contradict!


For  $z_0$  is an isolated singularity,  $z_0 \in C_r \subseteq C_R$ ,  $0 < r < R$ .

where  $f \in \mathcal{O}(C_R \setminus \{z_0\})$ . Then  $f(z) = \frac{1}{2\pi i} \int_{C_r \setminus C_1} \frac{f(s) ds}{s - z}$ ,  $z \in C_R \setminus C_r$ .

$$i) \int_{C_R} \frac{f(s)}{s - z} ds = \int_{C_R} \frac{1}{s - z_0} \frac{f(s) ds}{1 - \frac{z - z_0}{s - z_0}}$$

$$= \int_{C_R} \frac{1}{s - z_0} \sum_{n=0}^{\infty} \left( \frac{z - z_0}{s - z_0} \right)^n f(s) ds.$$

Since  $\left| \frac{z - z_0}{s - z_0} \right| < 1$ ,  $s \in \partial C_R$ .





$$\begin{aligned} \text{ii) } \int_{C_r} \frac{f(s) ds}{s-z} &= \int_{C_r} \frac{1}{z-z_0} \frac{f(s) ds}{1 - \frac{s-z_0}{z-z_0}} \\ &= \int_{C_r} \frac{1}{z-z_0} \sum_0^{\infty} \left( \frac{s-z_0}{z-z_0} \right)^n f(s) ds. \end{aligned}$$

$$\text{where } \left| \frac{s-z_0}{z-z_0} \right| < 1, s \in C_r$$

$$\therefore f(z) = \sum_{n \in \mathbb{Z}} a_n (z-z_0)^n, z \in C_r \setminus \{z_0\}.$$

$a_n$  is determined by above!

It's called Laurent Series.

Thm.

For Laurent series of  $f(z)$  at singularity

$z_0$ . We have following criteria:

i) If  $a_n = 0, \forall n < 0$ . Then  $z_0$  is removable

ii) If only finite  $n < 0$ , i.e.  $a_n \neq 0$ . Then  $z_0$  is a pole

iii) If there exists infinite  $n < 0$ , i.e.  $a_n \neq 0$ .

Then  $z_0$  is essential singularity.

Remark: The criteria holds only when  $z_0$  is a isolated singularity.

For  $z_0 = \infty$ . We can let  $g(z) = f\left(\frac{1}{z}\right)$

Expand  $g(z)$  at  $z=0$ . Replace " $n > 0$ "

with " $z < 0$ " in i), ii), iii).



### (3) Residue:

#### ① Zeros and poles:

$\mathcal{L}$  is connected

i) If  $f \in \mathcal{O}(\mathcal{L})$ ,  $f(z_0) = 0$ . Then  $\exists U(z_0)$  neighbourhood of  $z_0$  st.  $f(z) = (z - z_0)^n g(z)$ ,  $n \geq 1$ ,  $g \neq 0$  on  $U(z_0)$ .

ii) If  $z_0$  is a pole of  $f$  in  $\mathcal{L}$ . Then  $\exists U(z_0)$ .

st.  $f(z) = (z - z_0)^{-n} g(z)$ ,  $n \geq 1$ ,  $g \in \mathcal{O}(U(z_0))$ .

$$\therefore f(z) = \sum_{k=-n}^{\infty} a_k (z - z_0)^k, \quad \forall z \in U(z_0) \setminus \{z_0\}.$$

Note that  $\int_C f(z)/z^{n+1} dz = a_n$ ,  $z_0 \in \mathbb{C}$ .

We call it residue of  $f$  at  $z_0$ . Denote  $\text{Res}(f, z_0)$ .

Remark:  $\text{Res}(f, z_0) = \frac{1}{(n+1)!} \lim_{z \rightarrow z_0} \left( \frac{\partial}{\partial z} \right)^{n+1} ((z - z_0)^{n+1} f(z))$

#### ② Residue Formula:

i) Thm.  $f \in \mathcal{O}(\mathcal{L} \setminus \{z_i\}_1^N)$ ,  $C$  contour  $\{z_i\}_1^N \in \mathcal{L}$ .

Then  $\frac{1}{2\pi i} \int_C f(z) dz = \sum_{i=1}^N \text{Res}(f, z_i)$ , where  $\{z_i\}_1^N$  are poles of  $f$ .

ii) On  $\overline{\mathbb{C}_\infty}$ :

Consider the residue at  $z = \infty$ .

$$\text{Res}(f, \infty) = -\frac{1}{2\pi i} \oint_C f(z) dz, \quad C: |z| > r.$$

Where  $\infty$  is the isolated singularity in  $\mathbb{C}$ .



Let  $g(z) = f(\frac{1}{z})$ . Laurent series of  $g$  at 0 is:

$$g(z) = \sum a_n z^n. \therefore f(z) = \sum a_{-n} z^n.$$

$$\text{Res}(f, \infty) = -a_1$$

Remark: When  $z=\infty$  is removable,  $\text{Res}(f, \infty)$  may not be zero. e.g.  $\text{Res}(\frac{1}{z}, \infty) = -1$ .

Actually, we can expand  $f(z)$  at  $z=\infty$ .


$$\therefore f = \sum a_n z^n. \text{ Then } \text{Res}(f, \infty) = -a_1.$$

Thm.  $f$  defined on  $\bar{C}_\infty$  meromorphic. Then

$f$  has finite poles. Moreover, we have:

$$2\pi i \left( \sum_1^r \text{Res}(f, z_i) + \text{Res}(f, \infty) \right) = 0.$$

Pf: Note that  $\bar{C}_\infty \text{ cpt} \Rightarrow \text{reg cpt}$ .



The latter:

$$\int_{C-C} f(z) dz = 0.$$

prop.  $\text{Res}(f, \infty) = -\text{Res}\left(f\left(\frac{1}{z}\right) \frac{1}{z^2}, 0\right)$

Pf.  $\text{Res}(f, \infty) = \frac{1}{2\pi i} \oint_{C_r^-} f(z) dz$

$$= \frac{-1}{2\pi i} \int_0^{2\pi} f(re^{i\theta}) i r e^{i\theta} d\theta.$$

$$\stackrel{\theta \rightarrow -\theta}{=} \frac{-1}{2\pi i} \int_0^{2\pi} f\left(\frac{1}{\frac{1}{r} e^{i\theta}}\right) i \frac{1}{\frac{1}{r} e^{i\theta}} d\theta.$$

$$= -\frac{1}{2\pi i} \int_{C_{\frac{1}{r}}} f\left(\frac{1}{z}\right) \frac{1}{z^2} dz$$



### ③ Integration Calculate:

i) For  $\int_0^{2\pi} R(\cos\theta, \sin\theta) d\theta$ .  $R$  is a rational Func.:

If  $R(x, y) \neq \infty$  on  $x^2 + y^2 = 1$ . Let  $z = e^{i\theta}$ .

Then  $\begin{cases} \cos\theta = \frac{z^2+1}{2z} \\ \sin\theta = \frac{z^2-1}{2iz} \end{cases} \quad d\theta = \frac{dz}{iz}$ . We obtain:

$$\int_0^{2\pi} R d\theta = \oint_{|z|=1} R\left(\frac{z^2+1}{2z}, \frac{z^2-1}{2iz}\right) \frac{1}{iz} dz$$

$$= 2\pi i \sum_{|z_k| < 1} \text{Res}\left(R/iz, z_k\right)$$

ii) For  $\int_{-\infty}^{\infty} R(x) dx$ .  $R(x) = \frac{p(x)}{q(x)}$ , where  $p, q$  are polynomials, s.t.  $\deg q \geq \deg p + 2$ ,  $q \neq 0$ .


Lemma  $f$  conti,  $\lim_{R \rightarrow \infty} z f(z) = \lambda$ , when  $z$  is

on  $C_R = \{z = r e^{i\theta} \mid \theta_1 \leq \theta \leq \theta_2\}$ , uniformly with

$\theta$ . Then  $\lim_{R \rightarrow \infty} \int_{C_R} f(z) dz = i(\theta_2 - \theta_1)\lambda$ .

Pf:  $\exists R_0, \forall R > R_0, |z f(z) - \lambda| < \varepsilon$ .

$$\therefore \left| \int_{C_R} f(z) dz - i(\theta_2 - \theta_1)\lambda \right| \leq \frac{\varepsilon \ell}{R}. \quad \square$$



$$\oint_{C_{R+\varepsilon \cdot R}} R(z) dz = 2\pi i \sum_{\substack{|z_k| < R \\ \text{Im } z_k > 0}} \text{Res}(R, z_k)$$

$$= \int_{-R}^R R(x) dx + \int_{C_R} R dz.$$

Let  $R \rightarrow \infty$ . Since  $z f(z) \rightarrow \lambda$ .

$$\therefore \int_{-\infty}^{\infty} R(x) dx = 2\pi i \sum \text{Res}(R, z_k)$$

Remark: For  $\int_0^{\infty}$ , use The idea of

lemma is from:  $\int_{C_R} f(z) dz = \int_0^{\theta_2} i f(z) z d\theta$



$$= f(z_0) z_0 \cdot i(\theta_2 - \theta_1) \text{ mean value}$$

Thm of Integral.

iii) For  $\int_{-\infty}^{+\infty} R(x) e^{i\alpha x} dx$ ,  $\alpha > 0$ ,  $R = \frac{P}{Q}$ , where  $P(x), Q(x)$  are polynomials.  $\deg Q \geq \deg P + 1$  and  $Q(x) \neq 0 \forall x \in \mathbb{R}$ .

Jordan Lemma:

$g$  conti.  $\lim_{R \rightarrow \infty} g(Re^{i\theta}) = 0$ , uniformly with


$\theta \in [\theta_1, \theta_2]$ . Then  $\int_{C_R} g(z) e^{i\alpha z} dz \rightarrow 0$  ( $R \rightarrow \infty$ )

Pf: Denote  $M(R) = \max_{z \in C_R} |g(z)|$ .  $M(R) \rightarrow 0$  ( $R \rightarrow \infty$ )

$$|\int_{C_R} g(z) e^{i\alpha z} dz| \leq M(R) \int_{\theta_1}^{\theta_2} e^{-R \sin \theta} d\theta.$$

$$\leq R M(R) \int_0^{\frac{\pi}{2}} e^{-R \sin \theta} d\theta \leq C M(R) \rightarrow 0.$$

$$\text{by } \sin \theta \geq \frac{2}{\pi} \theta, \text{ when } 0 \leq \theta \leq \frac{\pi}{2}. \quad \square$$



$$\Rightarrow \int_{-\infty}^{+\infty} R(z) e^{i\alpha z} dz = 2\pi i \sum \text{Res}(Re^{i\alpha z}, z_k)$$

Since  $R(z) \rightarrow 0$  ( $R \rightarrow \infty$ )

iv) A special case:

when there's a pole on contour.

Lemma.  $f$  conti.  $C_r: |z-a|=r$ ,  $\theta_1 \leq \theta \leq \theta_2$ .

$\lim_{r \rightarrow a} (z-a) f(z) = \lambda$ , uniformly with  $\theta$ .

$z \in C_r$ . Then  $\lim_{r \rightarrow a} \int_{C_r} f(z) dz = i(\theta_2 - \theta_1) \lambda$


Pf:  $\exists r_0, \forall r < r_0, |(z-a)f(z) - \lambda| < \epsilon$ .

Similar Argument.



v) For  $\int_0^\infty \frac{R(x)}{x^\alpha} dx$ ,  $\alpha \in (1,1)$ ,  $R(x)$  is rational function.  $R = \frac{P}{Q}$

Case one:  $\alpha \in (0,1)$ . require  $\deg Q \geq \deg P + 1$ .



0 and  $\infty$  are pivot.

Choose  $\epsilon > 0$  to be partition line.

$$C_{r,\epsilon} = I_r \cup -I_\epsilon \cup [\epsilon, r] \cup [r, \epsilon].$$

$$\therefore \oint_{C_{r,\epsilon}} \frac{P(z)}{(z^\alpha)_0} dz = 2\pi i \sum \text{Res}(P/(z^\alpha)_0, z_k)$$

$$\text{where } (z^\alpha)_0 = e^{\alpha(\ln|z| + i\arg(z))}$$


$$\Rightarrow \text{LHS} = \int_{I_r} - \int_{I_\epsilon} + \int_\epsilon^r \frac{R(x)dx}{x^\alpha} + e^{-i\pi\alpha} \int_r^\epsilon \frac{R(x)dx}{x^\alpha}$$

Let  $\epsilon \rightarrow 0$ ,  $r \rightarrow \infty$ . Then  $\int_{I_r} - \int_{I_\epsilon} \rightarrow 0$ .

$$\therefore \int_0^\infty \frac{R(x)dx}{x^\alpha} + e^{-i2\pi\alpha} \int_\infty^0 \frac{R(x)dx}{x^\alpha} = 2\pi i \sum \text{Res}(P/(z^\alpha)_0, z_k)$$

Case two:  $\alpha \in (-1,0)$ . require:  $\deg Q \geq \deg P + 2$ .

Alternative method:




$$\text{Let } z = e^u.$$

$$\therefore \int R/x^\alpha = \int_{-\infty}^{+\infty} R(e^u) e^{u(1-\alpha)} du.$$

Remark: For multivalued Functions. The contour we choose should detour the pivots and partition lines:

e.g. For  $\frac{1}{x^\alpha(1+x)^\beta}$   
 $\alpha, \beta \in (0,1)$






vi) For  $\int_0^\infty R(x) (\ln x)^m dx$ ,  $m \geq 1$ .  $R(x)$  is rational function.  $R(x) = \frac{p}{a}$ .  $\deg p \geq \deg a + 2$ .  $a \neq 0$ .

For complexization:

Consider:  $F(z) = R(z) (\ln z)^{m+1}$ .



Choose  $\ln z = \ln |z| + i \arg(z)$ .

$$\therefore \oint_{C_m} F(z) dz = \int_{\Gamma_r} - \int_{\Gamma_R}$$

$$+ \int_1^r R(x) (\ln x)^{m+1} + \int_r^1 R(x) (\ln x + i2\pi)^{m+1}$$

$\Rightarrow$  Then  $R(x) (\ln x)^{m+1}$  will be affected.

Remark: We should calculate  $\int_0^\infty R(x) \ln x^k$ .

From  $k=1$  to  $m+1$  gradually.

Vii) Summary:

a. Check the integrand is a meromorphic function first.


That's why we substitute  $\cos \theta = \frac{z+\bar{z}}{2}$

$$\text{with } \frac{z^2 + \bar{z}^2}{2z} = \frac{z^2 + 1}{2z}$$


b. For  $\cos x / R(x)$  or  $\sin x / R(x)$ .

We only need to consider  $e^{iz} / R(x)$

Figure its real and imaginary part.



For integral contain  $e^z$ .



For rational function.



#### (4) Argument Principle:

##### ① Winding number:

Note that if  $\gamma \subseteq_{\text{open}} \mathbb{C}$ ,  $f \in \mathcal{O}(\gamma)$ .

$f$  nonvanishes and has no poles on  $\partial\gamma$ .

(\*)

→ If it holds,  
Then there may  
exist accumulation  
points on  $\partial\gamma$ .  
Then  $N$  or  $P$   
will be  $\infty$ !


$$\text{Then: } \frac{1}{2\pi i} \oint_{\partial\gamma} \frac{f'(z) dz}{f(z)} = N - P \stackrel{\Delta}{=} n(f(\partial\gamma), 0)$$

$N$  is total number of zeros.  $P$  is for poles

Pf: Easy to check by expansion of series.

Use contours to surround poles and zeros

##### Integration:



$$\oint_{\partial\gamma} \frac{f'(z) dz}{f(z)} = \oint \frac{dw}{w} = n(f(\partial\gamma), 0)$$

##### ② Application: Rouché's Thm:

$f, g \in \mathcal{O}(\gamma)$ ,  $C \subseteq_{\text{cycle}} \gamma$ . If  $|f| > |g|$  on  $C$ .


Then  $N_{f+g} = N_f$ .

Pf: Since  $f+g$  and  $f \in \mathcal{O}(\gamma)$ ,

$$\text{Then } \Delta_{\partial\gamma} \arg \left( \frac{f+g}{f} \right)$$

$$= \frac{1}{2\pi i} \oint_{\partial\gamma} \frac{(f+g/f)'}{f+g/f} dz = N_{f+g} - N_f.$$

Note that  $\Delta_{\partial\gamma} \arg(f+g/f) = \Delta_{\partial\gamma} \arg(1 + \frac{g}{f})$





since  $1 + \frac{z}{f} \neq 0, \forall z \in D. \therefore \Delta_{\partial D} \arg(1 + \frac{z}{f}) = 0.$

Other Form:

$f, g \in \mathcal{O}(D), \gamma \subseteq D, \text{ Jordan curve.}$

If  $|f| + |g| > |f + g|, \forall z \in \gamma.$

Then  $N_f = N_g.$

Pf: Note that  $f$  or  $g \neq 0$  on  $\gamma.$

$$\therefore 1 + \left| \frac{g}{f} \right| > \left| 1 + \frac{g}{f} \right|.$$

$$g/f \notin \mathbb{R}^+. \therefore \Delta_{\partial D} \arg\left(\frac{g}{f}\right) = 0.$$

since  $\frac{g}{f}(0)$  won't wind around  $z=0$  a whole circle.

because  $g/f$  won't walk through  $\mathbb{R}^+.$

(8) Application: Hurwitz Thm:

$f_n \in \mathcal{O}(D), f_n \neq 0, \forall n. f_n \xrightarrow{n.c.c} f$  on  $D.$

Then  $f \equiv 0$  or  $f \neq 0$  on  $D.$

Pf: Note that  $\forall K \subseteq_{cpt} D, \frac{f}{f_n} \xrightarrow{n} 1.$

$$\text{For } \varepsilon_0 > 0, \exists N_0, \forall n > N_0, \left| \frac{f}{f_n} - 1 \right| < \varepsilon_0.$$

$$\therefore N_f = N_{f-f_n+f_n} = \Delta_{\partial K} \arg(f-f_n+f_n)$$

$$= \Delta_{\partial K} \arg(f_n) + \Delta_{\partial K} \arg\left(\frac{f-f_n}{f_n} + 1\right)$$

for some  $n > N_0.$

$$\text{since } \left| 1 + \frac{f-f_n}{f_n} \right| > 1 - \varepsilon_0 > 0.$$

$$\therefore N_f = N_{f_n} = 0, \text{ when } f \neq 0.$$

When  $f \equiv 0$ , it holds automatically.

Remark: Analogously.

For  $f_n$  one-to-one,  $\mathcal{O}(D)$

$f_n \xrightarrow{n.c.c} f$ , then  $f \equiv c$

or  $f$  is one-to-one!



#### ④ Open mapping Thm:

If  $f \in \mathcal{O}(\mathcal{N})$ ,  $f \neq c$ , then  $f$  is open mapping

Pf:  $U \subseteq \mathcal{N}$   $\xrightarrow{f}$   $f(U) \ni w_0 = f(z_0)$

PROVE: the points surround  $w_0$  with " $\varepsilon$ " dist will have inverse image.

Note that  $f(z) - w = f(z) - w_0 + w_0 - w$

Choose  $\delta, \varepsilon$ :  $|f - w_0| > \varepsilon > |w_0 - w|$  when  $|z - z_0| = \delta$

Since:  $z_0$  is isolated zero of  $f - w_0 \in \mathcal{O}(\mathcal{N})$

#### Cor. (Maximal Modulus Principle)

$f \in \mathcal{O}(\mathcal{N})$ ,  $f \neq c$ , continuous on  $\partial \mathcal{N}$ . Then  $\max_{\bar{\mathcal{N}}} |f| = \max_{\partial \mathcal{N}} |f|$ .

Pf:  $f$  can't attain its maximal in  $\mathcal{N}^\circ$ .

Since it's an open mapping.

Remark: The minimal one won't hold if

$f$  has zeros, then  $\frac{1}{f} \notin \mathcal{O}(\mathcal{N})$ .

Cor.  $f \in \mathcal{O}(\mathcal{N})$ ,  $f \neq 0$  in  $\mathcal{N}$ . If  $f \equiv c$  on  $\partial \mathcal{N}$ .

Then  $f \equiv c$  on  $\bar{\mathcal{N}}$ .

Pf: If  $c = 0$ , it holds. Otherwise, let  $g = \frac{f(z)}{c}$

#### (5) $m$ to 1 Functions:

##### ① One to one:



Thm.  $f \in \mathcal{O}(D)$ , one-to-one. Then  $f' \neq 0$  on  $D$

Pf: WLOG. Suppose  $f(0) = 0$ . (By translation)

and  $f'(0) = 0$ . it will come into contrad.

Expand at  $z=0$ .  $\therefore f = \sum_m a_m z^m = z^m \varphi(z)$ ,  $m \geq 2$

Since  $z=0$  is isolated zero of  $f$ ,  $f'$

$\therefore \exists \bar{B}(0, \epsilon)$ .  $f \cdot f' \neq 0$  on  $\bar{B}(0, \epsilon)$

If  $a = \min_{\bar{B}} |f|$ . Then  $\forall w < \frac{a}{2}$

$N_{f-w} = N_f = 2$ . contradiction!

Remark: Converse is false. e.g.  $e^z$

But it will hold locally.

Thm.  $f \in \mathcal{O}(D)$ ,  $z_0 \in D$ . If  $f'(z_0) \neq 0$ . Then

locally near  $z_0$ ,  $f$  is one-to-one.

Pf: Expand  $f$  at  $z = z_0$ . then  $a_1 \neq 0$ :

$$f(z) = \sum_n a_n (z - z_0)^n$$

Estimate  $|f(z_1) - f(z_2)|$ , when  $s = |z_1 - z_2|$  is small enough.

Thm. (Darboux-Picard Thm)


$f \in \mathcal{O}(D)$ . conti on  $\partial D$ . If  $f$  is one-to-one on  $\partial D$ . Then  $f$  is one-to-one on  $\bar{D}$ .

( $\partial D$  is Jordan curve)



Pf: Note that  $f(z_0) \in \text{int } f(D) \Leftrightarrow z_0 \in \text{int } (D)$

by opening map Thm.



$\forall w_0 \in \text{int } f(D), \exists z_0 \in \text{int } D, \text{ st. } f(z_0) = w_0.$

$$\frac{1}{2\pi i} \oint_{\partial D} \frac{f'(z) dz}{f(z) - w_0} = \frac{1}{2\pi i} \oint_I \frac{dw}{w - w_0} = n(I, w_0) = 1$$

Since  $f$  is one-to-one on  $\partial f(D) \leftrightarrow \partial D$ . So when  $f$  walk around  $\partial D$  a circle, then  $f(\partial D)$  contour  $w_0$  a circle.

Remark: It holds when  $f$  is multicomplex  
(By proper map)

Def:  $f: D \rightarrow f(D)$  is biholomorphism on  $D$ .

When  $f \in \mathcal{O}(D)$ , one-to-one. Then:

$$\begin{aligned} f^{-1}: f(D) &\rightarrow D, (f^{-1}(z_0))' = \lim_{z \rightarrow z_0} \frac{f^{-1}(z) - f^{-1}(z_0)}{z - z_0} \\ &= \lim_{w \rightarrow w_0} \frac{w - w_0}{f(w) - f(w_0)} = \frac{1}{f'(w_0)}, w_0 = f^{-1}(z_0) \end{aligned}$$

Thm.  $f: \mathcal{R} \rightarrow f(\mathcal{R})$ , biholomorphism. If  $D \subseteq \mathcal{R}$  simply connected. Then  $f(D)$  is simply connected.

Pf: By Darboux - Picard Thm.



② m to one :

Thm.  $f \in \mathcal{O}(D)$ . Then  $f$  is a m-to-one covering map, locally.

Lemma.  $f \in \mathcal{O}(D)$ , nonvanishes on  $D$  simply connex.

Then exists  $g \in \mathcal{O}(D)$ , s.t.  $f = e^g$

$$\text{pf: Def } g(z) = \int_{z_0}^z \frac{f'}{f} dz + c_0 \in \mathcal{O}(D)$$

$$\therefore g'(z) = f'/f \Rightarrow (f e^{-g})' = 0$$

$$\therefore f = c e^g, \text{ let } e^{c_0} = f(z_0)$$

$$\therefore c=1, f(z) = e^{g(z)}, \forall z \in D.$$

$\Rightarrow$  Expand  $f$  at  $z = z_1$ ,  $f(z) = \sum a_n (z - z_1)^n$ .

$$\therefore f(z) = (z - z_1)^m \varphi(z), \varphi \text{ nonvanishes locally}$$

$$\therefore \varphi(z) = e^{g(z)}, \exists g \in \mathcal{O}(U(z_1)).$$

$$\therefore f(z) = ((z - z_1) e^{\frac{g(z)}{m}})^m, (m=0, \text{ holds initially suppose } m \geq 1)$$

(6) Entire Func and Meromorphic on  $\bar{\mathbb{C}}$  :

① Entire Func on  $\bar{\mathbb{C}}$  :

Note that entire function has the unique singularity :  $z = \infty$ . it must be a pole



or essential singularity (otherwise it will degenerate to const.)

Only when  $z=a$  is a pole, then  $f$  can be extended to  $\bar{\mathbb{C}}$ . (e.g.  $e^z$  can't)

$\Rightarrow f \in \mathcal{O}(\bar{\mathbb{C}})$ . Then  $f$  is a polynomial

(we call  $f \in \mathcal{O}(\mathbb{C})$ , but  $f$  isn't a polynomial by transcendental entire function)

properties:

i) Thm.  $f \in \mathcal{O}(\mathbb{C})$ .  $\lim_{z \rightarrow \infty} \frac{f(z)}{z^n} = 0$ . Then  $f$

is a polynomial with degree  $< n$

Pf:  $\exists R, \forall |z| > R, \left| \frac{f(z)}{z^n} \right| < \varepsilon$ .

By Cauchy Inequality.

Remark: It can be extended to  $\bar{\mathbb{C}}$ .

ii) Thm (Picard's little Thm)

$f \in \mathcal{O}(\mathbb{C})$ ,  $f \neq c$ . Then  $f$  takes

all points  $\in \mathbb{C}$  except one.

Remark: Picard's Great Thm:

$z_0$  is essential singularity of

$f(z)$ . Then  $\forall U(z_0)$  of  $z_0$ ,  $f(U(z_0))$

only misses at most one point.



iii)  $f \in \theta(\mathbb{C})$ .  $\forall z_0 \in \mathbb{C}$ , at least one coefficient is zero in its local expansion:  
 $\sum_0^{\infty} a_n (z - z_0)^n$ . Then  $f$  is polynomial.

Pf: When  $a_n = \frac{f^{(n)}(z_0)}{n!} = 0$ , then  $f^{(n)}(z_0) = 0$

If  $f$  isn't a polynomial.

Then  $f^{(n)}(z) \neq 0, \forall n$ .

For each  $n$ ,  $f^{(n)}(z) \in \theta(\mathbb{C})$ , has countable zeros<sup>(\*)</sup>

Denote  $Z_n = \{z_n^k \mid f^{(n)}(z_n^k) = 0\}$

$\therefore \bigcup_n Z_n$  is set of zeros of  $\{f^{(n)}(z)\}$ , countable.

But  $\forall n \subseteq \mathbb{C}$ ,  $|n| = 2^{\aleph_0} > |\bigcup_n Z_n| = \aleph_0$

Which is a contradiction!

Remark: (\*) : Since the zero is isolated,

which can correspond a neighborhood.

## ② Meromorphic on $\overline{\mathbb{C}}$ :

i) Thm.  $f$  is meromorphic on  $\overline{\mathbb{C}}$ . Then it's rational function.

Pf: Since  $\overline{\mathbb{C}}$  is cpt. So poles of  $f$  are finite.

Expand  $f$  at each pole  $z_k$ .

$$f(z) = f_k(z) + q_k\left(\frac{1}{z - z_k}\right), q_k \text{ is polynomial}$$

$$f\left(\frac{1}{z}\right) = f_n(z) + g_n(z),$$

$$\therefore f - g_n\left(\frac{1}{z}\right) - \sum q_k\left(\frac{1}{z - z_k}\right) \in \theta(\overline{\mathbb{C}}_n)$$



$$\therefore f(z) = g_n\left(\frac{1}{z}\right) + \sum g_k\left(\frac{1}{z-z_k}\right) + \text{Const.}$$

Remark: A meromorphic isn't polynomial will be called transcendental meromorphic func. e.g.

$$\frac{1}{e^{z+1}} \text{ has infinite poles: } (2k+1)\pi i \rightarrow \infty.$$

so it can't be extended to  $\overline{\mathbb{C}}$ .

ii) Thm.  $f$  is meromorphic on  $\overline{\mathbb{C}}$ ,  $f \neq \text{const.}$

The  $|f'(p)|$  is indep't with  $p$ .

$$\underline{pf:} \quad \forall p \in \overline{\mathbb{C}}, \exists z_0 \in \overline{\mathbb{C}}, \varepsilon > 0, \text{ s.t.}$$

$$f(z) - p \in \mathcal{O}(\overline{D}(z_0, \varepsilon)), \quad f(z) - p \neq 0 \text{ on } \overline{D}(z_0, \varepsilon)$$

$$\therefore \oint_{\partial D(z_0, \varepsilon)} \frac{f'(z) dz}{f(z) - p} = 0 = \oint_{\partial D(z_0, \varepsilon)} \frac{f'(z)}{f(z) - p} = N \cdot p.$$

$$\therefore N = |f'(p)| = p. \text{ poles of } f - p.$$

$$\therefore N_{f-p} = P_{f-p} = P_{f \text{ mer.}}$$