

Prova 3

① GSC para $L = \{a^i b^{2j} c^{3k} d^{4l} \mid i, j, k, l \geq 0\}$

$S \rightarrow aSBCD \mid \epsilon$

$S \rightarrow aBCD$

$DB \rightarrow BD$

$CB \rightarrow BC$

$DC \rightarrow CD$

$aB \rightarrow abb$

$bB \rightarrow bbb$

$bC \rightarrow bccc$

$cC \rightarrow cccc$

$cD \rightarrow cdddd$

$dD \rightarrow ddddd$

② GNC para AP, critério de parada-fim não, descreva as transições:

$S \rightarrow A0 \mid B1$

$A \rightarrow A00 \mid B10$

$B \rightarrow B11 \mid A01$

$\delta(q_0, \epsilon, S) \rightarrow (q_0, A0)$

$\delta(q_0, \epsilon, S) \rightarrow (q_0, B1)$

$\delta(q_0, \epsilon, A) \rightarrow (q_0, A00)$

$\delta(q_0, \epsilon, A) \rightarrow (q_0, B1)$

$\delta(q_0, \epsilon, A) \rightarrow (q_0, 0)$

$\delta(q_0, \epsilon, B) \rightarrow (q_0, B11)$

$\delta(q_0, \epsilon, B) \rightarrow (q_0, A0)$

$\delta(q_0, \epsilon, B) \rightarrow (q_0, 1)$

$\delta(q_0, 1, 1) \rightarrow (q_0, \epsilon)$

$\delta(q_0, 0, 0) \rightarrow (q_0, \epsilon)$

$(\epsilon, S, A0)$

$(\epsilon, S, B1)$

$(\epsilon, A, A00)$

$(\epsilon, A, B1)$

$(\epsilon, A, 0)$

$(\epsilon, B, B11)$

$(\epsilon, B, A0)$

$(\epsilon, B, 1)$

$(1, 1, \epsilon)$

$(0, 0, \epsilon)$

